

Tables of Calabi–Yau equations

Gert Almkvist, Christian van Enkevort,
Duco van Straten, and Wadim Zudilin

21 July 2005

Abstract

The main part of this paper is a big table (see Appendix A) containing what we believe to be a complete list of all fourth order equations of Calabi–Yau type known so far. In the text preceding the tables we explain what a differential equation of Calabi–Yau type is and we briefly discuss how we found these equations. We also describe an electronic version of this list.

Contents

1	Differential equations of Calabi–Yau type	1
2	Construction of Calabi–Yau equations	2
2.1	Computer searches	3
2.2	How to sum for $k = -n$ to $k = -1$	7
3	Electronic database of Calabi–Yau equations	9
A	Table of Calabi–Yau equations	10
B	Table of powers	93
C	Superseeker of Calabi–Yau differential equations	96

1 Differential equations of Calabi–Yau type

The differential equations we will be investigating are all of the following form

$$D_{n,k} := \sum_{i=0}^k z^i P_i(\theta). \quad (1)$$

Here $\theta := z \frac{\partial}{\partial z}$ and the $P_i(\theta)$ are polynomials with integral coefficients and order n . We will mostly restrict ourselves to equations of order 4, but the number of terms $k+1$ will vary. Independently of the number of terms we can rewrite the differential equation $D_{n,k}y = 0$ in the following way

$$y^{(n)} + a_{n-1}(z)y^{(n-1)} + \cdots + a_2(z)y'' + a_1(z)y' + a_0(z)y = 0,$$

where the $a_i(z)$ are rational functions of z . Recall that at any point $z = a$ (including $z = \infty$ via the transformation $w = z^{-1}$) we can define the so called indicial equation which governs the existence of solutions of the form $z^\lambda \times$ (power series in z). The set of solutions (with multiplicities) to the indicial equation at a certain point will be called the spectrum at that point.

In [1] the authors define Calabi–Yau equations to be differential equations of the form $D_{4,k}y = 0$ satisfying certain extra conditions. Let us briefly list the conditions satisfied by the equations from Appendix A. For an explanation of these conditions we refer to [1, 5]. Together they can be considered to define the notion of a Calabi–Yau equation.

Condition 1. The singular point $z = 0$ is a point of maximal unipotent monodromy, i.e., the indicial equation at $z = 0$ should have 0 as its only solution.

Condition 2. The coefficients $a_i(z)$ satisfy the following equation

$$a_1 = \frac{1}{2}a_2a_3 - \frac{1}{8}a_3^3 + a_2' - \frac{3}{4}a_3a_3' - \frac{1}{2}a_3''.$$

Condition 3. The solutions $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \lambda_4$ of the indicial equation at $z = \infty$ are positive rational numbers satisfying $\lambda_1 + \lambda_4 = \lambda_2 + \lambda_3 = s$ for some $s \in \mathbb{Q}$ (symmetry). We also suppose that the eigenvalues $e^{2\pi i\lambda}$ of the monodromy around $z = \infty$ are the zeros (counted with multiplicity) of a product of cyclotomic polynomials, which can be interpreted as the characteristic polynomial of the monodromy around $z = \infty$.

Condition 4. The power series solution near $z = 0$ has integral coefficients.

Condition 5. The genus zero instanton numbers computed by the standard recipe are integral (up to multiplication by an overall positive integer corresponding to the degree, which is not apparent from the differential equation alone).

2 Construction of Calabi–Yau equations

Some of the equations we collected in this paper have a geometric origin. These account for equations #1–28 except #9 in our list (see introduction to the main table for the precise references). Such geometric equations served as the starting point of our list. They are automatically of Calabi–Yau type because of their geometric origin. We observed that the expressions for the differential operators are very similar. This encouraged us to start looking for more equations that share the properties listed above. To find such equations we used various constructions:

- Translation to other singular points (not $z = 0$) with spectrum $\{a, a, a, a\}$
- Manipulation of binomial expressions for the coefficients of the power series solution
- Hadamard product
- ‘Pullbacks’ from fifth order equations
- Computer searches

The first method of obtaining new equations is rather trivial. Note that we defined a Calabi–Yau equation to have a point of maximal unipotent monodromy at $z = 0$. However, it can have other singular points where the spectrum consists of a single value a with multiplicity 4. These points can be found by computing the spectrum at all singular points. By translating to such a point (i.e., by substitution $z = z_0 + w$ if z_0 is the MUM-point or $z = 1/w$ if the MUM-point is at infinity) and simultaneously replacing y by $w^{-a}\tilde{y}$ we get a new equation for $\tilde{y}$, that has maximal unipotent monodromy at $w = 0$. Of course in some sense it is still the same equation. However, it can happen that one arrives at the same equation at a different point of maximal unipotent monodromy in entirely different ways. Then it may not always be clear that two equations are just translates of each other (in the sense described above). An example of a Picard–Fuchs equation with two points of maximal unipotent monodromy was discussed in [12, 13]. In that case both points have a geometric interpretation. By including translates it is easier to check if we include a certain equation.

The next three constructions are discussed in more detail in [1]. We will restrict ourselves to a few brief remarks. In many cases explicit formulas are known for the coefficients of the power series solution around $z = 0$. They satisfy a recursion relation coming from the differential equation. As can be seen from the list, in many cases the coefficients can be expressed in terms of sums or double sums of binomial expressions. There are many ways of transforming such expressions to other expressions which still satisfy a recursion relation. Translating this recursion relation back to a differential equation we obtain a new differential equation which in many cases turns out to be of Calabi–Yau type again.

The Hadamard product of two equations with power series solutions around $z = 0$ given by $\sum_{n=0}^{\infty} A_n z^n$ and $\sum_{n=0}^{\infty} B_n z^n$ is the differential equation that has $\sum_{n=0}^{\infty} A_n B_n z^n$ as its power series solution. As explained in [1] the Hadamard product can be used to construct higher order equations from lower order ones. The lower order ones come from [16, 6] and computer searches.

As discussed in [1] we can construct a fifth order equation from a fourth order one. This construction also goes the other way: given a suitable fifth order equation the corresponding fourth order equation can be constructed. The fifth order equations are found by other methods similar to the ones discussed above for fourth order equations.

2.1 Computer searches

A look at the list of differential equations shows that certain types occur very often. If we restrict to equations with $k = 1$, then the list reduces to the 14 hypergeometric cases. These are all of the form

$$D_{4,1}^{\text{hg}} = \theta^4 - cz(v\theta + u)(x\theta + w)(x\theta + x - w)(v\theta + v - u), \quad (2)$$

for integers c, u, v, w , and x . The term without z has to be $a\theta^4$ for some integer $a \neq 0$ because we want the equation to have maximal unipotent monodromy at $z = 0$. For some reason only $a = 1$ occurs. The last term determines the indicial equation at $z = \infty$, as can be seen by transforming (1) to $w = z^{-1}$.

$$D_{n,k}^w := \sum_{i=0}^k w^i P_{k-i}(-\theta_w), \quad (3)$$

where $\theta_w := w \frac{\partial}{\partial w}$. Therefore the indicial equation at $z = \infty$ is

$$P_k(-\lambda) = 0.$$

Hence Condition 3 tells us among other things that $P_k(-\lambda)$ has rational roots. Another part of this condition is the symmetry $\lambda_1 + \lambda_4 = \lambda_2 + \lambda_3 = s$ for some $s \in \mathbb{Q}$. Take $s = 1$ and write $\lambda_1 = \frac{u}{v}$ and $\lambda_2 = \frac{w}{x}$, then we find the expression from (2).

However, we still have not fully exploited Condition 3. The characteristic polynomial has to be a product of cyclotomic polynomials. As it has fixed degree 4, there are only finitely many possibilities (see Table 1). The λ_i are defined by the property that $e^{2\pi i \lambda_1}, \dots, e^{2\pi i \lambda_4}$ is the full set of zeros of the characteristic polynomial. This determines the λ_i up to integer shifts. Because they also have to be positive and we have the symmetry condition $\lambda_1 + \lambda_4 = \lambda_2 + \lambda_3 = s$, there are only finitely many choices for the λ_i 's for a fixed characteristic polynomial and fixed s . As mentioned above, for the hypergeometric equations s turns out to be 1. We listed the 14 possible spectra in Table 1. These correspond exactly to the first 14 equations in our list. Given the spectrum at $z = \infty$ the only thing that is not fixed is the constant c . This constant can easily be found by requiring the coefficients of the power series solution around $z = 0$ to be integral.

One step up from the hypergeometric equations are the Calabi–Yau equations with $k = 2$. In this case the most general form is

$$\begin{aligned} D_{4,2}^{\text{gen}} = & \theta^4 - cz(A\theta^4 + 2A\theta^3 + (B+A)\theta^2 + B\theta + C) \\ & - dz^2(v\theta + u)(x\theta + w)(x\theta + 2x - w)(v\theta + 2v - u). \end{aligned} \quad (4)$$

The first and the last term are again restricted because of the required monodromies (Conditions 1 and 3), where again we suppose that the coefficient of θ^4 is 1. The only difference is that for the spectrum at $z = \infty$ we now use $s = 2$ instead of $s = 1$. This is just based on the experimental fact that for $k = 2$ most Calabi–Yau equations turn out to have $s = 2$. In fact our database contains one exception ($s = \frac{3}{2}$). For larger k we observe that several values of s occur (based on the equations in our database).

Condition 2 forces the middle term (with z) to be of the given form, as can be checked by computing the coefficients $a_i(z)$ for a general polynomial $P_1(\theta)$. In practice in most cases the middle term factors and the operator is of the form

$$\begin{aligned} D_{4,2}^{\text{fact}} = & \theta^4 - cz(v\theta + u)(v\theta + v - u)(A\theta^2 + A\theta + B) \\ & - dz^2(v\theta + u)(x\theta + w)(x\theta + 2x - w)(v\theta + 2v - u). \end{aligned} \quad (5)$$

Using this form reduces the number of parameters by one which makes the computer search a lot faster.

As mentioned above, we also search for equations of lower order ($n = 2$ or $n = 3$) because they can be used as factors in the Hadamard product to find interesting fourth order equations. We have searched for operators of the following forms (see [1])

$$D_{2,2}^{\text{had}} = \theta^2 - c(A\theta^2 + A\theta + B) - d(\theta + 1)^2$$

and

$$D_{3,2}^{\text{had}} = \theta^3 - c(2\theta + 1)(A\theta^2 + A\theta + B) - d(\theta + 1)^3.$$

char. polyn.	spectra ($s = 1$)	spectra ($s = 2$)
$\phi_1\phi_2\phi_3$	—	—
$\phi_1\phi_2\phi_6$	—	—
$\phi_1\phi_2\phi_6$	—	—
$\phi_1\phi_2^3$	—	—
$\phi_1^2\phi_2^2$	—	$\{\frac{1}{2}, 1, 1, \frac{3}{2}\}^*$
$\phi_1^2\phi_3$	—	$\{\frac{2}{3}, 1, 1, \frac{4}{3}\}^*, \{\frac{1}{3}, 1, 1, \frac{5}{3}\}^*$
$\phi_1^2\phi_4$	—	$\{\frac{3}{4}, 1, 1, \frac{5}{4}\}, \{\frac{1}{4}, 1, 1, \frac{7}{4}\}^*$
$\phi_1^2\phi_6$	—	$\{\frac{5}{6}, 1, 1, \frac{7}{6}\}^*, \{\frac{1}{6}, 1, 1, \frac{11}{6}\}^*$
$\phi_1^3\phi_2$	—	—
ϕ_1^4	—	$\{1, 1, 1, 1\}^*$
$\phi_2^2\phi_3$	$\{\frac{1}{3}, \frac{1}{2}, \frac{1}{2}, \frac{2}{3}\}$	$\{\frac{1}{2}, \frac{2}{3}, \frac{4}{3}, \frac{3}{2}\}, \{\frac{1}{3}, \frac{1}{2}, \frac{3}{2}, \frac{5}{3}\}$
$\phi_2^2\phi_4$	$\{\frac{1}{4}, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}\}$	$\{\frac{1}{2}, \frac{3}{4}, \frac{5}{4}, \frac{3}{2}\}, \{\frac{1}{4}, \frac{1}{2}, \frac{3}{2}, \frac{7}{4}\}$
$\phi_2^2\phi_6$	$\{\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, \frac{5}{6}\}$	$\{\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{11}{6}\}, \{\frac{1}{2}, \frac{5}{6}, \frac{7}{6}, \frac{3}{2}\},$
ϕ_2^4	$\{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\}$	$\{\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}\}$
$\phi_3\phi_4$	$\{\frac{1}{4}, \frac{1}{3}, \frac{2}{3}, \frac{3}{4}\}$	$\{\frac{2}{3}, \frac{3}{4}, \frac{5}{4}, \frac{4}{3}\}, \{\frac{1}{4}, \frac{1}{3}, \frac{5}{3}, \frac{7}{4}\}, \{\frac{1}{3}, \frac{3}{4}, \frac{5}{4}, \frac{5}{3}\},$ $\{\frac{1}{4}, \frac{2}{3}, \frac{4}{3}, \frac{7}{4}\}$
$\phi_3\phi_6$	$\{\frac{1}{6}, \frac{1}{3}, \frac{2}{3}, \frac{5}{6}\}$	$\{\frac{2}{3}, \frac{5}{6}, \frac{7}{6}, \frac{4}{3}\}, \{\frac{1}{6}, \frac{1}{3}, \frac{5}{3}, \frac{11}{6}\}, \{\frac{1}{3}, \frac{5}{6}, \frac{7}{6}, \frac{5}{3}\},$ $\{\frac{1}{6}, \frac{2}{3}, \frac{4}{3}, \frac{11}{6}\}$
ϕ_3^2	$\{\frac{1}{3}, \frac{1}{3}, \frac{2}{3}, \frac{2}{3}\}$	$\{\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \frac{5}{3}\}, \{\frac{2}{3}, \frac{2}{3}, \frac{4}{3}, \frac{4}{3}\}, \{\frac{1}{3}, \frac{1}{3}, \frac{5}{3}, \frac{5}{3}\}$
$\phi_4\phi_6$	$\{\frac{1}{6}, \frac{1}{4}, \frac{3}{4}, \frac{5}{6}\}$	$\{\frac{1}{6}, \frac{3}{4}, \frac{5}{4}, \frac{11}{6}\}, \{\frac{3}{4}, \frac{5}{6}, \frac{7}{6}, \frac{5}{4}\}, \{\frac{1}{4}, \frac{5}{6}, \frac{7}{6}, \frac{7}{4}\},$ $\{\frac{1}{6}, \frac{1}{4}, \frac{7}{4}, \frac{11}{6}\}$
ϕ_4^2	$\{\frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}\}$	$\{\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, \frac{7}{4}\}, \{\frac{3}{4}, \frac{3}{4}, \frac{5}{4}, \frac{5}{4}\}, \{\frac{1}{4}, \frac{1}{4}, \frac{7}{4}, \frac{7}{4}\}$
ϕ_5	$\{\frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}\}$	$\{\frac{2}{5}, \frac{4}{5}, \frac{6}{5}, \frac{8}{5}\}, \{\frac{3}{5}, \frac{4}{5}, \frac{6}{5}, \frac{7}{5}\}, \{\frac{1}{5}, \frac{3}{5}, \frac{7}{5}, \frac{9}{5}\},$ $\{\frac{1}{5}, \frac{2}{5}, \frac{8}{5}, \frac{9}{5}\}$
ϕ_6^2	$\{\frac{1}{6}, \frac{1}{6}, \frac{5}{6}, \frac{5}{6}\}$	$\{\frac{1}{6}, \frac{5}{6}, \frac{7}{6}, \frac{11}{6}\}, \{\frac{1}{6}, \frac{1}{6}, \frac{11}{6}, \frac{11}{6}\}, \{\frac{5}{6}, \frac{5}{6}, \frac{7}{6}, \frac{7}{6}\}$
ϕ_8	$\{\frac{1}{8}, \frac{3}{8}, \frac{5}{8}, \frac{7}{8}\}$	$\{\frac{1}{8}, \frac{3}{8}, \frac{13}{8}, \frac{15}{8}\}, \{\frac{1}{8}, \frac{5}{8}, \frac{11}{8}, \frac{15}{8}\},$ $\{\frac{3}{8}, \frac{7}{8}, \frac{9}{8}, \frac{13}{8}\}, \{\frac{5}{8}, \frac{7}{8}, \frac{9}{8}, \frac{11}{8}\}$
ϕ_{10}	$\{\frac{1}{10}, \frac{3}{10}, \frac{7}{10}, \frac{9}{10}\}$	$\{\frac{7}{10}, \frac{9}{10}, \frac{11}{10}, \frac{13}{10}\}, \{\frac{1}{10}, \frac{3}{10}, \frac{17}{10}, \frac{19}{10}\},$ $\{\frac{1}{10}, \frac{7}{10}, \frac{13}{10}, \frac{19}{10}\}, \{\frac{3}{10}, \frac{9}{10}, \frac{11}{10}, \frac{17}{10}\}$
ϕ_{12}	$\{\frac{1}{12}, \frac{5}{12}, \frac{7}{12}, \frac{11}{12}\}$	$\{\frac{1}{12}, \frac{5}{12}, \frac{19}{12}, \frac{23}{12}\}, \{\frac{1}{12}, \frac{7}{12}, \frac{17}{12}, \frac{23}{12}\},$ $\{\frac{5}{12}, \frac{11}{12}, \frac{13}{12}, \frac{19}{12}\}, \{\frac{7}{12}, \frac{11}{12}, \frac{13}{12}, \frac{17}{12}\}$

Table 1: Possible characteristic polynomials and spectra

The special forms of the operator discussed above allow us to do a computer search for Calabi–Yau equations. For fourth order equations we use the following two step process.

Step 1. A program written in C searches a certain range of the parameters in the expressions for the operators discussed above (e.g., $A, B, C, c, d, \frac{u}{v}$, and $\frac{w}{x}$ for $D_{4,2}^{\text{gen}}$). The C program only checks for an integral solution to the recursion relation. If an integral power series solution exists, the corresponding values of the parameters are written to a Maple array.

Step 2. A Maple program reads the candidate parameter combinations and computes the instanton numbers. These need not be integral, because we can still multiply with the degree. Therefore, we compute the lowest common multiple of denominators of the first 15 and the first 20 instanton numbers. If they are equal, we consider this to be acceptable. In practice we observe that in these cases the lowest common multiple is also very small (below 100). We also check if the differential operator factors as a differential operator (using the `DEtools` package). If does, we remove the operator from our list.

For second and third order equations we cannot directly define the instanton numbers. Instead, we check if using them as factors in a Hadamard product yields a fourth order equation of Calabi–Yau type.

The first step is the most time consuming one. So we use a few tricks to save some time. First note that replacing z by λz multiplies the coefficients A_n of the power series solution by λ^n . In terms of the parameters this amounts to multiplying c and d by λ and λ^2 respectively. Therefore we can put $c = 1$ and compute the first N coefficients of the power series. In general the coefficients will be rational. We will require the denominators to be a product of powers of a limited number of small primes. We can then compute the scaling factor to make the first N coefficients integral. Using the rescaled parameters c and d we can compute the first M ($M > N$) coefficients and check that they are integral. We always used $N = 50$ and $M = 1000$, but it never happened that with the rescaled parameters we found non integral coefficients. Note that we do not *prove* that all the coefficients of the power series solution are integral, but as we check the first 1000 coefficients, we are pretty confident that they are.

In this way we did extensive searches using up to a dozen computers. The results are summarized in Table 2 on the following page. The last column contains the number of solutions found by the C program. For the fourth order equations we also indicate between parentheses the number of solutions that survive the second step of the searching process and are contained in our database. For the spectra searched we refer to Table 1 on the page before. The star indicates that we restrict to the subset of the spectra from Table 1 labelled by a star.

The difference between the programs `search4` and `search4q` is that the former is completely general, whereas the latter assumes that apart from $z = 0$ and $z = \infty$ there is just one rational critical point. This yields big restrictions on the parameters and allows us to compute d . Therefore this program is much faster and we were able to search a much larger parameter range. Of course it also means that we cannot find any equations which do not satisfy this extra condition.

We do not claim that our search has been exhaustive. First of all the parameter ranges are rather arbitrary. It is to be expected that enlarging these ranges

Program	Operator	Parameter range	Spectra	#
search2	$D_{2,2}^{\text{had}}$	$1 \leq A, B \leq 500, d = 2^p 3^q 5^r 7^s$ with $0 \leq p \leq 20, 0 \leq q \leq 6,$ $0 \leq r \leq 5, 0 \leq s \leq 2$	—	126
search3	$D_{3,2}^{\text{had}}$	$1 \leq A, B \leq 500, d = 2^p 3^q 5^r 7^s$ with $0 \leq p \leq 20, 0 \leq q \leq 6,$ $0 \leq r \leq 5, 0 \leq s \leq 2$	—	139
search4	$D_{4,2}^{\text{gen}}$	$1 \leq A, B, C \leq 200, d = 2^p 3^q 5^r 7^s$ with $0 \leq p \leq 15, 0 \leq q \leq 5,$ $0 \leq r \leq 4, 0 \leq s \leq 2$	$s = 2^*$	66 (15)
search4f	$D_{4,2}^{\text{fact}}$	$1 \leq A, B \leq 100, d = 2^p 3^q 5^r 7^s$ with $0 \leq p \leq 15, 0 \leq q \leq 6,$ $0 \leq r \leq 5, 0 \leq s \leq 1$	$s = 2$	364 (58)
search4q	$D_{4,2}^{\text{gen}}$	$1 \leq A, B, C \leq 1000, d$ is com- puted	$s = 2$	1321 (30)

Table 2: Computer searches

we will find more equations. Furthermore, the expressions (4) and (5) that we use are not completely general. Both the assumption that the coefficient of θ^4 is 1 and the restriction to $s = 2$ are arbitrary. As mentioned above, we already have one example with $s = \frac{3}{2}$. There are also 32 equations where the coefficient of θ^4 is bigger than 1. However, in all these cases k is at least 4. All in all, this is still very much work in progress. We are still adding equations to our list every now and then and we keep enhancing the information on the equations in our database.

2.2 How to sum for $k = -n$ to $k = -1$

As already mentioned, the work [1] explains manipulating binomial expressions to deduce new examples of Calabi–Yau equations. This is certainly a “hypergeometric” machinery, but, surprisingly, it works for certain unusual series $y_0 = \sum_{n=0}^{\infty} A_n z^n$ as well, namely, when coefficients A_n involve not only binomials but also harmonic sums and consist of several summations.

In some formulas for A_n (particular cases are #211 and #264) it is necessary to sum also for $k = -n$ to $k = -1$ to get the correct coefficients of the solution y_0 to the differential equation. This requires expressing binomial coefficients near $k = -\bar{k}$, where $\bar{k}$ is a positive integer:

$$\begin{aligned}
\binom{n}{k} &\approx \binom{n}{-\bar{k} - \varepsilon} = \frac{n!}{\Gamma(1 - \bar{k} - \varepsilon)\Gamma(1 + n + \bar{k} + \varepsilon)} \\
&= (-1)^{\bar{k}} \bar{k}^{-1} \binom{n + \bar{k}}{n}^{-1} \varepsilon + O(\varepsilon^2), \\
\binom{2k}{k} &\approx \binom{-2\bar{k} - 2\varepsilon}{-\bar{k} - \varepsilon} = \frac{\Gamma(1 - 2\bar{k} - 2\varepsilon)}{\Gamma(1 - \bar{k} - \varepsilon)^2} = \bar{k}^{-1} \binom{2\bar{k}}{\bar{k}}^{-1} \varepsilon + O(\varepsilon^2),
\end{aligned}$$

$$\binom{n+k}{n} \approx \binom{n-\bar{k}-\varepsilon}{n} = \frac{\Gamma(1+n-\bar{k}-\varepsilon)}{n!\Gamma(1-\bar{k}-\varepsilon)} = (-1)^{\bar{k}} \bar{k}^{-1} \left(\frac{n}{\bar{k}}\right)^{-1} \varepsilon + O(\varepsilon^2),$$

and so on.

The differential equation for case #211 was found in two ways: first, as the reflection of case #210 at infinity and, secondly, by taking Maple's **Zeilberger** on the following binomial sum:

$$“A_n” = \left(\frac{2n}{n}\right)^4 \sum_k (-1)^{n+k} \binom{n}{k}^2 \binom{2k}{k} \binom{4n-2k}{2n-k} \binom{n+k}{n}^{-2} \left(\frac{2n}{k}\right)^{-1}.$$

Usual summing for $k = 0$ to $k = n$ does not give the correct coefficient of y_0 . To remove the latter quotation one should sum for $k = -n$ to n . So we have to consider also negative k . The 'difficult' (i.e., negative) part of the summand is

$$\binom{n}{k}^2 \binom{2k}{k} \binom{n+k}{n}^{-2} \left(\frac{2n}{k}\right)^{-1},$$

which near $k = -\bar{k}$ becomes (after simple reduction)

$$(-1)^{\bar{k}} \binom{n+\bar{k}}{n}^{-2} \left(\frac{2\bar{k}}{\bar{k}}\right)^{-1} \binom{n}{\bar{k}}^2 \left(\frac{2n+\bar{k}}{2n}\right) + O(\varepsilon),$$

thus giving us the formula indicated below in Table A.

In case #264 we use **Zeilberger** on

$$\begin{aligned} “A_n” &= 16^{-n} \left(\frac{2n}{n}\right)^2 \sum_k (n-2k) \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2n+2k}{n+k}^2 \\ &\quad \times \left(\frac{4n-2k}{2n-k}\right)^2 \left(\frac{2n}{k}\right)^{-1} \left(\frac{2n}{n-k}\right)^{-1} \end{aligned}$$

and proceed, as before, for negative k to obtain for the 'difficult' part

$$\binom{n}{k} \binom{2k}{k} \left(\frac{2n}{k}\right)^{-1}$$

near $k = -\bar{k}$ the following result:

$$\bar{k}^{-1} \binom{n+\bar{k}}{n}^{-1} \left(\frac{2\bar{k}}{\bar{k}}\right)^{-1} \left(\frac{2n+\bar{k}}{2n}\right) \varepsilon + O(\varepsilon^2).$$

Therefore, for negative k , we get the sum (this formula was found by C. Krattenthaler [9])

$$\begin{aligned} \frac{d“A_n”}{d\varepsilon} &= 16^{-n} \left(\frac{2n}{n}\right)^2 \sum_{k=1}^n \frac{n+2k}{k} \binom{2n+k}{2n} \binom{2n+2k}{n+k} \\ &\quad \times \left(\frac{2n-2k}{n-k}\right)^2 \left(\frac{4n+2k}{2n+k}\right)^2 \binom{n+k}{n}^{-1} \left(\frac{2k}{k}\right)^{-1}, \end{aligned}$$

where we replace $\bar{k} = -k$ by k . Finally, adding the derivative of “ A_n ” for positive k , we deduce our formula in Table A for case #264.

3 Electronic database of Calabi–Yau equations

All equations we found are also available in electronic form. The web interface to our database is at:

`http://enriques.mathematik.uni-mainz.de/enckevort/db`

The database includes some extra information that is not contained in the tables in this paper. Furthermore we hope to be able to keep it up to date so that it can serve as a list of all known Calabi–Yau equations. Therefore we encourage anybody who finds new equations of Calabi–Yau type to send them to us so that we can include them into our database. Comments and suggestions are also welcome. The database and the web interface are still a work in progress.

Information about the use of the database can be found at the web address mentioned above. A small warning for people comparing the table in this article and the database: for equations past #180 the numbering of the equations in the database is different from the one used in this article. However, by searching for `almkvist[n]` in the source field, where n is the number used in this article one can easily find the equation in the database.

Part of the extra information that we do not provide in this article, but is contained in the database, is related to numerical computations of the monodromies around the singular points of the differential equations. In many cases we succeeded in guessing the exact monodromies using the approximate monodromies obtained numerically. In turn the exact monodromies allowed us to compute some geometric invariants of potential underlying Calabi–Yau manifolds. These, in combination with further information obtained from the monodromies, enabled us to compute elliptic invariants in many cases. This work is discussed in detail in [5].

Remark. We feel that our table in Appendix A is not perfectly organized, and we plan to systemize it in the nearest future. On the other hand, the papers [1], [5], as well as the electronic database have crucial references to the absolute numbering in Appendix A. Therefore, we have decided to keep the present numeration at least in the first arXiv-version of this work.

Acknowledgments

We would like to thank frankly C. Krattenthaler who has provided [9] us with several explicit formulas for A_n appearing now in Table A. The work of the fourth-named author is partially supported by grant no. 03-01-00359 of the Russian Foundation for Basic Research.

A Table of Calabi–Yau equations

The first column is reserved for numeration of cases. The second column of the following table contains a 4th-order linear differential operator D (by means of $\theta = z \frac{d}{dz}$). The third column indicates coefficients $\{A_n\}_{n=0,1,2,\dots}$ of the analytic solution $y_0(z) = \sum_{n=0}^{\infty} A_n z^n$ to the MUM differential equation $Dy = 0$ normalized by the condition $y_0(0) = A_0 = 1$.

See [11], [7] for cases #1, #2; [10] for cases #3–6; [7] cases #7, #8; [8] for cases #10–14; [3] for cases #15–23, and [4] for cases #24–28.

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$	
1	$D = \theta^4 - 5^5 z \left(\theta + \frac{1}{5}\right) \left(\theta + \frac{2}{5}\right) \left(\theta + \frac{3}{5}\right) \left(\theta + \frac{4}{5}\right)$	$A_n = \frac{(5n)!}{n!^5}$
2	$D = \theta^4 - 8 \cdot 10^5 z \left(\theta + \frac{1}{10}\right) \left(\theta + \frac{3}{10}\right) \left(\theta + \frac{7}{10}\right) \left(\theta + \frac{9}{10}\right)$	$A_n = \frac{(10n)!}{n!^3 (2n)! (5n)!}$
3	$D = \theta^4 - 256 z \left(\theta + \frac{1}{2}\right)^4$	$A_n = \binom{2n}{n}^4$
4	$D = \theta^4 - 3^6 z \left(\theta + \frac{1}{3}\right)^2 \left(\theta + \frac{2}{3}\right)^2$	$A_n = \left(\frac{(3n)!}{n!^3}\right)^2$
5	$D = \theta^4 - 432 z \left(\theta + \frac{1}{2}\right)^2 \left(\theta + \frac{1}{3}\right) \left(\theta + \frac{2}{3}\right)$	$A_n = \binom{2n}{n}^2 \frac{(3n)!}{n!^3}$
6	$D = \theta^4 - 2^{10} z \left(\theta + \frac{1}{2}\right)^2 \left(\theta + \frac{1}{4}\right) \left(\theta + \frac{3}{4}\right)$	$A_n = \binom{2n}{n} \frac{(4n)!}{n!^4}$
7	$D = \theta^4 - 2^{16} z \left(\theta + \frac{1}{8}\right) \left(\theta + \frac{3}{8}\right) \left(\theta + \frac{5}{8}\right) \left(\theta + \frac{7}{8}\right)$	$A_n = \frac{(8n)!}{n!^4 (4n)!}$
8	$D = \theta^4 - 11664 z \left(\theta + \frac{1}{6}\right) \left(\theta + \frac{1}{3}\right) \left(\theta + \frac{2}{3}\right) \left(\theta + \frac{5}{6}\right)$	$A_n = \frac{(6n)!}{n!^4 (2n)!}$
9	$D = \theta^4 - 12^6 z \left(\theta + \frac{1}{12}\right) \left(\theta + \frac{5}{12}\right) \left(\theta + \frac{7}{12}\right) \left(\theta + \frac{11}{12}\right)$	$A_n = \binom{2n}{n} \frac{(12n)!}{n!^2 (4n)! (6n)!}$
10	$D = \theta^4 - 2^{12} z \left(\theta + \frac{1}{4}\right)^2 \left(\theta + \frac{3}{4}\right)^2$	$A_n = \left(\frac{(4n)!}{n!^2 (2n)!}\right)^2$
11	$D = \theta^4 - 12^3 z \left(\theta + \frac{1}{4}\right) \left(\theta + \frac{3}{4}\right) \left(\theta + \frac{1}{3}\right) \left(\theta + \frac{2}{3}\right)$	$A_n = \binom{3n}{n} \frac{(4n)!}{n!^4}$
12	$D = \theta^4 - 2^{10} \cdot 3^3 z \left(\theta + \frac{1}{4}\right) \left(\theta + \frac{3}{4}\right) \left(\theta + \frac{1}{6}\right) \left(\theta + \frac{5}{6}\right)$	$A_n = \binom{4n}{n} \frac{(6n)!}{n!^2 (2n)!^2}$
13	$D = \theta^4 - 2^8 \cdot 3^6 z \left(\theta + \frac{1}{6}\right)^2 \left(\theta + \frac{5}{6}\right)^2$	$A_n = \left(\frac{(6n)!}{n! (2n)! (3n)!}\right)^2$
14	$D = \theta^4 - 2^8 \cdot 3^3 z \left(\theta + \frac{1}{2}\right)^2 \left(\theta + \frac{1}{6}\right) \left(\theta + \frac{5}{6}\right)$	$A_n = \binom{2n}{n} \frac{(6n)!}{n!^3 (3n)!}$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
15	$D = \theta^4 - 3z(3\theta + 1)(3\theta + 2)(7\theta^2 + 7\theta + 2) - 72z^2(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$
	$A_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{n}{k}^3$
16	$D = \theta^4 - 4z(2\theta + 1)^2(5\theta^2 + 5\theta + 2) + 2^8 z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_{j+k+l+m=n} \left(\frac{n!}{j!k!l!m!} \right)^2 = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k}$
17	$D = 25\theta^4 - 15z(51\theta^4 + 84\theta^3 + 72\theta^2 + 30\theta + 5) + 6z^2(531\theta^4 + 828\theta^3 + 541\theta^2 + 155\theta + 15) - 54z^3(423\theta^4 + 2160\theta^3 + 4399\theta^2 + 3795\theta + 1170) + 3^5 z^4(279\theta^4 + 1368\theta^3 + 2270\theta^2 + 1586\theta + 402) - 3^{10} z^5(\theta + 1)^4$
	$A_n = \sum_{j+k+l=n} \left(\frac{n!}{j!k!l!} \right)^3$
18	$D = \theta^4 - 4z(2\theta + 1)^2(3\theta^2 + 3\theta + 1) - 4z^2(4\theta + 2)(4\theta + 3)(4\theta + 5)(4\theta + 6)$
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^4$
19	$D = 23^2\theta^4 - 23z(921\theta^4 + 2046\theta^3 + 1644\theta^2 + 621\theta + 92) - z^2(380851\theta^4 + 1328584\theta^3 + 1772673\theta^2 + 1033528\theta + 221168) - 2z^3(475861\theta^4 + 1310172\theta^3 + 1028791\theta^2 + 208932\theta - 27232) - 2^2 \cdot 17z^4(8873\theta^4 + 14020\theta^3 + 5139\theta^2 - 1664\theta - 976) + 2^3 \cdot 3 \cdot 17z^5(\theta + 1)^2(3\theta + 2)(3\theta + 4)$
	$A_n = \sum_{k=0}^n \binom{n}{k}^3 \binom{n+k}{n} \binom{2n-k}{n}$
20	$D = \theta^4 - 3z(48\theta^4 + 60\theta^3 + 53\theta^2 + 23\theta + 4) + 9z^2(873\theta^4 + 1980\theta^3 + 2319\theta^2 + 1344\theta + 304) - 2 \cdot 3^4 z^3(1269\theta^4 + 3888\theta^3 + 5259\theta^2 + 3348\theta + 800) + 2^2 \cdot 3^6 z^4(891\theta^4 + 3240\theta^3 + 4653\theta^2 + 2952\theta + 688) - 2^3 \cdot 3^{11} z^5(\theta + 1)^2(3\theta + 2)(3\theta + 4)$
	$A_n = \sum_{k=0}^n \binom{n}{k} \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
21	$ \begin{aligned} D &= 25\theta^4 - 20z(36\theta^4 + 84\theta^3 + 72\theta^2 + 30\theta + 5) \\ &\quad - 2^4 z^2(181\theta^4 + 268\theta^3 + 71\theta^2 - 70\theta - 35) \\ &\quad + 2^8 z^3(\theta + 1)(37\theta^3 + 248\theta^2 + 375\theta + 165) \\ &\quad + 2^{10} z^4(39\theta^4 + 198\theta^3 + 331\theta^2 + 232\theta + 59) + 2^{15} z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^3 \binom{2k}{k} \binom{2n-2k}{n-k}$
22	$ \begin{aligned} D &= 49\theta^4 - 7z(155\theta^4 + 286\theta^3 + 234\theta^2 + 91\theta + 14) \\ &\quad - z^2(16105\theta^4 + 68044\theta^3 + 102261\theta^2 + 66094\theta + 15736) \\ &\quad + 2^3 z^3(2625\theta^4 + 8589\theta^3 + 9071\theta^2 + 3759\theta + 476) \\ &\quad - 2^4 z^4(465\theta^4 + 1266\theta^3 + 1439\theta^2 + 806\theta + 184) + 2^9 z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^5$
23	$ \begin{aligned} D &= 9\theta^4 - 12z(64\theta^4 + 80\theta^3 + 73\theta^2 + 33\theta + 6) \\ &\quad + 2^7 z^2(194\theta^4 + 440\theta^3 + 527\theta^2 + 315\theta + 75) \\ &\quad - 2^{12} z^3(94\theta^4 + 288\theta^3 + 397\theta^2 + 261\theta + 66) \\ &\quad + 2^{17} z^4(22\theta^4 + 80\theta^3 + 117\theta^2 + 77\theta + 19) - 2^{23} z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$
24	$ \begin{aligned} D &= \theta^4 - 3z(3\theta + 2)(3\theta + 1)(11\theta^2 + 11\theta + 3) \\ &\quad - 9z^2(3\theta + 5)(3\theta + 2)(3\theta + 4)(3\theta + 1) \end{aligned} $
	$A_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}$
25	$D = \theta^4 - 4z(2\theta + 1)^2(11\theta^2 + 11\theta + 3) - 16z^2(2\theta + 3)^2(1 + 2\theta)^2$
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}$
26	$ \begin{aligned} D &= \theta^4 - 2z(2\theta + 1)^2(13\theta^2 + 13\theta + 4) \\ &\quad - 12z^2(2\theta + 3)(2\theta + 1)(3\theta + 2)(3\theta + 4) \end{aligned} $
	$ \begin{aligned} A_n &= \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{n+k}{k} \binom{2k}{n} \\ &= \binom{2n}{n} \sum_{k,l} \binom{n}{k}^2 \binom{n}{l}^2 \binom{k+l}{n} \end{aligned} $

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
27	$ \begin{aligned} D = & 9\theta^4 - 3z(173\theta^4 + 340\theta^3 + 272\theta^2 + 102\theta + 15) \\ & - 2z^2(1129\theta^4 + 5032\theta^3 + 7597\theta^2 + 4773\theta + 1083) \\ & + 2z^3(843\theta^4 + 2628\theta^3 + 2353\theta^2 + 675\theta + 6) \\ & - z^4(295\theta^4 + 608\theta^3 + 478\theta^2 + 174\theta + 26) + z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k,l} \binom{n}{k}^2 \binom{n}{l}^2 \binom{k+l}{n} \binom{2n-k}{n}$
28	$ \begin{aligned} D = & \theta^4 - z(65\theta^4 + 130\theta^3 + 105\theta^2 + 40\theta + 6) \\ & + 4z^2(4\theta + 3)(4\theta + 5)(\theta + 1)^2 \end{aligned} $
	$A_n = \sum_{k,l,m} \binom{n}{k} \binom{n}{l} \binom{m}{k} \binom{m}{l} \binom{k+l}{k} \binom{n}{m}^2 = \sum_{k,l} \binom{n}{k}^2 \binom{n}{l}^2 \binom{k+l}{n}^2$
29	$ \begin{aligned} D = & \theta^4 - 2z(2\theta + 1)^2(17\theta^2 + 17\theta + 5) \\ & + 4z^2(2\theta + 3)(2\theta + 1)(\theta + 1)^2 \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$

Remark. The original formulas for A_n in cases #24, 25, 27–29 were simplified with the help of the binomial identity

$$\sum_j \binom{n}{j} \binom{k}{j} = \binom{n+k}{k}.$$

The origin of the following new cases is explained in the introductory part and [1, 5]. In particular, cases #30 and #31 are obtained in [1], Section 6; case #32 (the pullback of the differential equation related to $\zeta(4)$) is the subject of [1], Section 4; case #130 was communicated to us by H. Verrill [14] and case #186 by Tjøtta [13].

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
2^*	$D = \theta^4 - 2^4 \cdot 5z(2\theta + 1)^2(50\theta^2 + 50\theta + 33) + 2^{10} \cdot 5^4 z^2(2\theta + 1)(2\theta + 3)(5\theta + 4)(5\theta + 6)$
	$A_n = 2000^n \binom{2n}{n} \sum_k \binom{-1/10}{k} \binom{-3/10}{k} \binom{-7/10}{n-k} \binom{-9/10}{n-k}$
	the instanton numbers lie in $\mathbb{Z}[\sqrt{5}]$
3^*	$D = \theta^4 - 2^4 z(2\theta + 1)^2(2\theta^2 + 2\theta + 1) + 2^{10} z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = 16^n \binom{2n}{n} \sum_k \binom{-1/2}{k}^2 \binom{-1/2}{n-k}^2 = \binom{2n}{n} \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$
4^*	$D = \theta^4 - 6z(2\theta + 1)^2(9\theta^2 + 9\theta + 5) + 2^2 \cdot 3^6 z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = 27^n \binom{2n}{n} \sum_k \binom{-1/3}{k}^2 \binom{-2/3}{n-k}^2$
4^{**}	$D = \theta^4 - 6z(2\theta + 1)^2(9\theta^2 + 9\theta + 4) + 2^2 \cdot 3^4 z^2(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4)$
	$A_n = 27^n \binom{2n}{n} \sum_k \binom{-1/3}{k} \binom{-2/3}{k} \binom{-1/3}{n-k} \binom{-2/3}{n-k}$ $= \binom{2n}{n} \sum_{k=0}^n \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3}$
6^*	$D = \theta^4 - 2^5 z(2\theta + 1)^2(2\theta^2 + 2\theta + 1) + 2^8 z^2(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$
	$A_n = 32^n \binom{2n}{n} \sum_k \binom{-1/2}{k} \binom{-1/4}{k} \binom{-1/2}{n-k} \binom{-3/4}{n-k}$
7^*	$D = \theta^4 - 2^4 z(2\theta + 1)^2(32\theta^2 + 32\theta + 19) + 2^{14} z^2(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$
	$A_n = 256^n \binom{2n}{n} \sum_k \binom{-1/8}{k} \binom{-3/8}{k} \binom{-5/8}{n-k} \binom{-7/8}{n-k}$
7^{**}	$D = \theta^4 - 2^4 z(2\theta + 1)^2(32\theta^2 + 32\theta + 13) + 2^{16} z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = 256^n \binom{2n}{n} \sum_k \binom{-1/8}{k} \binom{-5/8}{k} \binom{-3/8}{n-k} \binom{-7/8}{n-k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
8^*	$D = \theta^4 - 12z(2\theta + 1)^2(18\theta^2 + 18\theta + 11) + 2^4 \cdot 3^4 z^2(2\theta + 1)(2\theta + 3)(6\theta + 5)(6\theta + 7)$
	$A_n = 108^n \binom{2n}{n} \sum_k \binom{-1/6}{k} \binom{-1/3}{k} \binom{-5/6}{n-k} \binom{-2/3}{n-k}$
8^{**}	$D = \theta^4 - 12z(2\theta + 1)^2(18\theta^2 + 18\theta + 7) + 2^4 \cdot 3^6 z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = 108^n \binom{2n}{n} \sum_k \binom{-1/6}{k} \binom{-2/3}{k} \binom{-5/6}{n-k} \binom{-1/3}{n-k}$
9^*	$D = \theta^4 - 48z(2\theta + 1)^2(72\theta^2 + 72\theta + 41) + 2^{14} \cdot 3^4 z^2(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4)$
	$A_n = 1728^n \binom{2n}{n} \sum_k \binom{-1/12}{k} \binom{-5/12}{k} \binom{-7/12}{n-k} \binom{-11/12}{n-k}$
9^{**}	$D = \theta^4 - 48z(2\theta + 1)^2(72\theta^2 + 72\theta + 31) + 2^{12} \cdot 3^6 z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = 432^n \binom{2n}{n}^2 \sum_k (-1)^k \binom{-5/6}{k} \binom{-1/6}{n-k}^2$
10^*	$D = \theta^4 - 2^4 z(2\theta + 1)^2(8\theta^2 + 8\theta + 5) + 2^{14} z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = 64^n \binom{2n}{n} \sum_k \binom{-1/4}{k}^2 \binom{-3/4}{n-k}^2$
10^{**}	$D = \theta^4 - 2^4 z(2\theta + 1)^2(8\theta^2 + 8\theta + 3) + 2^{12} z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = 64^n \binom{2n}{n} \sum_k \binom{-1/4}{k} \binom{-3/4}{k} \binom{-3/4}{n-k} \binom{-1/4}{n-k}$ $= \binom{2n}{n} \sum_{k=0}^n \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
13*	$D = \theta^4 - 48z(2\theta + 1)^2(18\theta^2 + 18\theta + 13) + 2^{10} \cdot 3^6 z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = 432^n \binom{2n}{n} \sum_k \binom{-1/6}{k}^2 \binom{-5/6}{n-k}^2$
13**	$D = \theta^4 - 48z(2\theta + 1)^2(18\theta^2 + 18\theta + 5) + 2^{10} \cdot 3^4 z^2(2\theta + 1)(2\theta + 3)(3\theta + 1)(3\theta + 5)$
	$A_n = 432^n \binom{2n}{n} \sum_k \binom{-1/6}{k} \binom{-5/6}{k} \binom{-5/6}{n-k} \binom{-1/6}{n-k}$ $= \binom{2n}{n} \sum_{k=0}^n \frac{(6k)!}{k!(2k)!(3k)!} \frac{(6n-6k)!}{(n-k)!(2n-k)!(3n-3k)!}$
14*	$D = \theta^4 - 2^4 \cdot 3^2 z(2\theta + 1)^2(2\theta^2 + 2\theta + 1) + 2^{10} \cdot 3^2 z^2(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4)$
	$A_n = 144^n \binom{2n}{n} \sum_k \binom{-1/6}{k} \binom{-1/2}{k} \binom{-5/6}{n-k} \binom{-1/2}{n-k}$
	the instanton numbers lie in $\mathbb{Z}[\sqrt{3}]$
$\hat{1}$	$D = \theta^4 - 5z(10000\theta^4 + 12500\theta^3 + 9500\theta^2 + 3250\theta + 399) + 5^8 z^2(2400\theta^4 + 6000\theta^3 + 6290\theta^2 + 2800\theta + 399) - 4 \cdot 5^{14} z^3(4\theta + 3)(80\theta^3 + 240\theta^2 + 221\theta + 42) + 5^{20} z^4(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 10z(2\theta + 1)(3125\theta^4 + 6250\theta^3 + 7875\theta^2 + 4750\theta + 1226) + 2^4 5^6 z^2(\theta + 1)(6250\theta^4 + 25000\theta^3 + 4550\theta^2 + 41000\theta + 14851) - 2^6 5^{14} z^3(\theta + 1)(\theta + 2)(2\theta + 3)(25\theta^2 + 75\theta + 82) + 2^8 5^{19} z^4(\theta + 1)(\theta + 2)(\theta + 3)(25\theta^2 + 100\theta + 113) - 2^9 5^{25} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = (4 \cdot 5^5)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/5)_k (2/5)_k (3/5)_k (4/5)_k}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
$\widehat{2}$	$D = \theta^4 - 2^4 5 z (160000\theta^4 + 200000\theta^3 + 154000\theta^2 + 54000\theta + 7189)$ $+ 2^{14} 5^8 z^2 (9600\theta^4 + 24000\theta^3 + 25320\theta^2 + 1400\theta + 1669)$ $- 2^{24} 5^{14} z^3 (4\theta + 3)(320\theta^3 + 960\theta^2 + 888\theta + 171)$ $+ 2^{32} 5^{20} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5$ $- 2^5 5 z (2\theta + 1)(50000\theta^4 + 100000\theta^3 + 127000\theta^2 + 77000\theta + 19811)$ $+ 2^{16} 5^6 z^2 (\theta + 1)(100000\theta^4 + 400000\theta^3 + 731000\theta^2 + 662000\theta + 240811)$ $- 2^{28} 5^{14} z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(100\theta^2 + 300\theta + 331)$ $+ 2^{39} 5^{19} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(50\theta^2 + 200\theta + 227)$ $- 2^{49} 5^{25} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 3200000^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/10)_k (3/10)_k (7/10)_k (9/10)_k}{k!^5}$
$\widehat{3}$	$D = \theta^4 - 2^4 z (4\theta + 1)(64\theta^3 + 64\theta^2 + 44\theta + 9)$ $+ 2^{14} z^2 (4\theta + 1)(96\theta^3 + 216\theta^2 + 196\theta + 61)$ $- 2^{24} z^3 (4\theta + 1)(4\theta + 3)(16\theta^2 + 44\theta + 33)$ $+ 2^{32} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 z (2\theta + 1)(80\theta^4 + 160\theta^3 + 200\theta^2 + 120\theta + 31)$ $+ 5 \cdot 2^{16} z^2 (\theta + 1)(32\theta^4 + 128\theta^3 + 232\theta^2 + 208\theta + 75)$ $- 5 \cdot 2^{28} z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(4\theta^2 + 12\theta + 13)$ $+ 5 \cdot 2^{37} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(2\theta^2 + 8\theta + 9)$ $- 2^{49} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 1024^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k^5}{k!^5}$
$\widehat{4}$	$D = \theta^4 - 3^2 z (1296\theta^4 + 1620\theta^3 + 1224\theta^2 + 414\theta + 49)$ $+ 3^8 z^2 (7776\theta^4 + 19440\theta^3 + 20322\theta^2 + 9000\theta + 1267)$ $- 4 \cdot 3^{16} z^3 (4\theta + 3)(144\theta^3 + 432\theta^2 + 397\theta + 75)$ $+ 3^{24} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2 \cdot 3^2 z (2\theta + 1)(405\theta^4 + 810\theta^3 + 1017\theta^2 + 612\theta + 158)$ $+ 2^4 3^8 z^2 (\theta + 1)(810\theta^4 + 3240\theta^3 + 5886\theta^2 + 5292\theta + 1913)$ $- 2^6 3^{17} z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(15\theta^2 + 45\theta + 49)$ $+ 2^8 3^{22} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(45\theta^2 + 180\theta + 203)$ $- 2^9 3^{30} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = (4 \cdot 3^6)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/3)_k^2 (2/3)_k^2}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
$\widehat{5}$	$D = \theta^4 - 2^2 3z(576\theta^4 + 720\theta^3 + 542\theta^2 + 182\theta + 21)$ $+ 2^4 3^2 z^2(124416\theta^4 + 311040\theta^3 + 324576\theta^2 + 143280\theta + 20017)$ $- 2^{11} 3^7 z^3(4\theta + 3)(1152\theta^3 + 3456\theta^2 + 3172\theta + 597)$ $+ 2^{16} 3^{12} z^4(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^3 3z(2\theta + 1)(180\theta^4 + 360\theta^3 + 451\theta^2 + 271\theta + 70)$ $+ 2^{10} 3^4 z^2(\theta + 1)(360\theta^4 + 1440\theta^3 + 2613\theta^2 + 2346\theta + 847)$ $- 2^{15} 3^8 z^3(\theta + 1)(\theta + 2)(2\theta + 3)(120\theta^2 + 360\theta + 391)$ $+ 2^{22} 3^{10} z^4(\theta + 1)(\theta + 2)(\theta + 3)(180\theta^2 + 720\theta + 811)$ $- 2^{29} 3^{15} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 1728^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k^3 (1/3)_k (2/3)_k}{k!^5}$
$\widehat{6}$	$D = \theta^4 - 2^4 z(1024\theta^4 + 1280\theta^3 + 968\theta^2 + 328\theta + 39)$ $+ 2^{12} z^2(24576\theta^4 + 61440\theta^3 + 64256\theta^2 + 28480\theta + 4017)$ $- 2^{27} z^3(4\theta + 3)(512\theta^3 + 1536\theta^2 + 1412\theta + 267)$ $+ 2^{40} z^4(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 z(2\theta + 1)(320\theta^4 + 640\theta^3 + 804\theta^2 + 484\theta + 125)$ $+ 2^{18} z^2(\theta + 1)(640\theta^4 + 2560\theta^3 + 4652\theta^2 + 4184\theta + 1513)$ $- 2^{31} z^3(\theta + 1)(\theta + 2)(2\theta + 3)(160\theta^2 + 480\theta + 523)$ $+ 2^{44} z^4(\theta + 1)(\theta + 2)(\theta + 3)(80\theta^2 + 320\theta + 361)$ $- 2^{54} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 2^{12n} \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k^3 (1/4)_k (3/4)_k}{k!^5}$
$\widehat{7}$	$D = \theta^4 - 2^4 z(49152\theta^4 + 81920\theta^3 + 62720\theta^2 + 21760\theta + 2793)$ $+ 2^{26} z^2(4\theta + 3)(768\theta^3 + 1984\theta^2 + 1572\theta + 291)$ $- 2^{46} z^3(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 z(2\theta + 1)(20480\theta^4 + 40960\theta^3 + 51840\theta^2 + 31360\theta + 8087)$ $+ 2^{24} z^2(\theta + 1)(40960\theta^4 + 163840\theta^3 + 298880\theta^2 + 270080\theta + 98071)$ $- 2^{48} z^3(\theta + 1)(\theta + 2)(2\theta + 3)(64\theta^2 + 192\theta + 211)$ $+ 2^{67} z^4(\theta + 1)(\theta + 2)(\theta + 3)(32\theta^2 + 128\theta + 145)$ $- 2^{89} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 2^{18n} \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/8)_k (3/8)_k (5/8)_k (7/8)_k}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
$\widehat{8}$	$D = \theta^4 - 2^2 3^2 z(3888\theta^4 + 6480\theta^3 + 4950\theta^2 + 1710\theta + 217)$ $+ 2^4 3^{10} z^2(4\theta + 3)(1728\theta^3 + 4464\theta^2 + 3532\theta + 651)$ $- 2^{10} 3^{18} z^3(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^3 3^2 z(1620\theta^4 + 3240\theta^3 + 4095\theta^2 + 2475\theta + 638)$ $+ 2^{10} 3^8 z^2(\theta + 1)(3240\theta^4 + 12960\theta^3 + 23625\theta^2 + 21350\theta + 7739)$ $- 5 \cdot 2^{15} 3^{17} z^3(\theta + 1)(\theta + 2)(2\theta + 3)(24\theta^2 + 72\theta + 79)$ $+ 5 \cdot 2^{22} 3^{22} z^4(\theta + 1)(\theta + 2)(\theta + 3)(36\theta^2 + 144\theta + 163)$ $- 2^{29} 3^{30} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = (4 \cdot 11664)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/3)_k (2/3)_k (1/6)_k (5/6)_k}{k!^5}$
$\widehat{9}$	$D = \theta^4 - 2^4 3^2 z(248832\theta^4 + 41472\theta^3 + 318528\theta^2 + 11116\theta + 14497)$ $+ 2^{22} 3^{10} z^2(4\theta + 3)(432\theta^3 + 1116\theta^2 + 886\theta + 165)$ $- 2^{34} 3^{18} z^3(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 3^2 z(2\theta + 1)(103680\theta^4 + 207360\theta^3 + 262944\theta^2$ $+ 159264\theta + 41087)$ $+ 2^{20} 3^{17} z^2(\theta + 1)(207360\theta^4 + 829440\theta^3 + 1514592\theta^2$ $+ 1370304\theta + 498143)$ $- 2^{38} 3^{17} z^3(\theta + 1)(\theta + 2)(2\theta + 3)(240\theta^2 + 720\theta + 793)$ $+ 2^{53} 3^{22} z^4(\theta + 1)(\theta + 2)(\theta + 3)(360\theta^2 + 1440\theta + 1633)$ $- 2^{69} 3^{30} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 3456^{2n} \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/12)_k (5/12)_k (7/12)_k (11/12)_k}{k!^5}$
$\widehat{10}$	$D = \theta^4 - 2^4 z(3072\theta^4 + 5120\theta^3 + 3904\theta^2 + 1344\theta + 169)$ $+ 2^{23} z^2(4\theta + 3)(24\theta^3 + 62\theta^2 + 49\theta + 9)$ $- 2^{34} z^3(4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 z(2\theta + 1)(1280\theta^4 + 2560\theta^3 + 3232\theta^2 + 1952\theta + 503)$ $+ 2^{20} z^2(\theta + 1)(2560\theta^4 + 10240\theta^3 + 18656\theta^2 + 16832\theta + 6103)$ $- 2^{38} z^3(\theta + 1)(\theta + 2)(2\theta + 3)(80\theta^2 + 240\theta + 263)$ $+ 2^{53} z^4(\theta + 1)(\theta + 2)(\theta + 3)(40\theta^2 + 160\theta + 181)$ $- 2^{69} z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = 2^{14n} \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/4)_k^2 (3/4)_k^2}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
$\widehat{11}$	$D = \theta^4 - 2^2 3 z (2304\theta^4 + 2880\theta^3 + 2186\theta^2 + 746\theta + 91)$ $+ 2^4 3^3 z^2 (1990656\theta^4 + 4976640\theta^3 + 5213952\theta^2 + 2318400\theta + 329497)$ $- 2^{15} 3^7 z^3 (4\theta + 3)(4608\theta^3 + 13824\theta^2 + 12724\theta + 2415)$ $+ 2^{24} 3^{12} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^3 3 z (2\theta + 1)(720\theta^4 + 1440\theta^3 + 1813\theta^2 + 1093\theta + 282)$ $+ 2^{12} 3^5 z^2 (\theta + 1)(480\theta^4 + 1920\theta^3 + 3493\theta^2 + 3146\theta + 1139)$ $- 2^{19} 3^8 z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(480\theta^2 + 1440\theta + 1573)$ $+ 2^{28} 3^{10} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(720\theta^2 + 2880\theta + 3253)$ $- 2^{39} 3^{15} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = (4 \cdot 12^3)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/3)_k (2/3)_k (1/4)_k (3/4)_k}{k!^5}$
$\widehat{12}$	$D = \theta^4 - 2^4 3 z (9216\theta^4 + 11520\theta^3 + 8840\theta^2 + 3080\theta + 403)$ $+ 2^{12} 3^2 z^2 (1990656\theta^4 + 4976640\theta^3 + 5241600\theta^2 + 2352960\theta + 3420049)$ $- 2^{27} 3^7 z^3 (4\theta + 3)(4608\theta^3 + 13824\theta^2 + 12772\theta + 2451)$ $+ 2^{40} 3^{12} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 z (2\theta + 1)(2880\theta^4 + 5760\theta^3 + 7300\theta^2 + 4420\theta + 1137)$ $+ 2^{18} 3^5 z^2 (\theta + 1)(1920\theta^4 + 7680\theta^3 + 14020\theta^2 + 12680\theta + 4607)$ $- 5 \cdot 2^{31} 3^8 z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(96\theta^2 + 288\theta + 317)$ $+ 5 \cdot 2^{44} 3^{10} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(144\theta^2 + 576\theta + 653)$ $- 2^{59} 3^{15} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = (2^{12} 3^3)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/4)_k (3/4)_k (1/6)_k (5/6)_k}{k!^5}$
$\widehat{13}$	$D = \theta^4 - 2^4 3^2 z (20736\theta^4 + 25920\theta^3 + 20016\theta^2 + 7056\theta + 961)$ $+ 2^{14} 3^8 z^2 (31104\theta^4 + 77760\theta^3 + 82152\theta^2 + 37080\theta + 5461)$ $- 2^{24} 3^{16} z^3 (4\theta + 3)(576\theta^3 + 1728\theta^2 + 1600\theta + 309)$ $+ 2^{32} 3^{24} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 3^2 z (2\theta + 1)(6480\theta^4 + 12960\theta^3 + 16488\theta^2 + 10008\theta + 2567)$ $+ 2^{16} 3^8 z^2 (\theta + 1)(12960\theta^4 + 51840\theta^3 + 94824\theta^2 + 85968\theta + 31295)$ $- 2^{28} 3^{17} z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(60\theta^2 + 180\theta + 199)$ $+ 2^{39} 3^{22} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(90\theta^2 + 360\theta + 409)$ $- 2^{49} 3^{30} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$
	$A'_n = (2^{10} 3^6)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k (1/6)_k^2 (5/6)_k^2}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
$\widehat{14}$	$ \begin{aligned} D = & \theta^4 - 2^4 3 z (2304 \theta^4 + 2880 \theta^3 + 2192 \theta^2 + 752 \theta + 93) \\ & + 2^{14} 3^4 z^2 (31104 \theta^4 + 77760 \theta^3 + 81576 \theta^2 + 36360 \theta + 5197) \\ & - 2^{24} 3^7 z^3 (4\theta + 3)(576 \theta^3 + 1728 \theta^2 + 1592 \theta + 303) \\ & + 2^{32} 3^{12} z^4 (4\theta + 1)(4\theta + 3)(4\theta + 7)(4\theta + 9) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2^5 3 z (2\theta + 1)(720 \theta^4 + 1440 \theta^3 + 1816 \theta^2 + 1096 \theta + 283) \\ & + 2^{16} 3^4 z^2 (\theta + 1)(1440 \theta^4 + 5760 \theta^3 + 10488 \theta^2 + 9456 \theta + 3427) \\ & - 2^{28} 3^8 z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(60 \theta^2 + 180 \theta + 197) \\ & + 2^{39} 3^{10} z^4 (\theta + 1)(\theta + 2)(\theta + 3)(90 \theta^2 + 360 \theta + 407) \\ & - 2^{49} 3^{15} z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \end{aligned} $
	$A'_n = (2^{10} 3^3)^n \sum_k (-1)^k \binom{n}{k} \frac{(1/2)_k^3 (1/6)_k (5/6)_k}{k!^5}$
30	$ \begin{aligned} D = & \theta^4 - 2^4 z (4\theta + 1)(4\theta + 3)(8\theta^2 + 8\theta + 3) \\ & + 2^{12} z^2 (4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7) \end{aligned} $
	$A_n = \frac{(4n)!}{n!^2 (2n)!} \sum_{k=0}^n 2^{2(n-k)} \binom{2k}{k}^2 \binom{2n-2k}{n-k} \text{ (see [1], (6.7))}$
31	$ \begin{aligned} D = & \theta^4 - 2^4 z (4\theta + 1)(32 \theta^3 + 40 \theta^2 + 28 \theta + 7) \\ & + 2^{12} z^2 (4\theta + 1)(4\theta + 3)^2 (4\theta + 5) \end{aligned} $
	$A_n = A_n^{(-)} \text{ (see [1], (6.8))}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
32	$ \begin{aligned} D = & \theta^4 - 3z(360\theta^4 + 90\theta^3 + 27\theta^2 - 18\theta - 11) \\ & + 3z^2(145764\theta^4 + 72882\theta^3 + 24899\theta^2 + 6094\theta + 13224) \\ & - 3^4z^3(970920\theta^4 + 728190\theta^3 + 279069\theta^2 + 130766\theta - 16005) \\ & + 3^2z^4(587866086\theta^4 + 587866086\theta^3 + 249671835\theta^2 \\ & + 51395166\theta + 4547776) \\ & + 3^5z^5(8738280\theta^4 + 10922850\theta^3 + 5667111\theta^2 + 606852\theta - 464210) \\ & + 3^5z^6(1311876\theta^4 + 1967814\theta^3 + 1171557\theta^2 + 284652\theta + 134263) \\ & + 3^8z^7(3240\theta^4 + 5670\theta^3 + 3753\theta^2 + 1260\theta + 116) \\ & + 3^8z^8(3\theta + 1)^2(3\theta + 2)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 3z(2\theta + 1)(3\theta^2 + 3\theta + 1)(15\theta^2 + 15\theta + 4) \\ & - 3z^2(\theta + 1)^3(3\theta + 2)(3\theta + 4) \end{aligned} $
	$A'_n = \sum_{k,l} \binom{n}{k}^2 \binom{n}{l}^2 \binom{n+k}{n} \binom{n+l}{n} \binom{k+l}{n}$
33	$ \begin{aligned} D = & \theta^4 - 2^2z(324\theta^4 + 456\theta^3 + 321\theta^2 + 93\theta + 10) \\ & + 2^9z^2(584\theta^4 + 584\theta^3 + 4\theta^2 - 71\theta - 13) \\ & - 2^{16}z^3(324\theta^4 + 192\theta^3 + 123\theta^2 + 48\theta + 7) + 2^{24}z^4(2\theta + 1)^4 \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^{2n} \binom{2n}{k}^3$
34	$ \begin{aligned} D = & \theta^4 - z(35\theta^4 + 70\theta^3 + 63\theta^2 + 28\theta + 5) \\ & + z^2(\theta + 1)^2(259\theta^2 + 518\theta + 285) - 225z^3(\theta + 1)^2(\theta + 2)^2 \end{aligned} $
	$A_n = \sum_{i+j+k+l+m=n} \left(\frac{n!}{i!j!k!l!m!} \right)^2$
35	$ \begin{aligned} D = & \theta^4 - 2^2 \cdot 3z(192\theta^4 + 240\theta^3 + 191\theta^2 + 71\theta + 10) \\ & + 2^7 \cdot 3^2z^2(1746\theta^4 + 3960\theta^3 + 4323\theta^2 + 2247\theta + 395) \\ & - 2^{12} \cdot 3^4z^3(2538\theta^4 + 7776\theta^3 + 9915\theta^2 + 5643\theta + 1030) \\ & + 2^{17} \cdot 3^6z^4(1782\theta^4 + 6480\theta^3 + 8793\theta^2 + 4905\theta + 875) \\ & - 2^{23} \cdot 3^{11}z^5(\theta + 1)^2(3\theta + 1)(3\theta + 5) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \frac{(6k)!}{k!(2k)!(3k)!} \frac{(6n-6k)!}{(n-k)!(2n-2k)!(3n-3k)!}$
36	$D = \theta^4 - 2^4z(2\theta + 1)^2(3\theta^2 + 3\theta + 1) + 2^9z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
37	$ \begin{aligned} D = & \theta^4 - 4z(1280\theta^4 + 320\theta^3 + 94\theta^2 - 66\theta - 39) \\ & + 2^4 z^2(679936\theta^4 + 339968\theta^3 + 114304\theta^2 - 19008\theta + 11601) \\ & - 2^{13} z^3(1515520\theta^4 + 1136640\theta^3 + 430656\theta^2 + 6144\theta + 1329) \\ & + 2^{20} z^4(7868416\theta^4 + 7868416\theta^3 + 3318656\theta^2 + 156096\theta - 118695) \\ & - 2^{31} z^5(1515520\theta^4 + 1894400\theta^3 + 880576\theta^2 + 62592\theta - 4599) \\ & + 2^{40} z^6(679936\theta^4 + 1019904\theta^3 + 518016\theta^2 + 64704\theta + 19713) \\ & - 2^{57} z^7\theta(640\theta^3 + 1120\theta^2 + 617\theta + 105) \\ & + 2^{68} z^8\theta(\theta + 1)(4\theta + 1)(4\theta + 3) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 8z(2\theta + 1)(4\theta + 1)(4\theta + 3)(5\theta^2 + 5\theta + 2) \\ & + 2^{10} z^2(\theta + 1)(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7) \end{aligned} $
	$A'_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k}$
38	$ \begin{aligned} D = & \theta^4 - 2^4 z(4\theta + 1)(4\theta + 3)(3\theta^2 + 3\theta + 1) \\ & + 2^9 z^2(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7) \end{aligned} $
	$A_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$
39	$ \begin{aligned} D = & \theta^4 - 4z(320\theta^4 + 80\theta^3 + 26\theta^2 - 14\theta - 9) \\ & + 2^4 z^2(42496\theta^4 + 21248\theta^3 + 7808\theta^2 - 560\theta + 825) \\ & - 2^{12} z^3(47360\theta^4 + 35520\theta^3 + 14568\theta^2 + 1212\theta + 141) \\ & + 2^{19} z^4(61472\theta^4 + 61472\theta^3 + 27848\theta^2 + 2992\theta - 783) \\ & - 2^{27} z^5(23680\theta^4 + 29600\theta^3 + 14684\theta^2 + 1858\theta - 15) \\ & + 2^{32} z^6(42496\theta^4 + 63744\theta^3 + 34368\theta^2 + 6000\theta + 1293) \\ & - 2^{45} z^7\theta(160\theta^3 + 280\theta^2 + 163\theta + 35) + 2^{54} z^8\theta(\theta + 1)(2\theta + 1)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 8z(2\theta + 1)^3(5\theta^2 + 5\theta + 2) + 2^{10} z^2(\theta + 1)(2\theta + 1)^2(2\theta + 3)^2$
	$A'_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k}$
40	$ \begin{aligned} D = & \theta^4 - 2^4 z(64\theta^4 + 32\theta^3 + 12\theta^2 - 4\theta - 3) \\ & + 2^{13} z^2(48\theta^4 + 48\theta^3 + 22\theta^2 + 2\theta + 3) \\ & - 2^{22} z^3\theta(4\theta + 3)(4\theta^2 + 3\theta + 1) + 2^{30} z^4\theta(\theta + 1)(2\theta + 1)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5 z(2\theta + 1)^3(2\theta^2 + 2\theta + 1) + 2^{12} z^2(\theta + 1)(2\theta + 1)^2(2\theta + 3)^2$
	$A'_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
41	$D = \theta^4 - 2z(2\theta + 1)^2(7\theta^2 + 7\theta + 3) + 324z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{n+k}{k} \frac{(3k)!}{k!^3}$
42	$D = \theta^4 - 8z(2\theta + 1)^2(3\theta^2 + 3\theta + 1) + 64z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{2k}{n}^2$
43	$D = \theta^4 - 2^4z(256\theta^4 + 128\theta^3 + 44\theta^2 - 20\theta - 13) + 2^{14}z^2(384\theta^4 + 384\theta^3 + 164\theta^2 + 4\theta + 23) - 2^{26}z^3(64\theta^3 + 96\theta^2 + 49\theta + 9) + 2^{36}z^4\theta(\theta + 1)(4\theta + 1)(4\theta + 3)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 2^5z(2\theta + 1)(4\theta + 1)(4\theta + 3)(2\theta^2 + 2\theta + 1) + 2^{12}z^2(\theta + 1)(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7)$
	$A'_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$
44	$D = \theta^4 - 2^2z(544\theta^4 + 136\theta^3 + 37\theta^2 - 31\theta - 18) + 2^6z^2(27760\theta^4 + 13880\theta^3 + 4355\theta^2 + 823\theta + 2472) - 2^{10}z^3(630496\theta^4 + 472872\theta^3 + 168105\theta^2 + 75333\theta - 9924) + 2^{14}z^4(5400856\theta^4 + 5400856\theta^3 + 2146159\theta^2 + 393923\theta + 22032) - 2^{18}z^5(630496\theta^4 + 788120\theta^3 + 365135\theta^2 - 5981\theta - 55194) + 2^{22}z^6(27760\theta^4 + 41640\theta^3 + 21705\theta^2 + 2373\theta + 1752) - 2^{26}z^7\theta(544\theta^3 + 952\theta^2 + 547\theta + 119) + 2^{30}z^8\theta(\theta + 1)(2\theta + 1)^2$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 4z(2\theta + 1)^3(17\theta^2 + 17\theta + 5) + 16z^2(\theta + 1)(2\theta + 1)^2(2\theta + 3)^2$
	$A'_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$
45	$D = \theta^4 - 4z(2\theta + 1)^2(7\theta^2 + 7\theta + 2) - 2^7z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^3$
46	$D = \theta^4 - 6z(2\theta + 1)^2(9\theta^2 + 9\theta + 4) + 324z^2(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4)$
	$A_n = \binom{2n}{n} \sum_{k=0}^n \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
47	$D = \theta^4 - 2^4 3z(2\theta + 1)^2(18\theta^2 + 18\theta + 5) + 2^{10} 3^4 z^2(2\theta + 1)(2\theta + 3)(3\theta + 1)(3\theta + 5)$
	$A_n = \binom{2n}{n} \sum_{k=0}^n \frac{(6k)!}{k!(2k)!(3k)!} \frac{(6n-6k)!}{(n-k)!(2n-2k)!(3n-3k)!}$
48	$D = \theta^4 - 12z(3\theta + 1)(3\theta + 2)(3\theta^2 + 3\theta + 1) + 288z^2(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$
	$A_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$
49	$D = \theta^4 - 2^2 \cdot 3z(144\theta^4 + 72\theta^3 + 26\theta^2 - 10\theta - 7) + 2^4 \cdot 3^4 z^2(864\theta^4 + 864\theta^3 + 384\theta^2 + 24\theta + 53) - 2^{11} \cdot 3^8 z^3\theta(24\theta^3 + 36\theta^2 + 19\theta + 4) + 2^{16} \cdot 3^{10} z^4\theta(\theta + 1)(3\theta + 1)(3\theta + 2)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 24z(2\theta + 1)(3\theta + 1)(3\theta + 2)(2\theta^2 + 2\theta + 1) + 2304z^2(\theta + 1)(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$
	$A'_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$
50	$D = \theta^4 - 3z(720\theta^4 + 180\theta^3 + 56\theta^2 - 34\theta - 21) + 3^4 z^2(23904\theta^4 + 11952\theta^3 + 4226\theta^2 - 472\theta + 439) - 3^7 z^3(426240\theta^4 + 319680\theta^3 + 126672\theta^2 + 6828\theta + 868) + 3^{10} z^4(4425984\theta^4 + 4425984\theta^3 + 1943584\theta^2 + 158704\theta - 61035) - 2^7 \cdot 3^{13} z^5(213120\theta^4 + 266400\theta^3 + 128456\theta^2 + 13202\theta - 364) + 2^{13} \cdot 3^{16} z^6(11952\theta^4 + 17928\theta^3 + 9417\theta^2 + 1443\theta + 356) - 2^{20} \cdot 3^{19} z^7\theta(180\theta^3 + 315\theta^2 + 179\theta + 35) + 2^{24} \cdot 3^{22} z^8\theta(\theta + 1)(3\theta + 1)(3\theta + 2)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 6z(2\theta + 1)(3\theta + 1)(3\theta + 2)(5\theta^2 + 5\theta + 2) + 576z^2(\theta + 1)(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$
	$A'_n = \frac{(3n)!}{n!^3} \sum_{j+k+l+m=n} \left(\frac{n!}{j!k!l!m!} \right)^2 = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k}$ $= 4^{-n} \frac{(3n)!}{n!^3} \binom{2n}{n}^3 \sum_{k=0}^n \binom{n}{k}^4 \binom{2n}{2k}^{-3}$
51	$D = \theta^4 - 4z(4\theta + 1)(4\theta + 3)(11\theta^2 + 11\theta + 3) - 16z^2(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7)$
	$A_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
52	$ \begin{aligned} D = & \theta^4 - 2^2 z(2176\theta^4 + 544\theta^3 + 131\theta^2 - 141\theta - 78) \\ & + 2^7 z^2(222080\theta^4 + 111040\theta^3 + 31370\theta^2 + 3690\theta + 19557) \\ & - 2^{14} z^3(2521984\theta^4 + 1891488\theta^3 + 613311\theta^2 + 261807\theta - 37362) \\ & + 2^{20} z^4(21603424\theta^4 + 21603424\theta^3 + 7909529\theta^2 + 1234089\theta + 58191) \\ & - 2^{26} z^5(2521984\theta^4 + 3152480\theta^3 + 1362025\theta^2 - 102855\theta - 233532) \\ & + 2^{31} z^6(222080\theta^4 + 333120\theta^3 + 163230\theta^2 + 9150\theta + 13317) \\ & - 2^{38} z^7\theta(2176\theta^3 + 3808\theta^2 + 2069\theta + 357) \\ & + 2^{44} z^8\theta(\theta + 1)(4\theta + 1)(4\theta + 3) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 4z(2\theta + 1)(4\theta + 1)(4\theta + 3)(17\theta^2 + 17\theta + 5) \\ & + 16z^2(\theta + 1)(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7) \end{aligned} $
	$A'_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$
53	$ \begin{aligned} D = & \theta^4 - 3z(1224\theta^4 + 306\theta^3 + 79\theta^2 - 74\theta - 42) \\ & + 3^4 z^2(62460\theta^4 + 31230\theta^3 + 9365\theta^2 + 1490\theta + 5534) \\ & - 3^7 z^3(1418616\theta^4 + 1063962\theta^3 + 363459\theta^2 + 159618\theta - 21788) \\ & + 3^{10} z^4(12151926\theta^4 + 12151926\theta^3 + 4660081\theta^2 + 800926\theta + 41364) \\ & - 3^{13} z^5(1418616\theta^4 + 1773270\theta^3 + 796925\theta^2 - 33190\theta - 127418) \\ & + 3^{16} z^6(62460\theta^4 + 93690\theta^3 + 47535\theta^2 + 4110\theta + 3854) \\ & - 3^{19} z^7\theta(1224\theta^3 + 2142\theta^2 + 1201\theta + 238) \\ & + 3^{22} z^8\theta(\theta + 1)(3\theta + 1)(3\theta + 2) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 3z(2\theta + 1)(3\theta + 1)(3\theta + 2)(17\theta^2 + 17\theta + 5) \\ & + 9z^2(\theta + 1)(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5) \end{aligned} $
	$A'_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$
54	$ \begin{aligned} D = & \theta^4 - 2z(6\theta^4 + 10\theta^3 + 10\theta^2 + 5\theta + 1) \\ & + 4z^2(1039\theta^4 + 4146\theta^3 + 5707\theta^2 + 3127\theta + 592) \\ & - 8z^3(\theta + 1)(4116\theta^3 + 18512\theta^2 + 27768\theta + 13889) \\ & + 16z^4(\theta + 1)(\theta + 2)(6159\theta^2 + 24631\theta + 25143) \\ & - 64z^5(\theta + 1)(\theta + 2)(\theta + 3)(2051\theta + 5127) \\ & + 65600z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_k (-1)^k 2^{n-2k} \binom{n}{2k} \left(\frac{(4k)!}{k!^2(2k)!} \right)^2$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
55	$D = 9\theta^4 - 12z(208\theta^4 + 224\theta^3 + 163\theta^2 + 51\theta + 6)$ $+ 2^9 z^2(32\theta^4 - 928\theta^3 - 1606\theta^2 - 837\theta - 141)$ $+ 2^{16} z^3(144\theta^4 + 576\theta^3 + 467\theta^2 + 144\theta + 15) - 2^{24} z^4(2\theta + 1)^4$
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^2 \binom{2n}{2k}$
56	$D = \theta^4 - 2^4 z(22\theta^4 + 8\theta^3 + 9\theta^2 + 5\theta + 1)$ $+ 2^9 z^2(94\theta^4 + 88\theta^3 + 97\theta^2 + 45\theta + 8)$ $- 2^{14} z^3(194\theta^4 + 336\theta^3 + 371\theta^2 + 195\theta + 41)$ $+ 2^{19} \cdot 3z^4(64\theta^4 + 176\theta^3 + 217\theta^2 + 129\theta + 30) - 2^{27} \cdot 3^2 z^5(\theta + 1)^4$
	$A_n = \sum_{k=0}^n \binom{n}{k}^{-1} \binom{2k}{k}^3 \binom{2n-2k}{n-k}^3$
57	$D = \theta^4 - 2^2 z(8\theta^4 + 10\theta^3 + 10\theta^2 + 5\theta + 1)$ $+ 2^4 z^2(28\theta^4 + 70\theta^3 + 95\theta^2 + 65\theta + 18)$ $- 2^7 z^3(28\theta^4 + 105\theta^3 + 180\theta^2 + 150\theta + 49)$ $+ 2^8 z^4(799\theta^4 + 6182\theta^3 + 15629\theta^2 + 13660\theta + 3856)$ $- 2^{11} z^5(\theta + 1)(1486\theta^3 + 11082\theta^2 + 25388\theta + 17397)$ $+ 2^{13} z^6(\theta + 1)(\theta + 2)(2201\theta^2 + 13185\theta + 18827)$ $- 2^{16} z^7(\theta + 1)(\theta + 2)(\theta + 3)(731\theta + 2557)$ $+ 2^{17} \cdot 5 \cdot 73 z^8(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)$
	$A_n = \sum_k (-1)^k 4^{n-4k} \binom{n}{4k} \left(\frac{(6k)!}{k!(2k)!(3k)!} \right)^2$
58	$D = \theta^4 - 4z(2\theta + 1)^2(10\theta^2 + 10\theta + 3) + 144z^2(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$
59	$D = 49\theta^4 - 2z(1799\theta^4 + 3640\theta^3 + 3045\theta^2 + 1225\theta + 196)$ $+ 2^2 z^2(13497\theta^4 + 55536\theta^3 + 81222\theta^2 + 50337\theta + 11396)$ $- 2^3 z^3(17201\theta^4 + 114996\theta^3 + 248466\theta^2 + 202629\theta + 55412)$ $- 2^4 z^4(5762\theta^4 + 29668\theta^3 + 48150\theta^2 + 31741\theta + 7412)$ $- 2^5 \cdot 3z^5(4\theta + 5)(3\theta + 2)(3\theta + 4)(4\theta + 3)$
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2n-k}{n} \binom{n+k}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
60	$ \begin{aligned} D = & \theta^4 - 2z(248\theta^4 + 62\theta^3 + 23\theta^2 - 8\theta - 6) \\ & + 2^2 z^2(24792\theta^4 + 12396\theta^3 + 5153\theta^2 + 704\theta + 1126) \\ & - 2^2 z^3(2549440\theta^4 + 1912080\theta^3 + 882790\theta^2 + 230890\theta + 21527) \\ & + 2^3 z^4(71646752\theta^4 + 71646752\theta^3 + 36508992\theta^2 + 8176202\theta - 785881) \\ & - 2^4 \cdot 3z^5(367119360\theta^4 + 458899200\theta^3 + 256824520\theta^2 \\ & + 54290940\theta + 1905253) \\ & + 2^4 \cdot 3^2 z^6(2056347648\theta^4 + 3084521472\theta^3 + 1880419968\theta^2 \\ & + 499802112\theta + 140357665) \\ & - 2^9 \cdot 3^5 z^7(20570112\theta^4 + 35997696\theta^3 + 23709888\theta^2 \\ & + 7437024\theta + 640447) \\ & + 2^{12} \cdot 3^8 z^8(24\theta + 5)(24\theta + 11)(24\theta + 13)(24\theta + 19) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(31\theta^4 + 62\theta^3 + 54\theta^2 + 23\theta + 4) \\ & + 12z^2(\theta + 1)(3\theta + 2)(3\theta + 4)(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{2n-k}{n} \binom{n+k}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$
61	$ \begin{aligned} D = & \theta^4 - 2^4 3^2 z(6\theta + 1)(6\theta + 5)(72\theta^2 + 72\theta + 31) \\ & + 2^{12} 3^8 z^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11) \end{aligned} $
	$A_n = \frac{(6n)!}{n!(2n)!(3n)!} \cdot 432^n \sum_k (-1)^k \binom{-5/6}{k} \binom{-1/6}{n-k}^2$
62	$ \begin{aligned} D = & \theta^4 - 12z(6\theta + 1)(6\theta + 5)(7\theta^2 + 7\theta + 2) \\ & + 1152z^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11) \end{aligned} $
	$A_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{n}{k}^3$
63	$ \begin{aligned} D = & \theta^4 - 12z(6\theta + 1)(6\theta + 5)(11\theta^2 + 11\theta + 3) \\ & + 144z^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11) \end{aligned} $
	$A_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
64	$D = \theta^4 - 12z(6\theta + 1)(6\theta + 5)(10\theta^2 + 10\theta + 3) + 1296z^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11)$
	$A_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$
65	$D = \theta^4 - 48z(6\theta + 1)(6\theta + 5)(3\theta^2 + 3\theta + 1) + 4608z^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11)$
	$A_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$
66	$D = \theta^4 - 2^2 \cdot 3z(2880\theta^4 + 720\theta^3 + 194\theta^2 - 166\theta - 93) + 2^4 \cdot 3^4 z^2(382464\theta^4 + 191232\theta^3 + 59648\theta^2 - 15088\theta + 5833) - 2^{12} \cdot 3^7 z^3(426240\theta^4 + 319680\theta^3 + 113352\theta^2 - 5412\theta - 305) + 2^{19} \cdot 3^{10} z^4(553248\theta^4 + 553248\theta^3 + 219896\theta^2 - 1432\theta - 9327) - 2^{28} \cdot 3^{13} z^5(106560\theta^4 + 133200\theta^3 + 58678\theta^2 + 1321\theta - 518) + 2^{32} \cdot 3^{16} z^6(382464\theta^4 + 573696\theta^3 + 277440\theta^2 + 22704\theta + 10669) - 2^{45} \cdot 3^{19} z^7\theta(1440\theta^3 + 2520\theta^2 + 1327\theta + 175) + 2^{54} \cdot 3^{22} z^8\theta(\theta + 1)(6\theta + 1)(6\theta + 5)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 24z(2\theta + 1)(6\theta + 1)(6\theta + 5)(5\theta^2 + 5\theta + 2) + 9216z^2(\theta + 1)(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11)$
	$A'_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k}$
67	$D = \theta^4 - 2^4 \cdot 3z(576\theta^4 + 288\theta^3 + 92\theta^2 - 52\theta - 31) + 2^{13} \cdot 3^4 z^2(432\theta^4 + 432\theta^3 + 174\theta^2 - 6\theta + 25) - 2^{22} \cdot 3^8 z^3\theta(4\theta + 1)(12\theta^2 + 15\theta + 5) + 2^{30} \cdot 3^{10} z^4\theta(\theta + 1)(6\theta + 1)(6\theta + 5)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 96z(2\theta + 1)(6\theta + 1)(6\theta + 5)(3\theta^2 + 3\theta + 1) + 36864z^2(\theta + 1)(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11)$
	$A'_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$
68	$D = \theta^4 - 4z(4\theta + 1)(4\theta + 3)(7\theta^2 + 7\theta + 2) - 128z^2(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7)$
	$A_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{n}{k}^3$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
69	$D = \theta^4 - 4z(4\theta + 1)(4\theta + 3)(10\theta^2 + 10\theta + 3) + 144z^2(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7)$
	$A_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$
70	$D = \theta^4 - 3z(3\theta + 1)(3\theta + 2)(10\theta^2 + 10\theta + 3) + 81z^2(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$
	$A_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$
71	$D = \theta^4 + 2^4z(39\theta^4 - 42\theta^3 - 29\theta^2 - 8\theta - 1) + 2^{11}z^2\theta(37\theta^3 - 137\theta^2 - 10\theta - 1) - 2^{16}z^3(181\theta^4 + 456\theta^3 + 353\theta^2 + 132\theta + 19) - 2^{23}5z^4(36\theta^4 + 60\theta^3 + 36\theta^2 + 6\theta - 1) + 2^{30}5^2z^5(\theta + 1)^4$
	$A_n = 2^{-n} \sum_{k=0}^n \binom{2k}{k}^4 \binom{2n-2k}{n-k}^4 \binom{n}{k}^{-3}$
72	$D = \theta^4 - 2z(136\theta^4 + 260\theta^3 + 198\theta^2 + 68\theta + 9) + 2^2z^2(1048\theta^4 + 2584\theta^3 + 3102\theta^2 + 2188\theta + 703) - 2^4z^3(1552\theta^4 + 4632\theta^3 + 6552\theta^2 + 4614\theta + 1089) + 2^4z^4(4112\theta^4 + 14368\theta^3 + 21528\theta^2 + 14344\theta + 3585) - 2^{16}z^5(\theta + 1)^4$
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}^3 \binom{2n}{2k}^{-1}$
73	$D = \theta^4 - 18z(42\theta^4 + 60\theta^3 + 45\theta^2 + 15\theta + 2) + 3 \cdot 18^2z^2(180\theta^4 + 432\theta^3 + 453\theta^2 + 222\theta + 40) - 3 \cdot 18^4z^3(2\theta + 1)^2(13\theta^2 + 29\theta + 20) + 18^6z^4(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n} \sum_{k=0}^n \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3} \binom{2n}{k} \binom{n}{k}^{-1}$
74	$D = \theta^4 - 2 \cdot 3z(99\theta^4 + 36\theta^3 + 39\theta^2 + 21\theta + 4) + 2^2 \cdot 3^2z^2(3807\theta^4 + 3564\theta^3 + 3798\theta^2 + 1683\theta + 284) - 2^3 \cdot 3^5z^3(7857\theta^4 + 13608\theta^3 + 14562\theta^2 + 7317\theta + 1444) + 2^4 \cdot 3^9z^4(2592\theta^4 + 7128\theta^3 + 8550\theta^2 + 4851\theta + 1052) - 2^5 \cdot 3^{13}z^5(3\theta + 2)(3\theta + 4)(6\theta + 5)(6\theta + 7)$
	$A_n = \sum_{k=0}^n \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{n}{k}^{-1}$
75	$D = \theta^4 - 10z(2\theta + 1)^2(26\theta^2 + 26\theta + 5) + 4z^2(2\theta + 1)(2\theta + 3)(774\theta^2 + 1548\theta + 823) - 3088z^3(2\theta + 1)(2\theta + 3)^2(2\theta + 5) + 4112z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)$
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \frac{(4k)!}{k!^4}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
76	$ \begin{aligned} D = & \theta^4 - z(1029\theta^4 + 2058\theta^3 + 1482\theta^2 + 453\theta + 49) \\ & + z^2(\theta + 1)^2(4106\theta^2 + 8212\theta + 5007) \\ & - z^3(\theta + 1)(\theta + 2)(6154\theta^2 + 18462\theta + 14809) \\ & + 4101z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & - 1025z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(4k)!}{k!^4}$
77	$ \begin{aligned} D = & \theta^4 - z(16416\theta^4 + 4116\theta^3 + 906\theta^2 - 1152\theta - 625) \\ & + z^2(101122496\theta^4 + 50758192\theta^3 + 13326174\theta^2 + 773908\theta + 8758171) \\ & - z^3(277299334656\theta^4 + 209187183168\theta^3 + 63952323504\theta^2 \\ & + 25856143860\theta - 4222453166) \\ & + z^4(286996730758656\theta^4 + 290319472045824\theta^3 + 102144306403616\theta^2 \\ & + 13734541538384\theta + 29090804825) \\ & - 2^4 \cdot 3 \cdot 5^2 \cdot 41z^5(4\theta - 1)(23108277888\theta^3 + 52094682096\theta^2 \\ & + 44583010636\theta + 13882711785) \\ & + 2^5 \cdot 5^4 \cdot 41^2z^6(4\theta - 1)(4\theta + 3)(50561248\theta^2 + 126501592\theta + 87298415) \\ & - 2^8 \cdot 3 \cdot 5^6 \cdot 41^3z^7(4\theta - 1)(4\theta + 3)(4\theta + 7)(1368\theta + 2395) \\ & + 2^8 \cdot 5^8 \cdot 41^4z^8(4\theta - 1)(4\theta + 3)(4\theta + 7)(4\theta + 11) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2(2\theta + 1)(1029\theta^4 + 2058\theta^3 + 1482\theta^2 + 453\theta + 49) \\ & + 4(\theta + 1)(2\theta + 1)(2\theta + 3)(4106\theta^2 + 8212\theta + 5007) \\ & - 8(2\theta + 1)(2\theta + 3)(2\theta + 5)(6154\theta^2 + 18462\theta + 14809) \\ & + 48 \cdot 1367(\theta + 2)(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \\ & - 2^5 \cdot 5^2 \cdot 41(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)(2\theta + 9) \end{aligned} $
	$A'_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(4k)!}{k!^4}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
78	$ \begin{aligned} D = & \theta^4 - z(16392\theta^4 + 4102\theta^3 + 900\theta^2 - 1151\theta - 624) \\ & + z^2(100778012\theta^4 + 50454570\theta^3 + 13189539\theta^2 + 803211\theta + 8784048) \\ & - z^3(275482230840\theta^4 + 207014719614\theta^3 + 62883681150\theta^2 \\ & + 25936450350\theta - 4109625168) \\ & + z^4(282850876768326\theta^4 + 283952402645202\theta^3 + 98665243460597\theta^2 \\ & + 13790192460644\theta + 614356880784) \\ & - 17 \cdot 241z^5\theta(275482230840\theta^3 + 620238065874\theta^2 \\ & + 530195542680\theta + 165017893373) \\ & + 17^2 \cdot 241^2z^6\theta(\theta + 1)(100778012\theta^2 + 252010594\theta + 173831131) \\ & - 2 \cdot 17^3 \cdot 241^3z^7\theta(\theta + 1)(\theta + 2)(8196\theta + 14345) \\ & + 17^4 \cdot 241^4z^8\theta(\theta + 1)(\theta + 2)(\theta + 3) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - z(2\theta + 1)(2051\theta^4 + 4102\theta^3 + 2951\theta^2 + 900\theta + 97) \\ & + z^2(\theta + 1)(20495\theta^4 + 81980\theta^3 + 132457\theta^2 + 100954\theta + 30175) \\ & - 2z^3(\theta + 1)(\theta + 2)(2\theta + 3)(10245\theta^2 + 30735\theta + 27983) \\ & + z^4(\theta + 1)(\theta + 2)(\theta + 3)(40975\theta^2 + 163900\theta + 173889) \\ & - 10243z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \\ & + 4097z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k}^2 \frac{(4k)!}{k!^4}$
79	$ \begin{aligned} D = & \theta^4 - z(3130\theta^4 + 6260\theta^3 + 4385\theta^2 + 1255\theta + 121) \\ & + 15z^2(\theta + 1)^2(834\theta^2 + 1668\theta + 1001) \\ & - 5z^3(\theta + 1)(\theta + 2)(3752\theta^2 + 11256\theta + 9005) \\ & + 12505z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & - 3126z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	same $K(q)$ as in #1
	$A_n = \sum_{k=0}^n \binom{n}{k} \frac{(5k)!}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
80	$ \begin{aligned} D = & \theta^4 - 2z(25016\theta^4 + 6260\theta^3 + 1255\theta^2 - 1875\theta - 998) \\ & + 2^2 z^2 (234725112\theta^4 + 117512640\theta^3 + 28486025\theta^2 - 10620\theta + 20240412) \\ & - 2^4 z^3 (489688550224\theta^4 + 367970437920\theta^3 + 104723790940\theta^2 \\ & + 41242253625\theta - 7232306723) \\ & + 2^6 z^4 (383914650187780\theta^4 + 385383011813200\theta^3 + 126338801127175\theta^2 \\ & + 14531449104300\theta + 193021814409) \\ & - 16 \cdot 24 \cdot 521 z^5 (4\theta - 1)(489688550224\theta^3 + 1102503263256\theta^2 \\ & + 937734079014\theta + 289728842991) \\ & + 4 \cdot 24^2 \cdot 521^2 z^6 (4\theta - 1)(4\theta + 3)(234725112\theta^2 \\ & + 586962864\theta + 404126249) \\ & - 16 \cdot 24^3 \cdot 521^3 z^7 (4\theta - 1)(4\theta + 3)(4\theta + 7)(3127\theta + 5473) \\ & + 24^4 \cdot 521^4 z^8 (4\theta - 1)(4\theta + 3)(4\theta + 7)(4\theta + 11) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(3130\theta^4 + 6260\theta^3 + 4385\theta^2 + 1255\theta + 121) \\ & + 60z^2(\theta + 1)(2\theta + 1)(2\theta + 3)(834\theta^2 + 1668\theta + 1001) \\ & - 40z^3(2\theta + 1)(2\theta + 3)(2\theta + 5)(3752\theta^2 + 11256\theta + 9005) \\ & + 2^4 \cdot 5 \cdot 41 \cdot 61 z^4(\theta + 2)(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \\ & - 2^6 \cdot 3 \cdot 521 z^5(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)(2\theta + 9) \end{aligned} $
	$A'_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \frac{(5k)!}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
81	$ \begin{aligned} D = & \theta^4 - z(50008\theta^4 + 12506\theta^3 + 2504\theta^2 - 3749\theta - 1995) \\ & + z^2(937850028\theta^4 + 469125042\theta^3 + 113531285\theta^2 + 50011\theta + 81040890) \\ & - z^3(7818126050056\theta^4 + 5867345737626\theta^3 + 1665767387626\theta^2 \\ & + 660699781296\theta - 114681662235) \\ & + z^4(24453139064250070\theta^4 + 24484407817250210\theta^3 \\ & + 7988410528587745\theta^2 + 932188857443850\theta + 28724348623005) \\ & - 3^3 \cdot 463z^5\theta(7818126050056\theta^3 + 17594534812710\theta^2 \\ & + 14959207512780\theta + 4621083875125) \\ & + 3^6 \cdot 463^2z^6\theta(\theta + 1)(937850028\theta^2 + 2344825098\theta + 1614171341) \\ & - 2 \cdot 3^9 \cdot 463^3z^7\theta(\theta + 1)(\theta + 2)(25004\theta + 43759) \\ & + 3^{12} \cdot 463^4z^8\theta(\theta + 1)(\theta + 2)(\theta + 3) \end{aligned} $
	same $K(q)$ as in #80 the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - z(2\theta + 1)(6253\theta^4 + 12506\theta^3 + 8757\theta^2 + 2504\theta + 241) \\ & + z^2(\theta + 1)(62515\theta^4 + 250060\theta^3 + 402605\theta^2 + 305090\theta + 90511) \\ & - 70z^3(\theta + 1)(\theta + 2)(2\theta + 3)(893\theta^2 + 2679\theta + 2429) \\ & + 5z^4(\theta + 1)(\theta + 2)(\theta + 3)(25003\theta^2 + 100012\theta + 106013) \\ & - 31253z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \\ & + 3^3 \cdot 463z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(5k)!}{k!^5}$
82	$ \begin{aligned} D = & \theta^4 - z(186631\theta^4 + 46662\theta^3 + 8428\theta^2 - 14903\theta - 7812) \\ & + z^2(13061813781\theta^4 + 6531700068\theta^3 + 1452108415\theta^2 \\ & - 105618806\theta + 1116194184) \\ & - z^3(406305132943139\theta^4 + 304784361622746\theta^3 \\ & + 80477825393615\theta^2 + 30347763996276\theta - 5895787146228) \\ & + 13 \cdot 37 \cdot 97z^4\theta(101592610503011\theta^3 + 101629619768568\theta^2 \\ & + 31089416550198\theta + 2838662698240) \\ & - 13^2 \cdot 37^2 \cdot 97^2z^5\theta(\theta + 1)(6531186837\theta^2 + 10886104581\theta + 5020240741) \\ & + 13^3 \cdot 37^3 \cdot 97^3z^6\theta(\theta + 1)(\theta + 2)(139975\theta + 186639) \\ & - 13^4 \cdot 37^4 \cdot 97^4z^7\theta(\theta + 1)(\theta + 2)(\theta + 3) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 7z(2\theta + 1)(3333\theta^4 + 6666\theta^3 + 4537\theta^2 + 1204\theta + 103) \\ & + z^2(\theta + 1)(8233295\theta^4 + 933180\theta^3 + 1496985\theta^2 + 1127610\theta + 331951) \\ & - 10z^3(\theta + 1)(\theta + 2)(\theta + 3)(23329\theta^2 + 69987\theta + 63183) \\ & + 5z^4(\theta + 1)(\theta + 2)(\theta + 3)(93315\theta^2 + 373260\theta + 395293) \\ & - 3 \cdot 59 \cdot 659z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \\ & + 13 \cdot 37 \cdot 97z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{n}{k} \frac{(6k)!}{k!^6}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
83	$ \begin{aligned} D = & \theta^4 - 2^4 z(88\theta^4 + 32\theta^3 + 33\theta^2 + 17\theta + 3) \\ & + 2^9 z^2(1504\theta^4 + 1408\theta^3 + 1436\theta^2 + 596\theta + 93) \\ & - 2^{18} z^3(776\theta^4 + 1344\theta^3 + 1381\theta^2 + 651\theta + 117) \\ & + 3 \cdot 2^{23} z^4(2\theta + 1)(512\theta^3 + 1152\theta^2 + 1054\theta + 339) \\ & - 9 \cdot 2^{31} z^5(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \binom{2n}{2k}^{-1} \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!}$
84	$ \begin{aligned} D = & \theta^4 - 4z(32\theta^4 + 64\theta^3 + 63\theta^2 + 31\theta + 6) \\ & + 256z^2(\theta + 1)^2(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{2n}{2k}^{-1} \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!}$
85	$ \begin{aligned} D = & \theta^4 - 2z(2\theta + 1)^2(866\theta^2 + 866\theta + 121) \\ & + 4z^2(2\theta + 1)(2\theta + 3)(5190\theta^2 + 10380\theta + 5431) \\ & - 2^4 \cdot 1297z^3(2\theta + 1)(2\theta + 3)^2(2\theta + 5) \\ & + 2^4 \cdot 7 \cdot 13 \cdot 19z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(6k)!}{k!(2k)!(3k)!}$
86	$ \begin{aligned} D = & \theta^4 - z(6917\theta^4 + 13834\theta^3 + 9610\theta^2 + 2693\theta + 241) \\ & + z^2(\theta + 1)^2(27658\theta^2 + 55316\theta + 33039) \\ & - z^3(\theta + 1)(\theta + 2)(41482\theta^2 + 124446\theta + 99481) \\ & + 27653z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & - 31 \cdot 223z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	same $K(q)$ as in #85
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(6k)!}{k!^3(3k)!}$
87	$ \begin{aligned} D = & \theta^4 - z(3145\theta^4 + 6282\theta^3 + 4403\theta^2 + 1262\theta + 122) \\ & + 4z^2(2\theta + 1)(6270\theta^3 + 18804\theta^2 + 20058\theta + 7523) \\ & - 8z^3(2\theta + 1)(2\theta + 3)(9395\theta^2 + 28181\theta + 22544) \\ & + 16z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(6260\theta + 12519) \\ & - 2^4 \cdot 3 \cdot 7 \cdot 149z^5(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \frac{(5k)!}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
88	$ \begin{aligned} D = & \theta^4 - z(110624\theta^4 + 27668\theta^3 + 5386\theta^2 - 8448\theta - 4465) \\ & + z^2(4589568448\theta^4 + 2296111664\theta^3 + 542921054\theta^2 \\ & - 10713580\theta + 395878043) \\ & - z^3(84647953763840\theta^4 + 63541034834496\theta^3 + 17687257564080\theta^2 \\ & + 6901304092148\theta - 1219151757870) \\ & + z^4(586017416803534336\theta^4 + 587032971970653952\theta^3 \\ & + 188288528023623968\theta^2 + 20443050826683472\theta + 526999484846681) \\ & - 2^4 \cdot 31 \cdot 223z^5(4\theta - 1)(21161988440960\theta^3 + 47628241370064\theta^2 \\ & + 40437124384228\theta + 12464650057531) \\ & + 2^5 \cdot 31^2 \cdot 223^2z^6(4\theta - 1)(4\theta + 3)(2294784224\theta^2 \\ & + 5737624280\theta + 3947717231) \\ & - 2^8 \cdot 31^3 \cdot 223^3z^7(4\theta - 1)(4\theta + 3)(4\theta + 7)(27656\theta + 48401) \\ & + 2^8 \cdot 31^4 \cdot 223^4z^8(4\theta - 1)(4\theta + 3)(4\theta + 7)(4\theta + 11) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(6917\theta^4 + 13834\theta^3 + 9610\theta^2 + 2693\theta + 241) \\ & + 4z^2(\theta + 1)(2\theta + 1)(2\theta + 3)(27658\theta^2 + 55316\theta + 33039) \\ & - 8z^3(2\theta + 1)(2\theta + 3)(2\theta + 5)(41482\theta^2 + 124446\theta + 99481) \\ & + 442448z^4(\theta + 2)(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \\ & - 221216z^5(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)(2\theta + 9) \end{aligned} $
	$A'_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(6k)!}{k!^3(3k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
89	$ \begin{aligned} D = & \theta^4 - z(110600\theta^4 + 27654\theta^3 + 5380\theta^2 - 8447\theta - 4464) \\ & + z^2(4587245596\theta^4 + 2294065194\theta^3 + 542010659\theta^2 \\ & - 10506997\theta + 396053424) \\ & - z^3(84565362438200\theta^4 + 63442370368638\theta^3 + 17639445591678\theta^2 \\ & + 6905443579438\theta - 1214113199184) \\ & + z^4(584748316984283206\theta^4 + 585086560085496018\theta^3 \\ & + 187240905834613493\theta^2 + 20468485628797540\theta + 703267501469328) \\ & - 43 \cdot 643z^5\theta(84565362438200\theta^3 + 190290414025938\theta^2 \\ & + 161530840617496\theta + 49787404171133) \\ & + 43^2 \cdot 643^2z^6\theta(\theta + 1)(4587245596\theta^2 + 11468556386\theta + 7890274651) \\ & - 2 \cdot 43^3 \cdot 643^3z^7\theta(\theta + 1)(\theta + 2)(55300\theta + 96777) \\ & + 43^4 \cdot 643^4z^8\theta(\theta + 1)(\theta + 2)(\theta + 3) \end{aligned} $
	same $K(q)$ as in #88
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - z(2\theta + 1)(13827\theta^4 + 27654\theta^3 + 19207\theta^2 + 5380\theta + 481) \\ & + z^2(\theta + 1)(138255\theta^4 + 553020\theta^3 + 889449\theta^2 + 672858\theta + 199135) \\ & - 2z^3(\theta + 1)(\theta + 2)(2\theta + 3)(69125\theta^2 + 207375\theta + 187791) \\ & + z^4(\theta + 1)(\theta + 2)(\theta + 3)(276495\theta^2 + 1105980\theta + 1172033) \\ & - 69123z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \\ & + 27649z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k}^2 \frac{(6k)!}{k!^3(3k)!}$
90	$ \begin{aligned} D = & \theta^4 - 2z(2\theta + 1)^2(56\theta^2 + 56\theta + 13) \\ & + 20z^2(2\theta + 1)(2\theta + 3)(66\theta^2 + 132\theta + 71) \\ & - 1312z^3(2\theta + 1)(2\theta + 3)^2(2\theta + 5) \\ & + 1744z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(3k)!}{k!^3}$
91	$ \begin{aligned} D = & \theta^4 - z(437\theta^4 + 874\theta^3 + 646\theta^2 + 209\theta + 25) \\ & + z^2(\theta + 1)^2(1738\theta^2 + 3476\theta + 2151) \\ & - z^3(\theta + 1)(\theta + 2)(2602\theta^2 + 7806\theta + 6277) \\ & + 1733z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & - 433z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k}^2 \frac{(3k)!}{k!^3}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
92	$ \begin{aligned} D = & \theta^4 - 2^2 z(12\theta^4 + 16\theta^3 + 14\theta^2 + 6\theta + 1) \\ & + 2^4 z^2(789\theta^4 + 3076\theta^3 + 4167\theta^2 + 2222\theta + 385) \\ & - 2^7 z^3(2\theta + 1)(1498\theta^3 + 6701\theta^2 + 9987\theta + 4949) \\ & + 2^8 z^4(2\theta + 1)(2\theta + 3)(4434\theta^2 + 1769\theta + 18003) \\ & - 2^{11} z^5(2\theta + 1)(2\theta + 3)(2\theta + 5)(1470\theta + 3671) \\ & + 2^{12} \cdot 733 z^6(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_k (-1)^k 2^{n-2k} \binom{n}{2k} \binom{2k}{k}^2 \binom{6k}{2k}$
93	$ \begin{aligned} D = & \theta^4 - z(448\theta^4 + 872\theta^3 + 648\theta^2 + 212\theta + 26) \\ & + z^2(7008\theta^4 + 17376\theta^3 + 20664\theta^2 + 14496\theta + 4652) \\ & - z^3(41728\theta^4 + 124800\theta^3 + 175488\theta^2 + 122784\theta + 28576) \\ & + z^4(110848\theta^4 + 387584\theta^3 + 577920\theta^2 + 381824\theta + 94096) \\ & - 3072 z^5(\theta + 1)^2(6\theta + 5)(6\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n}{2k}^{-1} \frac{(3k)!}{k!^3}$
94	$ \begin{aligned} D = & \theta^4 - z(27656\theta^4 + 6918\theta^3 + 1420\theta^2 - 2039\theta - 1092) \\ & + z^2(286848028\theta^4 + 143534634\theta^3 + 35472587\theta^2 + 608435\theta + 24800580) \\ & - z^3(1322624277560\theta^4 + 993115489662\theta^3 + 287226757326\theta^2 \\ & + 115129154686\theta - 19751450796) \\ & + z^4(2289130534388806\theta^4 + 2294419884217554\theta^3 + 762036995515973\theta^2 \\ & + 93972178515892\theta + 2628910994892) \\ & - 31 \cdot 223 z^5\theta(1322624277560\theta^3 + 2977051906002\theta^2 \\ & + 2534884699840\theta + 784655821733) \\ & + 31^2 \cdot 223^2 z^6\theta(\theta + 1)(286848028\theta^2 + 717230690\theta + 494009059) \\ & - 2 \cdot 31^3 \cdot 223^3 z^7\theta(\theta + 1)(\theta + 2)(13828\theta + 24201) \\ & + 31^4 \cdot 223^4 z^8\theta(\theta + 1)(\theta + 2)(\theta + 3) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - z(2\theta + 1)(3459\theta^4 + 6918\theta^3 + 4879\theta^2 + 1420\theta + 145) \\ & + z^2(\theta + 1)(34575\theta^4 + 138300\theta^3 + 222873\theta^2 + 169146\theta + 50287) \\ & - 2z^3(\theta + 1)(\theta + 2)(2\theta + 3)(17285\theta^2 + 51855\theta + 47067) \\ & + z^4(\theta + 1)(\theta + 2)(\theta + 3)(69135\theta^2 + 276540\theta + 293201) \\ & - 17283 z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \\ & + 6913 z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \frac{(3k)!}{k!^3} \frac{(4k)!}{k!^2(2k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
95	$ \begin{aligned} D = & \theta^4 - z(1733\theta^4 + 3466\theta^3 + 2446\theta^2 + 713\theta + 73) \\ & + z^2(\theta + 1)^2(6922\theta^2 + 13844\theta + 8343) \\ & - z^3(\theta + 1)(\theta + 2)(10378\theta^2 + 31134\theta + 24925) \\ & + 6917z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & - 1729z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \frac{(3k)!}{k!^3} \frac{(4k)!}{k!^2(2k)!}$
96	$ \begin{aligned} D = & \theta^4 - z(1040\theta^4 + 2056\theta^3 + 1484\theta^2 + 456\theta + 50) \\ & + 4z^2(4120\theta^4 + 10264\theta^3 + 12062\theta^2 + 8396\theta + 2687) \\ & - 16z^3(6160\theta^4 + 18456\theta^3 + 25752\theta^2 + 17862\theta + 4081) \\ & + 16z^4(16400\theta^4 + 57376\theta^3 + 85016\theta^2 + 55560\theta + 13441) \\ & - 2^{14}z^5(\theta + 1)^2(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{2n}{2k}^{-1} \frac{(4k)!}{k!^4}$
97	$ \begin{aligned} D = & \theta^4 - z(16400\theta^4 + 4100\theta^3 + 902\theta^2 - 1148\theta - 623) \\ & + z^2(100925536\theta^4 + 50462768\theta^3 + 13232182\theta^2 + 847628\theta + 8796151) \\ & - z^3(276490092800\theta^4 + 207367569600\theta^3 + 63242834064\theta^2 \\ & \quad + 26362467316\theta - 3821932666) \\ & + z^4(285882691092736\theta^4 + 285882691092736\theta^3 + 100029707616352\theta^2 \\ & \quad + 15451954237456\theta + 1121739103233) \\ & - 2^{11}z^5(2211920742400\theta^4 + 2764900928000\theta^3 + 2006667953392\theta^2 \\ & \quad + 1165909352584\theta + 332723569061) \\ & + 2^{20}z^6(25836937216\theta^4 + 38755405824\theta^3 + 33098847360\theta^2 \\ & \quad + 13914543616\theta - 569939335) \\ & - 2^{33}z^7(8396800\theta^4 + 14694400\theta^3 + 12330304\theta^2 + 3444000\theta + 461113) \\ & + 2^{44}z^8(8\theta + 3)^2(8\theta + 5)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(2058\theta^5 + 5125\theta^4 + 5002\theta^3 + 2378\theta^2 + 549\theta + 49) \\ & + 2^2z^2(20520\theta^5 + 61480\theta^4 + 90950\theta^3 + 89906\theta^2 + 55877\theta + 15545) \\ & - 2^3z^3(82000\theta^5 + 286840\theta^4 + 507060\theta^3 + 532058\theta^2 + 277120\theta + 38601) \\ & + 2^4z^4(163920\theta^5 + 655520\theta^4 + 1259720\theta^3 + 1286224\theta^2 \\ & \quad + 611097\theta + 122882) \\ & - 2^5z^5(163872\theta^5 + 737360\theta^4 + 1464400\theta^3 + 1446952\theta^2 \\ & \quad + 709642\theta + 138241) \\ & + 2^{18}z^6(\theta + 1)^3(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A'_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}^2 \binom{2n}{2k}^{-1} \frac{(4k)!}{k!^2(2k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
98	$ \begin{aligned} D = & \theta^4 - z(6928\theta^4 + 1732\theta^3 + 414\theta^2 - 452\theta - 251) \\ & + z^2(18026592\theta^4 + 9013296\theta^3 + 2541558\theta^2 + 295932\theta + 1580631) \\ & - z^3(20926439680\theta^4 + 15694829760\theta^3 + 5119446672\theta^2 \\ & + 2221999476\theta - 262743618) \\ & + z^4(9248048087296\theta^4 + 9248048087296\theta^3 + 3470622495840\theta^2 \\ & + 668590646032\theta + 65658526849) \\ & - 2^4 \cdot 24z^5(376675914240\theta^4 + 470844892800\theta^3 + 348726894896\theta^2 \\ & + 202514342736\theta + 57000462817) \\ & + 2^6 \cdot 24^2z^6(23362463232\theta^4 + 35043694848\theta^3 + 30328003584\theta^2 \\ & + 12869932032\theta - 426340673) \\ & - 2^4 \cdot 24^5z^7(17957376\theta^4 + 31425408\theta^3 + 26673840\theta^2 \\ & + 7638120\theta + 1060093) \\ & + 24^8z^8(12\theta + 5)^2(12\theta + 7)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(874\theta^5 + 2165\theta^4 + 2146\theta^3 + 1054\theta^2 + 257\theta + 25) \\ & + 2^2z^2(8680\theta^5 + 25960\theta^4 + 38710\theta^3 + 38498\theta^2 + 23981\theta + 6689) \\ & - 2^3z^3(34640\theta^5 + 121080\theta^4 + 215220\theta^3 + 227034\theta^2 + 119160\theta + 17085) \\ & + 2^4z^4(69200\theta^5 + 276640\theta^4 + 533960\theta^3 + 548432\theta^2 + 263401\theta + 53762) \\ & - 2^5z^5(69152\theta^5 + 311120\theta^4 + 620240\theta^3 + 616936\theta^2 + 305866\theta + 60481) \\ & + 2^{14} \cdot 3z^6(\theta + 1)^3(6\theta + 5)(6\theta + 7) \end{aligned} $
	$A'_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}^2 \binom{2n}{2k}^{-1} \frac{(3k)!}{k!^3}$
99	$ \begin{aligned} D = & 13^2\theta^4 - z(59397\theta^4 + 117546\theta^3 + 86827\theta^2 + 28054\theta + 3380) \\ & + 2^4z^2(6386\theta^4 - 1774\theta^3 - 17898\theta^2 - 11596\theta - 2119) \\ & + 2^8z^3(67\theta^4 + 1248\theta^3 + 1091\theta^2 + 312\theta + 26) - 2^{12}z^4(2\theta + 1)^4 \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{n}{k}^2 \binom{2n+k}{n}$
100	$ \begin{aligned} D = & \theta^4 - z(73\theta^4 + 98\theta^3 + 77\theta^2 + 28\theta + 4) \\ & + z^2(520\theta^4 - 1040\theta^3 - 2904\theta^2 - 2048\theta - 480) \\ & + 2^6z^3(65\theta^4 + 390\theta^3 + 417\theta^2 + 180\theta + 28) \\ & - 2^9z^4(73\theta^4 + 194\theta^3 + 221\theta^2 + 124\theta + 28) + 2^{15}z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \left\{ \sum_{k=0}^n \binom{n}{k}^3 \right\}^2$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
101	$D = \theta^4 - z(124\theta^4 + 242\theta^3 + 187\theta^2 + 66\theta + 9)$ $+ z^2(123\theta^4 - 246\theta^3 - 787\theta^2 - 554\theta - 124)$ $+ z^3(123\theta^4 + 738\theta^3 + 689\theta^2 + 210\theta + 12)$ $- z^4(124\theta^4 + 254\theta^3 + 205\theta^2 + 78\theta + 12) + z^5(\theta + 1)^4$
	$A_n = \left\{ \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k} \right\}^2$
102	$D = \theta^4 - z(7\theta^2 + 7\theta + 2)(11\theta^2 + 11\theta + 3)$ $- z^2(1049\theta^4 + 4100\theta^3 + 5689\theta^2 + 3178\theta + 640)$ $+ 2^3 z^3(77\theta^4 - 462\theta^3 - 1420\theta^2 - 1053\theta - 252)$ $+ 2^4 z^4(1041\theta^4 + 2082\theta^3 - 1406\theta^2 - 2447\theta - 746)$ $+ 2^6 z^5(77\theta^4 + 770\theta^3 + 428\theta^2 - 93\theta - 80)$ $- 2^6 z^6(1049\theta^4 + 96\theta^3 - 317\theta^2 + 96\theta + 100)$ $- 2^9 z^7(7\theta^2 + 7\theta + 2)(11\theta^2 + 11\theta + 3) + 2^{12} z^8(\theta + 1)^4$
	$A_n = \sum_{k=0}^n \binom{n}{k}^3 \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}$
103	$D = \theta^4 - z(73\theta^4 + 200\theta^3 + 160\theta^2 + 60\theta + 9)$ $- z^2(738\theta^4 - 1476\theta^3 - 5274\theta^2 - 3816\theta - 918)$ $+ z^3(6642\theta^4 + 39852\theta^3 + 32238\theta^2 + 5832\theta - 1458)$ $+ 3^6 z^4(73\theta^4 + 92\theta^3 - 2\theta^2 - 48\theta - 18) - 3^{10} z^5(\theta + 1)^4$
	$A_n = \left\{ \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}^2 \right\}^2$
104	$D = \theta^4 - z(10\theta^2 + 10\theta + 3)(7\theta^2 + 7\theta + 2)$ $- z^2(71\theta^4 + 1148\theta^3 + 1591\theta^2 + 886\theta + 192)$ $- 2^3 \cdot 3^2 z^3(70\theta^4 - 420\theta^3 - 1289\theta^2 - 963\theta - 240)$ $- 2^4 \cdot 3^2 z^4(143\theta^4 + 286\theta^3 - 1138\theta^2 - 1281\theta - 414)$ $+ 2^6 \cdot 3^4 z^5(70\theta^4 + 700\theta^3 + 391\theta^2 - 75\theta - 76)$ $+ 2^6 \cdot 3^4 z^6(-71\theta^4 + 864\theta^3 + 1427\theta^2 + 864\theta + 180)$ $+ 2^9 \cdot 3^6 z^7(10\theta^2 + 10\theta + 3)(7\theta^2 + 7\theta + 2) + 2^{12} \cdot 3^8 z^8(\theta + 1)^4$
	$A_n = \sum_{k=0}^n \binom{n}{k}^3 \cdot \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$
105	$D = \theta^4 - 4z(3\theta^2 + 3\theta + 1)(7\theta^2 + 7\theta + 1)$ $+ 2^5 z^2(45\theta^4 + 84\theta^3 + 117\theta^2 + 66\theta + 12)$ $- 2^{10} z^3(21\theta^4 - 126\theta^3 - 386\theta^2 - 291\theta - 76)$ $+ 2^{14} z^4(37\theta^4 + 74\theta^3 + 50\theta^2 + 13\theta + 6)$ $+ 2^{18} z^5(21\theta^4 + 210\theta^3 + 118\theta^2 - 19\theta - 24)$ $+ 3 \cdot 2^{21} z^6(15\theta^4 + 32\theta^3 + 45\theta^2 + 32\theta + 8)$ $+ 2^{26} z^7(3\theta^2 + 3\theta + 1)(7\theta^2 + 7\theta + 1) + 2^{32} z^8(\theta + 1)^4$
	$A_n = \sum_{k=0}^n \binom{n}{k}^3 \cdot \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
106	$ \begin{aligned} D = & \theta^4 - 4z(3\theta^2 + 3\theta + 1)(11\theta^2 + 11\theta + 3) \\ & + 2^4 z^2(241\theta^4 + 940\theta^3 + 1303\theta^2 + 726\theta + 145) \\ & - 2^7 z^3(33\theta^4 - 198\theta^3 - 607\theta^2 - 456\theta - 117) \\ & + 2^{10} z^4(239\theta^4 + 478\theta^3 - 322\theta^2 - 561\theta - 169) \\ & + 2^{12} z^5(33\theta^4 + 330\theta^3 + 185\theta^2 - 32\theta - 37) \\ & + 2^{14} z^6(241\theta^4 + 24\theta^3 - 71\theta^2 + 24\theta + 23) \\ & + 2^{17} z^7(3\theta^2 + 3\theta + 1)(11\theta^2 + 11\theta + 3) + 2^{20} z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k} \cdot \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k}$
107	$ \begin{aligned} D = & \theta^4 - 2^4 z(3\theta^4 + 18\theta^3 + 15\theta^2 + 6\theta + 1) \\ & - 2^9 z^2(5\theta^4 - 10\theta^3 - 45\theta^2 - 34\theta - 9) \\ & + 2^{14} z^3(5\theta^4 + 30\theta^3 + 15\theta^2 - 6\theta - 5) \\ & + 2^{19} z^4(3\theta^4 - 6\theta^3 - 21\theta^2 - 18\theta - 5) - 2^{25} z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \left\{ \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \right\}^2$
108	$ \begin{aligned} D = & \theta^4 - 2z(6\theta^4 + 10\theta^3 + 10\theta^2 + 5\theta + 1) \\ & + 2^2 z^2(46671\theta^4 + 186674\theta^3 + 238539\theta^2 + 103735\theta + 14416) \\ & - 2^3 z^3(\theta + 1)(186644\theta^3 + 839888\theta^2 + 1223544\theta + 575489) \\ & + 2^4 z^4(\theta + 1)(\theta + 2)(279951\theta^2 + 1119788\theta + 1124983) \\ & - 2^6 z^5(\theta + 1)(\theta + 2)(\theta + 3)(93315\theta + 233287) \\ & + 2^6 \cdot 13 \cdot 37 \cdot 97 z^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_k (-1)^k 2^{n-2k} \binom{n}{2k} \left(\frac{(6k)!}{k!(2k)!(3k)!} \right)^2$
109	$ \begin{aligned} D = & 49\theta^4 - 2 \cdot 3z(8904\theta^4 + 17556\theta^3 + 12453\theta^2 + 3675\theta + 392) \\ & + 2^2 \cdot 3z^2(43704\theta^4 + 38088\theta^3 - 25757\theta^2 - 20608\theta - 3360) \\ & - 2^4 \cdot 3^3 z^3(2736\theta^4 - 1512\theta^3 - 1672\theta^2 - 357\theta - 14) \\ & - 2^6 \cdot 3^5 z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2) \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^{2n} \binom{n+k}{k} \binom{2n}{k}^2$
110	$ \begin{aligned} D = & \theta^4 - 12z(3\theta + 1)(3\theta + 2)(8\theta^2 + 8\theta + 3) \\ & + 2304z^2(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5) \end{aligned} $
	$A_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$
111	$ \begin{aligned} D = & \theta^4 - 16z(2\theta + 1)^2(8\theta^2 + 8\theta + 3) \\ & + 4096z^2(2\theta + 1)^2(2\theta + 3)^2 \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
112	$D = \theta^4 - 48z(6\theta + 1)(6\theta + 5)(8\theta^2 + 8\theta + 3)$ $+ 36864z^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11)$ $A_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$
113	$D = \theta^4 - z(10\theta^2 + 10\theta + 3)(11\theta^2 + 11\theta + 3)$ $+ z^2(1025\theta^4 + 3992\theta^3 + 5533\theta^2 + 3082\theta + 615)$ $+ 3^2 z^3(-110\theta^4 + 660\theta^3 + 2027\theta^2 + 1509\theta + 369)$ $+ 3^2 z^4(2032\theta^4 + 4064\theta^3 - 2726\theta^2 - 4758\theta - 1431)$ $+ 3^4 z^5(110\theta^4 + 1100\theta^3 + 613\theta^2 - 125\theta - 117)$ $+ 3^4 z^6(1025\theta^4 + 108\theta^3 - 293\theta^2 + 108\theta + 99)$ $+ 3^6 z^7(10\theta^2 + 10\theta + 3)(11\theta^2 + 11\theta + 3) + 3^8 z^8(\theta + 1)^4$ $A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k} \cdot \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$
114	$D = \theta^4 - 4z(7\theta^2 + 7\theta + 2)(8\theta^2 + 8\theta + 3)$ $+ 2^7 z^2(98\theta^4 + 200\theta^3 + 274\theta^2 + 148\theta + 23)$ $+ 2^{13} z^3(-56\theta^4 + 336\theta^3 + 1027\theta^2 + 783\theta + 216)$ $+ 2^{19} z^4(82\theta^4 + 164\theta^3 + 84\theta^2 + 2\theta + 11)$ $+ 2^{24} z^5(56\theta^4 + 560\theta^3 + 317\theta^2 - 39\theta - 68)$ $+ 2^{29} z^6(98\theta^4 + 192\theta^3 + 262\theta^2 + 192\theta + 47)$ $+ 2^{35} z^7(7\theta^2 + 7\theta + 2)(8\theta^2 + 8\theta + 3) + 2^{44} z^8(\theta + 1)^4$ $A_n = \sum_{k=0}^n \binom{n}{k}^3 \cdot \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$
115	$D = \theta^4 - 2^4 z(16\theta^4 + 128\theta^3 + 112\theta^2 + 48\theta + 9)$ $+ 2^{12} z^2(-32\theta^4 + 64\theta^3 + 304\theta^2 + 240\theta + 71)$ $+ 2^{20} z^3(32\theta^4 + 192\theta^3 + 80\theta^2 - 48\theta - 39)$ $+ 2^{28} z^4(16\theta^4 - 64\theta^3 - 176\theta^2 - 144\theta - 39) - 2^{40} z^5(\theta + 1)^4$ $A_n = \left\{ \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k} \right\}^2$
116	$D = \theta^4 - 2^5 z(10\theta^4 + 26\theta^3 + 20\theta^2 + 7\theta + 1)$ $+ 2^8 z^2(52\theta^4 + 472\theta^3 + 832\theta^2 + 492\theta + 103)$ $+ 2^{16} z^3(14\theta^4 + 12\theta^3 - 96\theta^2 - 105\theta - 29)$ $- 2^{18} z^4(2\theta + 1)(56\theta^3 + 468\theta^2 + 646\theta + 249)$ $- 2^{24} z^5(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$ $A_n = \binom{2n}{n}^2 \sum_{k=0}^n 4^{n-k} \binom{n}{k}^2 \binom{n+k}{k} \binom{2k}{k} \binom{2n}{2k}^{-1}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
117	$ \begin{aligned} D = & 3^2\theta^4 + 12z(256\theta^4 + 176\theta^3 + 133\theta^2 + 45\theta + 6) \\ & + 2^7z^2(2588\theta^4 + 1952\theta^3 + 584\theta^2 + 15\theta - 15) \\ & + 2^{12}z^3(3183\theta^4 + 2466\theta^3 + 1801\theta^2 + 711\theta + 111) \\ & + 2^{17} \cdot 7z^4(134\theta^4 + 250\theta^3 + 180\theta^2 + 55\theta + 5) \\ & - 2^{22} \cdot 7^2z^5(\theta + 1)^4 \end{aligned} $
	a formula for A_n is not known
118	$ \begin{aligned} D = & \theta^4 - z(465\theta^4 + 594\theta^3 + 431\theta^2 + 134\theta + 16) \\ & + 2^4z^2(2625\theta^4 + 1911\theta^3 - 946\theta^2 - 884\theta - 176) \\ & + 2^6z^3(-16105\theta^4 + 3624\theta^3 + 5241\theta^2 + 1284\theta + 36) \\ & - 2^{11} \cdot 7z^4(155\theta^4 + 334\theta^3 + 306\theta^2 + 139\theta + 26) + 2^{16} \cdot 7^2z^5(\theta + 1)^4 \end{aligned} $
	$A_n = 2^{-5n} \sum_{k=0}^n \binom{n}{k}^{-5} \binom{2k}{k}^5 \binom{2n-2k}{n-k}^5$
119	$ \begin{aligned} D = & 9\theta^4 - 12z(256\theta^4 + 320\theta^3 + 271\theta^2 + 111\theta + 18) \\ & + 2^7z^2(3104\theta^4 + 7040\theta^3 + 8012\theta^2 + 4452\theta + 927) \\ & - 2^{15}z^3(752\theta^4 + 2304\theta^3 + 3042\theta^2 + 1854\theta + 405) \\ & + 2^{21}z^4(2\theta + 1)(176\theta^3 + 552\theta^2 + 622\theta + 231) \\ & - 2^{31}z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!^2}$
120	$ \begin{aligned} D = & \theta^4 - 4z(8\theta^2 + 8\theta + 3)(10\theta^2 + 10\theta + 3) \\ & + 2^4z^2(1600\theta^4 + 8128\theta^3 + 11408\theta^2 + 6560\theta + 1473) \\ & + 2^{10} \cdot 3^2z^3(80\theta^4 - 480\theta^3 - 1466\theta^2 - 1122\theta - 315) \\ & - 2^{13} \cdot 3^2z^4(1744\theta^4 + 3488\theta^3 - 4256\theta^2 - 6000\theta - 2079) \\ & + 2^{18} \cdot 3^4z^5(80\theta^4 + 800\theta^3 + 454\theta^2 - 50\theta - 99) \\ & + 2^{20} \cdot 3^4z^6(1600\theta^4 - 1728\theta^3 - 3376\theta^2 - 1728\theta - 207) \\ & - 2^{26} \cdot 3^6z^7(8\theta^2 + 8\theta + 3)(10\theta^2 + 10\theta + 3) + 2^{32} \cdot 3^8z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \cdot \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$
121	$ \begin{aligned} D = & \theta^4 - 4z(8\theta^2 + 8\theta + 3)(11\theta^2 + 11\theta + 3) \\ & + 2^4z^2(1936\theta^4 + 7552\theta^3 + 10464\theta^2 + 5824\theta + 1159) \\ & + 2^{10}z^3(-88\theta^4 + 528\theta^3 + 1615\theta^2 + 1227\theta + 333) \\ & + 2^{13}z^4(1920\theta^4 + 3840\theta^3 - 2592\theta^2 - 4512\theta - 1353) \\ & + 2^{18}z^5(88\theta^4 + 880\theta^3 + 497\theta^2 - 67\theta - 105) \\ & + 2^{20}z^6(1936\theta^4 + 192\theta^3 - 576\theta^2 + 192\theta + 183) \\ & + 2^{26}z^7(8\theta^2 + 8\theta + 3)(11\theta^2 + 11\theta + 3) + 2^{30}z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k} \cdot \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
122	$ \begin{aligned} D = & \theta^4 - 2^4 z (3\theta^2 + 3\theta + 1)(8\theta^2 + 8\theta + 3) \\ & + 2^9 z^2 (72\theta^4 + 480\theta^3 + 680\theta^2 + 400\theta + 97) \\ & + 2^{17} z^3 (24\theta^4 - 144\theta^3 - 439\theta^2 - 339\theta - 99) \\ & + 2^{23} z^4 (-88\theta^4 - 176\theta^3 + 320\theta^2 + 408\theta + 151) \\ & + 2^{30} z^5 (24\theta^4 + 240\theta^3 + 137\theta^2 - 11\theta - 31) \\ & + 2^{35} z^6 (72\theta^4 - 192\theta^3 - 328\theta^2 - 192\theta - 31) \\ & - 2^{43} z^7 (3\theta^2 + 3\theta + 1)(8\theta^2 + 8\theta + 3) + 2^{52} z^8 (\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \cdot \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}$
123	$ \begin{aligned} D = & \theta^4 - 4z(3\theta^2 + 3\theta + 1)(10\theta^2 + 10\theta + 3) \\ & + 2^4 z^2 (209\theta^4 + 1052\theta^3 + 1471\theta^2 + 838\theta + 183) \\ & + 2^7 \cdot 3^2 z^3 (30\theta^4 - 180\theta^3 - 551\theta^2 - 417\theta - 111) \\ & - 2^{10} \cdot 3^2 z^4 (227\theta^4 + 454\theta^3 - 550\theta^2 - 777\theta - 261) \\ & + 2^{12} \cdot 3^4 z^5 (30\theta^4 + 300\theta^3 + 169\theta^2 - 25\theta - 35) \\ & + 2^{14} \cdot 3^4 z^6 (209\theta^4 - 216\theta^3 - 431\theta^2 - 216\theta - 27) \\ & - 2^{17} \cdot 3^6 z^7 (3\theta^2 + 3\theta + 1)(10\theta^2 + 10\theta + 3) + 2^{20} \cdot 3^8 z^8 (\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n+k}{k}^2 \binom{2k}{k} \cdot \sum_{k=0}^n 4^{n-k} \binom{n}{k}^2 \binom{2k}{k}$
124	$ \begin{aligned} D = & 61^2 \theta^4 - 61z(3029\theta^4 + 5572\theta^3 + 4677\theta^2 + 1891\theta + 305) \\ & + z^2(1215215\theta^4 + 3428132\theta^3 + 4267228\theta^2 + 2572675\theta + 611586) \\ & - 3^4 z^3(39370\theta^4 + 140178\theta^3 + 206807\theta^2 + 142191\theta + 37332) \\ & + 3^8 z^4(566\theta^4 + 2230\theta^3 + 3356\theta^2 + 2241\theta + 558) - 3^{13} z^5 (\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k,l} \binom{n}{k}^2 \binom{n}{l} \binom{k}{l} \binom{k+l}{k} \binom{2n-k-l}{n-k}$
125	$ \begin{aligned} D = & \theta^4 - z(11669\theta^4 + 23338\theta^3 + 15886\theta^2 + 4217\theta + 361) \\ & + z^2(\theta + 1)^2(4666\theta^2 + 93332\theta + 55095) \\ & - z^3(\theta + 1)(\theta + 2)(69994\theta^2 + 209982\theta + 167533) \\ & + 46661z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & - 11665z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \frac{(6k)!}{k!^4 (2k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
126	$ \begin{aligned} D = & \theta^4 - z(186656\theta^4 + 46676\theta^3 + 8434\theta^2 - 14904\theta - 7813) \\ & + z^2(13065919936\theta^4 + 6535199792\theta^3 + 1453643198\theta^2 \\ & - 105990748\theta + 1115889227) \\ & - z^3(406553346039296\theta^4 + 305071791609408\theta^3 + 80614299041712\theta^2 \\ & + 30335098270388\theta - 5908917544614) \\ & + z^4(4746509269839709696\theta^4 + 4751387282863861504\theta^3 \\ & + 1455612685803281696\theta^2 + 132330243679138384\theta \\ & - 845722995408775) \\ & - 2^4 \cdot 5 \cdot 2333z^5(4\theta - 1)(101638336509824\theta^3 + 228725452667088\theta^2 \\ & + 193462702609348\theta + 59317557294355) \\ & + 2^5 \cdot 5^2 \cdot 2333^2z^6(4\theta - 1)(4\theta + 3)(6532959968\theta^2 \\ & + 16333519832\theta + 11222488415) \\ & - 2^8 \cdot 5^3 \cdot 2333^3z^7(4\theta - 1)(4\theta + 3)(4\theta + 7)(46664\theta + 81665) \\ & + 2^8 \cdot 5^4 \cdot 2333^4z^8(4\theta - 1)(4\theta + 3)(4\theta + 7)(4\theta + 11) \end{aligned} $
	same $K(q)$ as in #82 the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(11669\theta^4 + 23338\theta^3 + 15886\theta^2 + 4217\theta + 361) \\ & + 4z^2(\theta + 1)(2\theta + 1)(2\theta + 3)(4666\theta^2 + 93332\theta + 55095) \\ & - 8z^3(2\theta + 1)(2\theta + 3)(2\theta + 5)(69994\theta^2 + 209982\theta + 167533) \\ & + 746576z^4(\theta + 2)(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \\ & - 373280z^5(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)(2\theta + 9) \end{aligned} $
	$A'_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \frac{(6k)!}{k!^4 (2k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
127	$ \begin{aligned} D = & \theta^4 - z(186640\theta^4 + 46660\theta^3 + 8430\theta^2 - 14900\theta - 7811) \\ & + z^2(13063680096\theta^4 + 6531840048\theta^3 + 1452582774\theta^2 \\ & \quad - 105150948\theta + 1116311607) \\ & - z^3(406448815694080\theta^4 + 304836611770560\theta^3 + 80522819145360\theta^2 \\ & \quad + 30399976916340\theta - 5858109970482) \\ & + z^4(4744882429422797056\theta^4 + 4744882429422797056\theta^3 \\ & \quad + 1452477726672583776\theta^2 + 134807094643795216\theta \\ & \quad + 727318797282433) \\ & - 2^7 \cdot 3^2 z^5(65844708142440960\theta^4 + 82305885178051200\theta^3 \\ & \quad + 57162742271218800\theta^2 + 33149511110202480\theta + 9595488553244821) \\ & + 2^{12} \cdot 3^8 z^6(16930529404416\theta^4 + 25395794106624\theta^3 + 21045353440896\theta^2 \\ & \quad + 8640860120064\theta - 514138329761) \\ & - 2^{19} \cdot 3^{14} z^7(483770880\theta^4 + 846599040\theta^3 + 692048880\theta^2 \\ & \quad + 182253960\theta + 22046293) \\ & + 2^{24} \cdot 3^{22} z^8(4\theta + 1)(4\theta + 3)(12\theta + 5)(12\theta + 7) \end{aligned} $
	same $K(q)$ as in #82
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 + z(46676\theta^5 + 116650\theta^4 + 110180\theta^3 + 48620\theta^2 + 9874\theta + 722) \\ & - z^2(93328\theta^5 + 2799520\theta^4 + 4069720\theta^3 + 3965960\theta^2 \\ & \quad + 2452340\theta + 678212) \\ & + z^3(7465600\theta^5 + 26128320\theta^4 + 45620640\theta^3 + 47278800\theta^2 \\ & \quad + 24206400\theta + 3155112) \\ & - 2^4 z^4(1866320\theta^5 + 7465120\theta^4 + 14204360\theta^3 + 14307920\theta^2 \\ & \quad + 6634825\theta + 1290242) \\ & + 2^5 z^5(1866272\theta^5 + 8398160\theta^4 + 16537040\theta^3 + 16096360\theta^2 \\ & \quad + 7704010\theta + 1451521) \\ & - 4608z^6(\theta + 1)(3\theta + 2)(3\theta + 4)(6\theta + 5)(6\theta + 7) \end{aligned} $
	$A'_n = \sum_{k=0}^n \binom{2n-2k}{n-k} \frac{(6k)!}{k!^6}$
128	$ \begin{aligned} D = & \theta^4 - z(3145\theta^4 + 6282\theta^3 + 4403\theta^2 + 1262\theta + 122) \\ & + 4z^2(2\theta + 1)(6270\theta^3 + 18804\theta^2 + 20058\theta + 7523) \\ & - 8z^3(2\theta + 1)(2\theta + 3)(9395\theta^2 + 28181\theta + 22544) \\ & + 16z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(6260\theta + 2519) \\ & - 50064z^5(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	same $K(q)$ as in #1
	$A_n = \binom{2n}{n} \sum_{k=0}^n \binom{n}{k} \frac{(5k)!}{k!^3(2k)!}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
129	$ \begin{aligned} D = & \theta^4 - 2 \cdot 3z\theta(4\theta + 1)(6\theta^2 + 1) + 2^2 \cdot 3(3z)^2\theta(92\theta^3 + 46\theta^2 + 33\theta + 9) \\ & - 2^4(3z)^3\theta(1012\theta^3 + 759\theta^2 + 583\theta + 171) \\ & + 2^4 \cdot 7(3z)^4\theta(1518\theta^3 + 1518\theta^2 + 1243\theta + 388) \\ & - 2^5 \cdot 3 \cdot 7(3z)^5\theta(2024\theta^3 + 2530\theta^2 + 2200\theta + 725) \\ & + 2^5(3z)^6(269200\theta^4 + 403800\theta^3 + 371615\theta^2 + 128340\theta - 434) \\ & - 2^6(3z)^7(692352\theta^4 + 1211616\theta^3 + 1176528\theta^2 + 422499\theta - 6076) \\ & + 2^7(3z)^8(1472166\theta^4 + 2944332\theta^3 + 3008574\theta^2 + 1113993\theta - 39494) \\ & - 2^8(3z)^9(2621536\theta^4 + 5898456\theta^3 + 6326976\theta^2 + 2395311\theta - 157976) \\ & + 2^9 \cdot 7(3z)^{10}(563856\theta^4 + 1409640\theta^3 + 1583800\theta^2 + 608575\theta - 62062) \\ & - 2^{10} \cdot 7(3z)^{11}(722976\theta^4 + 1988184\theta^3 + 2335176\theta^2 + 906129\theta - 124124) \\ & + 2^8(3z)^{12}(44454688\theta^4 + 133364064\theta^3 + 163448032\theta^2 \\ & + 63949536\theta - 10415783) \\ & - 2^{12}(3z)^{13}(5247024\theta^4 + 17052828\theta^3 + 21770956\theta^2 \\ & + 8611822\theta - 1478855) \\ & + 2^{12}(3z)^{14}(8546736\theta^4 + 29913576\theta^3 + 39718716\theta^2 \\ & + 15991626\theta - 2532173) \\ & - 2^{14}(3z)^{15}(3006608\theta^4 + 11274780\theta^3 + 15546100\theta^2 \\ & + 6429885\theta - 794437) \\ & + 2^{13}(3z)^{16}(7307784\theta^4 + 29231136\theta^3 + 41794464\theta^2 \\ & + 17919876\theta - 1365581) \\ & - 2^{16}(3z)^{17}(956304\theta^4 + 4064292\theta^3 + 6017616\theta^2 + 2691333\theta - 83545) \\ & + 2^{17}(3z)^{18}(428800\theta^4 + 1929600\theta^3 + 2954715\theta^2 + 1380690\theta - 2429) \\ & - 2^{18}(3z)^{19}(163104\theta^4 + 774744\theta^3 + 1225444\theta^2 + 596629\theta + 4256) \\ & + 2^{16}(3z)^{20}(414336\theta^4 + 2071680\theta^3 + 3381120\theta^2 + 1704960\theta + 5957) \\ & - 2^{21}(3z)^{21}\theta(6688\theta^3 + 35112\theta^2 + 59068\theta + 30597) \\ & + 2^{23}(3z)^{22}\theta(\theta + 1)(672\theta^2 + 3024\theta + 3379) \\ & - 2^{27} \cdot 3(3z)^{23}\theta(\theta + 1)(\theta + 2)(4\theta + 11) + 2^{28}(3z)^{24}\theta(\theta + 1)(\theta + 2)(\theta + 3) \end{aligned} $ the pullback of the 5th-order differential equation $D'y = 0$, where $ \begin{aligned} D' = & \theta^5 - 6z\theta(\theta^2 + \theta + 1)(3\theta^2 + 3\theta + 1) \\ & + (6z)^2(\theta + 1)(15\theta^4 + 60\theta^3 + 105\theta^2 + 90\theta + 31) \\ & - 10(6z)^3(\theta + 1)(\theta + 2)(2\theta + 3)(\theta^2 + 3\theta + 3) \\ & + 5(6z)^4(\theta + 1)(\theta + 2)(\theta + 3)(3\theta^2 + 12\theta + 13) \\ & - 3(6z)^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5) \\ & + 2(6z)^6(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5) \end{aligned} $ $ A'_n = \sum_{k=0}^n (-1)^k 6^{n-6k} \binom{n}{6k} \frac{(6k)!}{k!^6} $

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
130	$ \begin{aligned} D = & \theta^4 - z(224\theta^4 + 56\theta^3 + 28\theta^2 - 3) \\ & + z^2(21952\theta^4 + 10976\theta^3 + 5998\theta^2 + 1020\theta + 261) \\ & - z^3(1238528\theta^4 + 928896\theta^3 + 551280\theta^2 + 139800\theta + 29316) \\ & + z^4(44574208\theta^4 + 44574208\theta^3 + 28575008\theta^2 + 8415456\theta + 1270689) \\ & - 2^4 z^5(67010048\theta^4 + 83762560\theta^3 + 57732992\theta^2 + 18330336\theta + 2276415) \\ & + 2^5 z^6(549092864\theta^4 + 823639296\theta^3 + 607886736\theta^2 \\ & \quad + 202347456\theta + 21851403) \\ & - 2^8 z^7(767502848\theta^4 + 1343129984\theta^3 + 1057767088\theta^2 \\ & \quad + 363949656\theta + 34797501) \\ & + 2^8 z^8(5778481408\theta^4 + 11556962816\theta^3 + 9682036960\theta^2 \\ & \quad + 3432553440\theta + 350349201) \\ & - 2^{13} \cdot 3^2 z^9(98832896\theta^4 + 222374016\theta^3 + 197614416\theta^2 \\ & \quad + 73072472\theta + 9376113) \\ & + 2^{16} \cdot 3^4 z^{10}(4204032\theta^4 + 10510080\theta^3 + 9877952\theta^2 + 3874976\theta + 592815) \\ & - 2^{21} \cdot 3^6 z^{11}(4\theta + 1)(6272\theta^3 + 15680\theta^2 + 13172\theta + 3783) \\ & + 2^{24} \cdot 3^8 z^{12}(4\theta + 1)(4\theta + 3)^2(4\theta + 5) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(14\theta^4 + 28\theta^3 + 28\theta^2 + 14\theta + 3) \\ & + 4z^2(\theta + 1)^3(196\theta^2 + 392\theta + 255) - 1152z^3(\theta + 1)^2(\theta + 2)^2(2\theta + 3) \end{aligned} $
	$A'_n = \sum_{i+j+k+l+m+s=n} \left(\frac{n!}{i!j!k!l!m!s!} \right)^2$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
131	$D = \theta^4 - 5z\theta(24\theta^3 + 6\theta^2 + 4\theta + 1) + \dots$ $+ 3^4 \cdot 5^{24} z^{24} \theta(\theta + 1)(\theta + 2)(\theta + 4)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 5z(2\theta + 1)(\theta^2 + \theta + 1)(3\theta^2 + 3\theta + 1)$ $+ 5^2 z^2 (\theta + 1)(15\theta^4 + 60\theta^3 + 105\theta^2 + 90\theta + 31)$ $- 2 \cdot 5^4 z^3 (\theta + 1)(\theta + 2)(2\theta + 3)(\theta^2 + 3\theta + 3)$ $+ 5^5 z^4 (\theta + 1)(\theta + 2)(\theta + 3)(3\theta^2 + 12\theta + 12)$ $- 5^5 z^5 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(2\theta + 5)$ $- 3 \cdot 5^6 z^6 (\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)(\theta + 5)$
	$A'_n = \sum_{k=0}^n (-1)^k 5^{n-5k} \binom{2k}{k} \binom{n}{5k} \frac{(5k)!}{k!^5}$
132	$D = \theta^4 - 5z(80\theta^4 + 20\theta^3 + 10\theta^2 - 1) + \dots$ $+ 2^{30} \cdot 5^{20} z^{16} \theta(\theta + 2)(2\theta + 1)(2\theta + 3)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 10z(2\theta + 1)(5\theta^4 + 10\theta^3 + 10\theta^2 + 5\theta + 1)$ $+ 500z^2 (\theta + 1)(2\theta + 1)(2\theta + 3)(2\theta^2 + 4\theta + 3)$ $- 5000z^3 (2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta^2 + 6\theta + 5)$ $+ 50000z^4 (\theta + 2)(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)$
	$A'_n = \binom{2n}{n} \sum_{k=0}^n (-1)^k 5^{n-5k} \binom{n}{5k} \frac{(5k)!}{k!^5}$
cases #133–143 are described in [1], Section 7	

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
144	$D = \theta^4 - z(73\theta^4 + 578\theta^3 + 493\theta^2 + 204\theta + 36)$ $+ 2^3 \cdot 3^2 z^2(-145\theta^4 + 290\theta^3 + 1387\theta^2 + 1072\theta + 300)$ $+ 2^6 \cdot 3^4 z^3(145\theta^4 + 870\theta^3 + 353\theta^2 - 252\theta - 180)$ $- 2^9 \cdot 3^6 z^4(-73\theta^4 + 286\theta^3 + 803\theta^2 + 660\theta + 180) - 2^{15} \cdot 3^{10} z^5(\theta + 1)^4$ $\sum_{n=0}^{\infty} A_n z^n = (g) * (g) \quad (\text{see [1], Section 7})$
145	$D = \theta^4 - 3^2 z(81\theta^4 + 648\theta^3 + 576\theta^2 + 252\theta + 49)$ $+ 2 \cdot 3^8 z^2(-81\theta^4 + 162\theta^3 + 765\theta^2 + 612\theta + 187)$ $+ 2 \cdot 3^{14} z^3(81\theta^4 + 486\theta^3 + 207\theta^2 - 108\theta - 97)$ $+ 3^{20} z^4(81\theta^4 - 324\theta^3 - 882\theta^2 - 720\theta - 194) - 3^{30} z^5(\theta + 1)^4$ $\sum_{n=0}^{\infty} A_n z^n = (h) * (h) \quad (\text{see [1], Section 7})$
146	$D = \theta^4 - 4z(5\theta^4 + 10\theta^3 + 10\theta^2 + 5\theta + 1)$ $+ 80z^2(\theta + 1)^2(2\theta^2 + 4\theta + 3) - 320z^3(\theta + 1)(\theta + 2)(2\theta^2 + 6\theta + 5)$ $+ 2304z^4(\theta + 1)(\theta + 2)^2(\theta + 3) - 5120z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4)$ $A_n = \sum_{k=0}^n (-1)^k 4^{n-4k} \binom{2k}{k} \binom{n}{4k} \frac{(4k)!}{k!^4}$
147	$D = \theta^4 - 8z(2\theta + 1)^2(2\theta^2 + 2\theta + 1)$ $+ 64z^2(2\theta + 1)(2\theta + 3)(6\theta^2 + 12\theta + 7)$ $- 1024z^3(2\theta + 1)(2\theta + 3)^2(2\theta + 5)$ $+ 8192z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)$ $A_n = \binom{2n}{n} \sum_{k=0}^n (-1)^k 4^{n-4k} \binom{n}{4k} \frac{(4k)!}{k!^4}$
148	$D = \theta^4 - 5z(5\theta^4 + 10\theta^3 + 10\theta^2 + 5\theta + 1) + 125z^2(\theta + 1)^2(2\theta^2 + 4\theta + 3)$ $- 625z^3(\theta + 1)(\theta + 2)(2\theta^2 + 6\theta + 5) - 3125z^4(\theta + 1)(\theta + 2)^2(\theta + 3)$ $A_n = \sum_{k=0}^n (-1)^k 5^{n-5k} \binom{n}{5k} \frac{(5k)!}{k!^5}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
149	$ \begin{aligned} D = & \theta^4 - 2^2 3z(4896\theta^4 + 1224\theta^3 + 265\theta^2 - 347\theta - 186) \\ & + 2^6 3^4 z^2(249840\theta^4 + 124920\theta^3 + 32255\theta^2 + 1619\theta + 21824) \\ & - 2^{10} 3^7 z^3(5674464\theta^4 + 4255848\theta^3 + 1276509\theta^2 + 519897\theta - 79028) \\ & + 2^{14} 3^{10} z^4(48607704\theta^4 + 486077040\theta^3 + 16615003\theta^2 \\ & + 2178895\theta + 94752) \\ & - 2^{18} 3^{13} z^5(5674464\theta^4 + 7093080\theta^3 + 2892155\theta^2 - 369553\theta - 546818) \\ & + 2^{22} 3^{16} z^6(249840\theta^4 + 374760\theta^3 + 174525\theta^2 + 1689\theta + 14384) \\ & - 2^{26} 3^{19} z^7\theta(4896\theta^3 + 8568\theta^3 + 4447\theta + 595) \\ & + 2^{30} 3^{22} z^8\theta(\theta + 1)(6\theta + 1)(6\theta + 5) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 12z(2\theta + 1)(6\theta + 1)(6\theta + 5)(17\theta^2 + 17\theta + 5) \\ & + 144z^2(\theta + 1)(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11) \end{aligned} $
	$A'_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$
150	$ \begin{aligned} D = & \theta^4 - 2^2 z(224\theta^4 + 56\theta^3 + 19\theta^2 - 9\theta - 6) \\ & + 2^6 z^2(6000\theta^4 + 3000\theta^3 + 1139\theta^2 - 537\theta - 420) \\ & - 2^{10} z^3(98336\theta^4 + 73752\theta^3 + 31007\theta^2 - 5469\theta + 1524) \\ & + 2^{14} z^4(1073176\theta^4 + 1073176\theta^3 + 495055\theta^2 + 34851\theta + 112248) \\ & - 2^{18} 3^4 z^5(98336\theta^4 + 122920\theta^3 + 61737\theta^2 + 13173\theta + 6210) \\ & + 2^{18} 3^8 z^6(96000\theta^4 + 144000\theta^3 + 78224\theta^2 + 19792\theta - 576) \\ & - 2^{26} 3^{12} z^7\theta(224\theta^3 + 392\theta^2 + 229\theta + 49) + 2^{30} 3^{16} z^8\theta(\theta + 1)(2\theta + 1)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 4z(2\theta + 1)^3(7\theta^2 + 7\theta + 3) + 1296z^2(\theta + 1)(2\theta + 1)^2(2\theta + 3)^2$
	$A'_n = \binom{2n}{n}^2 \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{n+k}{k} \frac{(3k)!}{k!^3}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
151	$ \begin{aligned} D = & \theta^4 - 3z(504\theta^4 + 126\theta^3 + 41\theta^2 - 22\theta - 14) \\ & + 3^2z^2(121500\theta^4 + 60750\theta^3 + 22221\theta^2 - 11790\theta - 8762) \\ & - 3^5z^3(1991304\theta^4 + 1493478\theta^3 + 607149\theta^2 - 133506\theta + 26716) \\ & + 3^8z^4(21731814\theta^4 + 21731814\theta^3 + 9723033\theta^2 + 378126\theta + 2244916) \\ & - 3^{15}z^5(1991304\theta^4 + 2489130\theta^3 + 1215603\theta^2 + 230166\theta + 125122) \\ & + 3^{22}z^6(121500\theta^4 + 182250\theta^3 + 96471\theta^2 + 22446\theta - 698) \\ & - 3^{31}z^7\theta(504\theta^3 + 882\theta^2 + 503\theta + 98) + 3^{38}z^8\theta(\theta + 1)(3\theta + 1)(3\theta + 2) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 3z(2\theta + 1)(3\theta + 1)(3\theta + 2)(7\theta^2 + 7\theta + 3) \\ & + 729z^2(\theta + 1)(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5) \end{aligned} $
	$A'_n = \frac{(3n)!}{n!^3} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{n+k}{k} \frac{(3k)!}{k!^3}$
152	$ \begin{aligned} D = & \theta^4 - 2^2z(896\theta^4 + 224\theta^3 + 69\theta^2 - 43\theta - 26) \\ & + 2^7z^2(48000\theta^4 + 24000\theta^3 + 8362\theta^2 - 5110\theta - 3589) \\ & - 2^{14}z^3(393344\theta^4 + 295008\theta^3 + 114809\theta^2 - 31991\theta + 4246) \\ & + 2^{20}z^4(4292704\theta^4 + 4292704\theta^3 + 1846073\theta^2 - 6199\theta + 436401) \\ & - 2^{26}3^4z^5(393344\theta^4 + 491680\theta^3 + 231583\theta^2 + 36431\theta + 24552) \\ & + 2^{31}3^8z^6(48000\theta^4 + 72000\theta^3 + 36862\theta^2 + 7582\theta - 261) \\ & - 2^{38}3^{12}z^7\theta(896\theta^3 + 1568\theta^2 + 867\theta + 147) \\ & + 2^{44}3^{16}z^8\theta(\theta + 1)(4\theta + 1)(4\theta + 3) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 4z(2\theta + 1)(4\theta + 1)(4\theta + 3)(7\theta^2 + 7\theta + 3) \\ & + 1296z^2(\theta + 1)(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7) \end{aligned} $
	$A'_n = \frac{(4n)!}{n!^2(2n)!} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{n+k}{k} \frac{(3k)!}{k!^3}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
153	$ \begin{aligned} D = & \theta^4 - 2^2 \cdot 3z(2016\theta^4 + 504\theta^3 + 143\theta^2 - 109\theta - 62) \\ & + 2^6 3^2 z^2(486000\theta^4 + 243000\theta^3 + 78759\theta^2 - 58149\theta - 38156) \\ & - 2^{10} 3^5 z^3(7965216\theta^4 + 5973912\theta^3 + 2179683\theta^2 - 807129\theta + 56452) \\ & + 2^{14} 3^8 z^4(86927256\theta^4 + 86927256\theta^3 + 35270163\theta^2 \\ & - 2418777\theta + 8633800) \\ & - 2^{18} 3^{15} z^5(7965216\theta^4 + 9956520\theta^3 + 4447557\theta^2 + 481617\theta + 492250) \\ & + 2^{22} 3^{21} z^6(1458000\theta^4 + 2187000\theta^3 + 1066527\theta^2 + 175635\theta - 7332) \\ & - 2^{26} 3^{31} z^7\theta(2016\theta^3 + 3528\theta^2 + 1865\theta + 245) \\ & + 2^{30} 3^{38} z^8\theta(\theta + 1)(6\theta + 1)(6\theta + 5) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 12z(2\theta + 1)(6\theta + 1)(6\theta + 5)(7\theta^2 + 7\theta + 3) \\ & + 11664z^2(\theta + 1)(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11) \end{aligned} $
	$A'_n = \frac{(6n)!}{n!(2n)!(3n)!} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{n+k}{k} \frac{(3k)!}{k!^3}$
154	$ \begin{aligned} D = & \theta^4 + 2^2 3z(36\theta^4 - 144\theta^3 - 99\theta^2 - 27\theta - 3) \\ & + 2^4 3^5 z^2(312\theta^2 + 156\theta + 25) \\ & + 2^{10} 3^8 z^3(12\theta^4 + 72\theta^3 + 57\theta^2 + 18\theta + 2) + 2^{12} 3^{12} z^4(2\theta + 1)^4 \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_{k=0}^n (-1)^k 3^{2n-3k} \binom{2n}{3k} \frac{(3k)!}{k!^3}$
155	$ \begin{aligned} D = & \theta^4 - 2^4 z(256\theta^4 + 2048\theta^3 + 1856\theta^2 + 832\theta + 169) \\ & - 2^{16} z^2(512\theta^4 - 1024\theta^3 - 4800\theta^2 - 3904\theta - 1239) \\ & + 2^{28} z^3(512\theta^4 + 3072\theta^3 + 1344\theta^2 - 576\theta - 599) \\ & + 2^{40} z^4(256\theta^4 - 1024\theta^3 - 2752\theta^2 - 2240\theta - 599) - 2^{60} z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \left\{ 64^n \sum_k (-1)^k \binom{-3/4}{k} \binom{-1/4}{n-k}^2 \right\}^2$
156	$ \begin{aligned} D = & \theta^4 - 6z(2\theta + 1)^2(2\theta^2 + 2\theta + 1) \\ & + 36z^2(2\theta + 1)(2\theta + 3)(6\theta^2 + 12\theta + 7) \\ & - 3888z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{2k}{k} \frac{(3k)!}{k!^3}$
157	$ \begin{aligned} D = & \theta^4 - 3z(5\theta^4 + 10\theta^3 + 10\theta^2 + 3\theta + 1) + 45z^2(\theta + 1)^2(2\theta^2 + 4\theta + 3) \\ & + 27z^3(\theta + 1)(\theta + 2)(6\theta^2 + 18\theta + 11) - 2187z^4(\theta + 1)(\theta + 2)^2(\theta + 3) \\ & + 3645z^5(\theta + 1)(\theta + 2)(\theta + 3)(\theta + 4) \end{aligned} $
	$A_n = \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{2k}{k}^2 \frac{(3k)!}{k!^3}$

#	differential operator D and coefficients $A_n, n = 0, 1, 2, \dots$
158	$ \begin{aligned} D = & \theta^4 - 2 \cdot 3z(12\theta^4 + 16\theta^3 + 14\theta^2 + 6\theta + 1) \\ & + 2^2 3^2 z^2 (60\theta^4 + 160\theta^3 + 198\theta^2 + 116\theta + 25) \\ & - 2^2 3^3 z^3 (2\theta + 1)(106\theta^3 + 263\theta^2 + 231\theta + 86) \\ & - 2^4 3^4 z^4 (2\theta + 1)(2\theta + 3)(21\theta^2 + 178\theta + 249) \\ & + 2^4 3^5 z^5 (2\theta + 1)(2\theta + 3)(2\theta + 5)(114\theta + 355) \\ & - 2^6 3^6 23z^6 (2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \binom{3k}{k} \frac{(3k)!}{k!^3}$
159	$ \begin{aligned} D = & \theta^4 - 2 \cdot 5z(14\theta^4 + 16\theta^3 + 14\theta^2 + 6\theta + 1) \\ & + 2^2 5^2 z^2 (84\theta^4 + 192\theta^3 + 226\theta^2 + 128\theta + 27) \\ & - 2^4 5^3 z^3 (140\theta^4 + 480\theta^3 + 710\theta^2 + 480\theta + 114) \\ & + 2^4 5^5 z^4 (2\theta + 1)(56\theta^3 + 228\theta^2 + 342\theta + 181) \\ & - 2^4 5^5 z^5 (2\theta + 1)(2\theta + 3)(332\theta^2 + 1216\theta + 1189) \\ & + 2^7 5^6 z^6 (2\theta + 1)(2\theta + 3)(2\theta + 5)(26\theta + 57) \\ & - 2^8 3 \cdot 5^7 z^7 (2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_{k=0}^n (-1)^k 5^{n-5k} \binom{n}{5k} \binom{2k}{k}^{-1} \frac{(5k)!}{k!^5}$
160	$ \begin{aligned} D = & \theta^4 - 3z(3\theta^2 + 3\theta + 1)(7\theta^2 + 7\theta + 3) \\ & + 3^2 z^2 (171\theta^4 + 396\theta^3 + 555\theta^2 + 318\theta + 64) \\ & + 2^3 3^4 z^3 (-21\theta^4 + 126\theta^3 + 386\theta^2 + 291\theta + 76) \\ & + 2^4 3^5 z^4 (147\theta^4 + 294\theta^3 + 102\theta^2 - 45\theta - 14) \\ & + 2^6 3^7 z^5 (21\theta^4 + 210\theta^3 + 118\theta^2 - 19\theta - 24) \\ & + 2^6 3^8 z^6 (171\theta^4 + 288\theta^3 + 393\theta^2 + 288\theta + 76) \\ & + 2^9 3^{10} z^7 (3\theta^2 + 3\theta + 1)(7\theta^2 + 7\theta + 3) + 2^{12} 3^{12} z^8 (\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^3 \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \frac{(3k)!}{k!^3}$
161	$ \begin{aligned} D = & \theta^4 - 3z(3\theta^2 + 3\theta + 1)(11\theta^2 + 11\theta + 3) \\ & + 3^2 z^2 (366\theta^4 + 1428\theta^3 + 1980\theta^2 + 1104\theta + 221) \\ & + 3^4 z^3 (-33\theta^4 + 198\theta^3 + 607\theta^2 + 456\theta + 117) \\ & + 3^5 z^4 (726\theta^4 + 1452\theta^3 - 978\theta^2 - 1704\theta - 515) \\ & + 3^7 z^5 (33\theta^4 + 330\theta^3 + 185\theta^2 - 32\theta - 37) \\ & + 3^8 z^6 (366\theta^4 + 36\theta^3 - 108\theta^2 + 36\theta + 35) \\ & + 3^{10} z^7 (3\theta^2 + 3\theta + 1)(11\theta^2 + 11\theta + 3) + 3^{12} z^8 (\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \frac{(3k)!}{k!^3}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
162	$ \begin{aligned} D = & \theta^4 - 3z(3\theta^2 + 3\theta + 1)(10\theta^2 + 10\theta + 3) \\ & + 3^3 z^2(91\theta^4 + 472\theta^3 + 659\theta^2 + 374\theta + 81) \\ & + 3^6 z^3(30\theta^4 - 180\theta^3 - 551\theta^2 - 417\theta - 111) \\ & + 3^8 z^4(-200\theta^4 - 400\theta^3 + 514\theta^2 + 714\theta + 237) \\ & + 3^{11} z^5(30\theta^4 + 300\theta^3 + 169\theta^2 - 25\theta - 35) \\ & + 3^{13} z^6(91\theta^4 - 108\theta^3 - 211\theta^2 - 108\theta - 15) \\ & - 3^{16} z^7(3\theta^2 + 3\theta + 1)(10\theta^2 + 10\theta + 3) + 3^{20} z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \frac{(3k)!}{k!^3}$
163	$ \begin{aligned} D = & \theta^4 - 2^2 3z(3\theta^2 + 3\theta + 1)^2 \\ & + 2^4 3^2 z^2(21\theta^4 + 156\theta^3 + 219\theta^2 + 126\theta + 29) \\ & + 2^7 3^4 z^3(3\theta^2 + 3\theta + 1)(3\theta^2 - 21\theta - 35) \\ & + 2^{10} 3^5 z^4(-27\theta^4 - 54\theta^3 + 114\theta^2 + 141\theta + 49) \\ & + 2^{12} 3^7 z^5(3\theta^2 + 3\theta + 1)(3\theta^2 + 27\theta - 11) \\ & + 2^{14} 3^8 z^6(21\theta^4 - 72\theta^3 - 123\theta^2 - 72\theta - 13) \\ & - 2^{17} 3^{10} z^7(3\theta^2 + 3\theta + 1)^2 + 2^{20} 3^{12} z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \cdot \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \frac{(3k)!}{k!^3}$
164	$ \begin{aligned} D = & \theta^4 - 2^2 3z(3\theta^2 + 3\theta + 1)(8\theta^2 + 8\theta + 3) \\ & + 2^4 3^2 z^2(144\theta^4 + 1152\theta^3 + 1632\theta^2 + 960\theta + 235) \\ & + 2^{10} 3^4 z^3(24\theta^4 - 144\theta^3 - 439\theta^2 - 339\theta - 99) \\ & + 2^{13} 3^5 z^4(-192\theta^4 - 384\theta^3 + 864\theta^2 + 1056\theta + 389) \\ & + 2^{18} 3^7 z^5(24\theta^4 + 240\theta^3 + 137\theta^2 - 11\theta - 31) \\ & + 2^{20} 3^8 z^6(144\theta^4 - 576\theta^3 - 960\theta^2 - 576\theta - 101) \\ & - 2^{26} 3^{10} z^7(3\theta^2 + 3\theta + 1)(8\theta^2 + 8\theta + 3) + 2^{32} 3^{12} z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{k=0}^n 4^{n-k} \binom{2k}{k}^2 \binom{2n-2k}{n-k} \cdot \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \frac{(3k)!}{k!^3}$
165	$ \begin{aligned} D = & \theta^4 - 3^2 z(3\theta + 1)(6\theta^2 + 3\theta + 1) \\ & + 3^5 z^2(-3\theta^4 + 6\theta^3 + 39\theta^2 + 30\theta + 8) \\ & + 3^8 z^3(3\theta^4 + 18\theta^3 - 3\theta^2 - 18\theta - 8) \\ & - 3^{11} z^4(3\theta + 2)(6\theta^2 + 9\theta + 4) - 3^{15} z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \left\{ \sum_{k=0}^n (-1)^k 3^{n-3k} \binom{n}{3k} \frac{(3k)!}{k!^3} \right\}^2$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
166	$ \begin{aligned} D = & \theta^4 - 2^4 3^2 z (1296\theta^4 + 10368\theta^3 + 9648\theta^2 + 4464\theta + 961) \\ & - 2^{12} 3^8 z^2 (2592\theta^4 - 5184\theta^3 - 24048\theta^2 - 20016\theta - 6671) \\ & + 2^{20} 3^{14} z^3 (2592\theta^4 + 15552\theta^3 + 7056\theta^2 - 2160\theta - 2927) \\ & + 2^{28} 3^{20} z^4 (1296\theta^4 - 5184\theta^3 - 13680\theta^2 - 11088\theta - 2927) \\ & - 2^{40} 3^{30} z^5 (\theta + 1)^4 \end{aligned} $
	$A_n = \left\{ 432^n \sum_k (-1)^k \binom{-5/6}{k} \binom{-1/6}{n-k}^2 \right\}^2$
167	$ \begin{aligned} D = & \theta^4 - 3z(7\theta^2 + 7\theta + 2)(18\theta^2 + 18\theta + 7) \\ & + 3^2 z^2 (3969\theta^4 + 8100\theta^3 + 11025\theta^2 + 5850\theta + 832) \\ & + 2^3 3^7 z^3 (-126\theta^4 + 756\theta^3 + 2309\theta^2 + 1767\theta + 496) \\ & + 2^4 3^8 z^4 (3321\theta^4 + 6642\theta^3 + 3330\theta^2 + 9\theta + 526) \\ & + 2^6 3^{13} z^5 (126\theta^4 + 1260\theta^3 + 715\theta^2 - 79\theta - 156) \\ & + 2^6 3^{14} z^6 (3969\theta^4 + 7776\theta^3 + 10593\theta^2 + 7776\theta + 1876) \\ & + 2^9 3^{19} z^7 (7\theta^2 + 7\theta + 2)(18\theta^2 + 18\theta + 7) + 2^{12} 3^{24} z^8 (\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (a) * (h) \quad (\text{see [1], Section 7})$
168	$ \begin{aligned} D = & \theta^4 - 3z(11\theta^2 + 11\theta + 3)(18\theta^2 + 18\theta + 7) \\ & + 3^2 z^2 (9801\theta^4 + 38232\theta^3 + 52965\theta^2 + 29466\theta + 5855) \\ & + 3^7 z^3 (-198\theta^4 + 1188\theta^3 + 3631\theta^2 + 2769\theta + 765) \\ & + 3^8 z^4 (19440\theta^4 + 38880\theta^3 - 26262\theta^2 - 45702\theta - 13679) \\ & + 3^{13} z^5 (198\theta^4 + 1980\theta^3 + 1121\theta^2 - 137\theta - 241) \\ & + 3^{14} z^6 (9801\theta^4 + 972\theta^3 - 2925\theta^2 + 972\theta + 923) \\ & + 3^{19} z^7 (11\theta^2 + 11\theta + 3)(18\theta^2 + 18\theta + 7) + 3^{24} z^8 (\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (b) * (h) \quad (\text{see [1], Section 7})$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
169	$ \begin{aligned} D = & \theta^4 - 3z(10\theta^2 + 10\theta + 3)(18\theta^2 + 18\theta + 7) \\ & + 3^4 z^2(900\theta^2 + 4592\theta^3 + 6426\theta^2 + 3708\theta + 841) \\ & + 3^9 z^3(180\theta^4 - 1080\theta^3 - 3296\theta^2 - 2532\theta - 723) \\ & + 3^{12} z^4(-1962\theta^4 - 3924\theta^3 + 4770\theta^2 + 6732\theta + 2359) \\ & + 3^{17} z^5(180\theta^4 + 1800\theta^3 + 1024\theta^2 - 100\theta - 227) \\ & + 3^{20} z^6(900\theta^4 - 972\theta^3 - 1890\theta^2 - 972\theta - 113) \\ & - 3^{25} z^7(10\theta^2 + 10\theta + 3)(18\theta^2 + 18\theta + 7) + 3^{32} z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (c) * (h) \quad (\text{see [1], Section 7})$
170	$ \begin{aligned} D = & \theta^4 - 2^2 3 z(3\theta^2 + 3\theta + 1)(18\theta^2 + 18\theta + 7) \\ & + 2^4 3^2 z^2(729\theta^4 + 4860\theta^3 + 6903\theta^2 + 4086\theta + 1007) \\ & + 2^7 3^7 z^3(54\theta^4 - 324\theta^3 - 987\theta^2 - 765\theta - 227) \\ & + 2^{10} 3^8 z^4(-891\theta^4 - 1782\theta^3 + 3222\theta^2 + 4113\theta + 1549) \\ & + 2^{12} 3^{13} z^5(54\theta^4 + 540\theta^3 + 309\theta^2 - 21\theta - 71) \\ & + 2^{14} 3^{14} z^6(729\theta^4 - 1944\theta^3 - 3303\theta^2 - 1944\theta - 307) \\ & - 2^{17} 3^{19} z^7(3\theta^2 + 3\theta + 1)(18\theta^2 + 18\theta + 7) + 2^{20} 3^{24} z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (d) * (h) \quad (\text{see [1], Section 7})$
171	$ \begin{aligned} D = & \theta^4 - 2^2 3 z(72\theta^4 + 288\theta^3 + 254\theta^2 + 110\theta + 21) \\ & + 2^4 3^2 z^2(-1296\theta^4 + 15552\theta^3 + 33696\theta^2 + 23472\theta + 6625) \\ & + 2^9 3^5 z^3(2592\theta^4 + 5184\theta^3 - 9000\theta^2 - 11592\theta - 4499) \\ & + 2^{12} 3^8 z^4(-1296\theta^4 - 20736\theta^3 - 20736\theta^2 - 7920\theta + 1) \\ & + 2^{18} 3^{13} z^5(-72\theta^4 + 178\theta^2 + 178\theta + 51) + 2^{24} 3^{18} z^6(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (e) * (h) \quad (\text{see [1], Section 7})$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
172	$ \begin{aligned} D = & \theta^4 - 3^2 z(3\theta^2 + 3\theta + 1)(18\theta^2 + 18\theta + 7) \\ & + 3^5 z^2(243\theta^4 + 1944\theta^3 + 2763\theta^2 + 1638\theta + 409) \\ & + 3^{11} z^3(54\theta^4 - 324\theta^3 - 987\theta^2 - 765\theta - 227) \\ & + 3^{14} z^4(-648\theta^4 - 1296\theta^3 + 2898\theta^2 + 3456\theta + 1333) \\ & + 3^{20} z^5(54\theta^4 + 540\theta^3 + 309\theta^2 - 21\theta - 7) \\ & + 3^{23} z^6(243\theta^4 - 972\theta^3 - 1611\theta^2 - 972\theta - 167) \\ & - 3^{29} z^7(3\theta^2 + 3\theta + 1)(18\theta^2 + 18\theta + 7) + 3^{36} z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (f) * (h) \quad (\text{see [1], Section 7})$
173	$ \begin{aligned} D = & \theta^4 - z(7\theta^2 + 7\theta + 2)(17\theta^2 + 17\theta + 6) \\ & + 2^6 z^2(55\theta^4 + 112\theta^3 + 155\theta^2 + 86\theta + 15) \\ & + 2^6 3^2 z^3(-119\theta^4 + 714\theta^3 + 2185\theta^2 + 1656\theta + 444) \\ & + 2^{12} 3^2 z^4(92\theta^4 + 184\theta^3 + 98\theta^2 + 6\theta + 9) \\ & + 2^{12} 3^4 z^5(119\theta^4 + 1190\theta^3 + 671\theta^2 - 96\theta - 140) \\ & + 2^{18} 3^4 z^6(55\theta^4 + 108\theta^3 + 149\theta^2 + 108\theta + 27) \\ & + 2^{18} 3^6 z^7(7\theta^2 + 7\theta + 2)(17\theta^2 + 17\theta + 6) + 2^{24} 3^8 z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (a) * (g) \quad (\text{see [1], Section 7})$
174	$ \begin{aligned} D = & \theta^4 - z(11\theta^2 + 11\theta + 3)(17\theta^2 + 17\theta + 6) \\ & + z^2(8711\theta^4 + 33980\theta^3 + 47095\theta^2 + 26230\theta + 5232) \\ & + 2^3 3^2 z^3(-187\theta^4 + 1122\theta^3 + 3436\theta^2 + 2595\theta + 684) \\ & + 2^4 3^2 z^4(8639\theta^4 + 17278\theta^3 - 11650\theta^2 - 20289\theta - 6102) \\ & + 2^6 3^4 z^5(187\theta^4 + 1870\theta^3 + 1052\theta^2 - 163\theta - 216) \\ & + 2^6 3^4 z^6(8711\theta^4 + 864\theta^3 - 2579\theta^2 + 864\theta + 828) \\ & + 2^9 3^6 z^7(11\theta^2 + 11\theta + 3)(17\theta^2 + 17\theta + 6) + 2^{12} 3^8 z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (b) * (g) \quad (\text{see [1], Section 7})$
175	$ \begin{aligned} D = & \theta^4 - z(10\theta^2 + 10\theta + 3)(17\theta^2 + 17\theta + 6) \\ & + 3^4 z^2(89\theta^4 + 452\theta^3 + 633\theta^2 + 362\theta + 80) \\ & + 2^3 3^4 z^3(170\theta^4 - 1020\theta^3 - 3119\theta^2 - 2373\theta - 648) \\ & + 2^4 3^8 z^4(-97\theta^4 - 194\theta^3 + 238\theta^2 + 335\theta + 114) \\ & + 2^6 3^8 z^5(170\theta^4 + 1700\theta^3 + 961\theta^2 - 125\theta - 204) \\ & + 2^6 3^{12} z^6(89\theta^4 - 96\theta^3 - 189\theta^2 - 96\theta - 12) \\ & - 2^9 3^{12} z^7(10\theta^2 + 10\theta + 3)(17\theta^2 + 17\theta + 6) + 2^{12} 3^{16} z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (c) * (g) \quad (\text{see [1], Section 7})$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
176	$ \begin{aligned} D = & \theta^4 - 2^2 z(3\theta^2 + 3\theta + 1)(17\theta^2 + 17\theta + 6) \\ & + 2^5 z^2(325\theta^4 + 2164\theta^3 + 3053\theta^2 + 1778\theta + 420) \\ & + 2^{10} z^3 z^3(51\theta^4 - 306\theta^3 - 934\theta^2 - 717\theta - 204) \\ & + 2^{14} z^2 z^4(-397\theta^4 - 794\theta^3 + 1454\theta^2 + 1851\theta + 666) \\ & + 2^{18} z^4 z^5(51\theta^4 + 510\theta^3 + 290\theta^2 - 29\theta - 64) \\ & + 2^{21} z^4 z^6(325\theta^4 - 864\theta^3 - 1489\theta^2 - 864\theta - 144) \\ & - 2^{26} z^6 z^7(3\theta^2 + 3\theta + 1)(17\theta^2 + 17\theta + 6) + 2^{32} z^8 z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (d) * (g) \quad (\text{see [1], Section 7})$
177	$ \begin{aligned} D = & \theta^4 - 2^2 z(8\theta^2 + 8\theta + 3)(17\theta^2 + 17\theta + 6) \\ & + 2^7 z^2(578\theta^4 + 4040\theta^3 + 5746\theta^2 + 3412\theta + 849) \\ & + 2^{13} z^2 z^3(136\theta^4 - 816\theta^3 - 2485\theta^2 - 1929\theta - 576) \\ & + 2^{19} z^2 z^4(-722\theta^4 - 1444\theta^3 + 2764\theta^2 + 3486\theta + 1323) \\ & + 2^{24} z^4 z^5(136\theta^4 + 1360\theta^3 + 779\theta^2 - 49\theta - 180) \\ & + 2^{29} z^4 z^6(578\theta^4 - 1728\theta^3 - 2906\theta^2 - 1728\theta - 279) \\ & - 2^{35} z^6 z^7(8\theta^2 + 8\theta + 3)(17\theta^2 + 17\theta + 6) + 2^{44} z^8 z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (e) * (g) \quad (\text{see [1], Section 7})$
178	$ \begin{aligned} D = & \theta^4 - 3z(3\theta^2 + 3\theta + 1)(17\theta^2 + 17\theta + 6) \\ & + 3^3 z^2(217\theta^4 + 1732\theta^3 + 2441\theta^2 + 1418\theta + 336) \\ & + 2^3 z^6 z^3(51\theta^4 - 306\theta^3 - 934\theta^2 - 717\theta - 204) \\ & + 2^4 z^8 z^4(-289\theta^4 - 578\theta^3 + 1310\theta^2 + 1599\theta + 570) \\ & + 2^6 z^{11} z^5(51\theta^4 + 510\theta^3 + 290\theta^2 - 29\theta - 64) \\ & + 2^6 z^{13} z^6(217\theta^4 - 864\theta^3 - 1453\theta^2 - 864\theta - 156) \\ & - 2^9 z^{16} z^7(3\theta^2 + 3\theta + 1)(17\theta^2 + 17\theta + 6) + 2^{12} z^{20} z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (f) * (g) \quad (\text{see [1], Section 7})$
179	$ \begin{aligned} D = & \theta^4 - 3z(17\theta^2 + 17\theta + 6)(18\theta^2 + 18\theta + 7) \\ & + 3^4 z^2(2601\theta^4 + 18180\theta^3 + 25929\theta^2 + 15498\theta + 3920) \\ & + 2^3 z^9 z^3(306\theta^4 - 1836\theta^3 - 5587\theta^2 - 4353\theta - 1320) \\ & + 2^4 z^{12} z^4(-3249\theta^4 - 6498\theta^3 + 12366\theta^2 + 15615\theta + 6034) \\ & + 2^6 z^{17} z^5(306\theta^4 + 3060\theta^3 + 1757\theta^2 - 89\theta - 412) \\ & + 2^6 z^{20} z^6(2601\theta^4 - 7776\theta^3 - 13005\theta^2 - 7776\theta - 1228) \\ & - 2^9 z^{25} z^7(17\theta^2 + 17\theta + 6)(18\theta^2 + 18\theta + 7) + 2^{12} z^{32} z^8(\theta + 1)^4 \end{aligned} $
	$\sum_{n=0}^{\infty} A_n z^n = (g) * (h) \quad (\text{see [1], Section 7})$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
180	$D = \theta^4 - 2^4 3z(198\theta^4 + 72\theta^3 + 69\theta^2 + 33\theta + 5)$ $+ 2^9 3^2 z^2(7614\theta^4 + 7128\theta^3 + 6813\theta^2 + 2529\theta + 340)$ $- 2^{14} 3^5 z^3(15714\theta^4 + 27216\theta^3 + 26343\theta^2 + 11151\theta + 1685)$ $+ 2^{19} 3^9 z^4(3\theta + 1)(3\theta + 2)(576\theta^2 + 1008\theta + 605)$ $- 2^{27} 3^{13} z^5(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k} \binom{2n}{2k}^{-1} \frac{(6k)!}{k!(2k)!(3k)!} \frac{(6n-6k)!}{(n-k)!(2n-2k)!(3n-3k)!}$
181	$D = \theta^4 - 18z(324\theta^4 + 648\theta^3 + 765\theta^2 + 441\theta + 97)$ $+ 236196z^2(\theta + 1)^2(6\theta + 5)(6\theta + 7)$
	a formula for A_n is not known
182	$D = \theta^4 - z(43\theta^4 + 86\theta^3 + 77\theta^2 + 34\theta + 6)$ $+ 12z^2(\theta + 1)^2(6\theta + 5)(6\theta + 7)$
	a formula for A_n is not known
183	$D = \theta^4 - 4z(2\theta + 1)^2(7\theta^2 + 7\theta + 3)$ $+ 48z^2(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$
	a formula for A_n is not known
184	$D = \theta^4 - 2z(2\theta + 1)^2(11\theta^2 + 11\theta + 5)$ $+ 500z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	a formula for A_n is not known
185	$D = \theta^4 - 6z(2\theta + 1)^2(3\theta^2 + 3\theta + 1)$ $- 108z^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_{k,l} \binom{n}{k} \binom{n}{l} \binom{k+l}{k}^2 \binom{n}{k+l}$
186	$D = 19^2\theta^4 - 19z(700\theta^4 + 1238\theta^3 + 999\theta^2 + 380\theta + 57)$ $- z^2(64745\theta^4 + 368006\theta^3 + 609133\theta^2 + 412756\theta + 102258)$ $+ 3^3 z^3(6397\theta^4 + 12198\theta^3 - 11923\theta^2 - 27360\theta - 11286)$ $+ 3^6 z^4(64\theta^4 + 1154\theta^3 + 2425\theta^2 + 1848\theta + 486)$ $- 3^{11} z^5(\theta + 1)^4$
	a formula for A_n is not known
187	$D = \theta^4 + z(-64\theta^4 + 898\theta^3 + 653\theta^2 + 204\theta + 27)$ $+ 3^2 z^2(-6397\theta^4 - 13390\theta^3 + 10135\theta^2 + 7492\theta + 1850)$ $+ 3^4 z^3(64745\theta^4 - 109026\theta^3 - 106415\theta^2 - 39528\theta - 4626)$ $+ 3^9 19z^4(700\theta^4 + 1562\theta^3 + 1485\theta^2 + 704\theta + 138)$ $- 3^{14} 19^2 z^5(\theta + 1)^4$
	the reflection of #186 at infinity
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
188	$ \begin{aligned} D = & \theta^4 - 2z(280\theta^4 + 70\theta^3 + 28\theta^2 - 7\theta - 6) \\ & + 2^2 z^2(33544\theta^4 + 16772\theta^3 + 7451\theta^2 + 223\theta + 354) \\ & - 2^5 z^3(562360\theta^4 + 421770\theta^3 + 206211\theta^2 + 30144\theta + 6888) \\ & + 2^6 z^4(23219644\theta^4 + 23219644\theta^3 + 12399919\theta^2 + 2343719\theta + 158340) \\ & - 2^{10} z^5(76599320\theta^4 + 95749150\theta^3 + 55494982\theta^2 + 11674319\theta + 584280) \\ & + 2^{11} z^6(1300767032\theta + 1951150548\theta^3 + 1220689587\theta^2 \\ & \quad + 274877463\theta + 6995856) \\ & - 2^{14} z^7(3491380760\theta^4 + 6109916330\theta^3 + 4107826961\theta^2 \\ & \quad + 935352670\theta - 69427776) \\ & + 2^{15} z^8(22736902622\theta^4 + 45473805244\theta^3 + 32745416803\theta^2 \\ & \quad + 7035711131\theta - 1066887690) \\ & - 2^{17} 15^2 z^9(188347832\theta^4 + 423782622\theta^3 + 326067072\theta^2 \\ & \quad + 67245937\theta - 10221330) \\ & + 2^{18} 15^4 z^{10}(1735944\theta^4 + 4339860\theta^3 + 3554495\theta^2 + 820435\theta + 24846) \\ & - 2^{21} 15^6 z^{11}\theta(\theta + 1)(1036\theta^2 + 2331\theta + 1308) \\ & + 2^{22} 15^8 z^{12}\theta(\theta + 1)(2\theta + 1)(2\theta + 3) \end{aligned} $ the pullback of the 5th-order differential equation $D'y = 0$, where $ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(35\theta^4 + 70\theta^3 + 63\theta^2 + 28\theta + 5) \\ & + 4z^2(\theta + 1)(2\theta + 1)(2\theta + 3)(259\theta^2 + 518\theta + 285) \\ & - 1800z^3(\theta + 1)(\theta + 2)(2\theta + 1)(2\theta + 3)(2\theta + 5) \end{aligned} $ $A'_n = \binom{2n}{n} \sum_{i+j+k+l+m=n} \left(\frac{n!}{i!j!k!l!m!} \right)^2$
189	$ \begin{aligned} D = & \theta^4 - z(1040\theta^4 + 260\theta^3 + 80\theta^2 - 50\theta - 31) \\ & + z^2(409696\theta^4 + 204848\theta^3 + 71714\theta^2 + 15832\theta + 34347) \\ & - 2^2 z^3(18374720\theta^4 + 13781040\theta^3 + 5417700\theta^2 + 2280015\theta - 228513) \\ & + z^4(5406720256\theta^4 + 5406720256\theta^3 + 2370020896\theta^2 \\ & \quad + 470740976\theta - 45885347) \\ & - 2^7 z^5(587991040\theta^4 + 734988800\theta^3 + 370894640\theta^2 \\ & \quad + 26874280\theta - 35246511) \\ & + 2^{12} z^6(104882176\theta^4 + 157323264\theta^3 + 88826496\theta^2 \\ & \quad + 16395264\theta + 8271969) \\ & - 2^{21} z^7(532480\theta^4 + 931840\theta^3 + 577600\theta^2 + 160160\theta + 10273) \\ & + 2^{28} z^8(8\theta + 1)(8\theta + 3)(8\theta + 5)(8\theta + 7) \end{aligned} $ the pullback of the 5th-order differential equation $D'y = 0$, where $ \begin{aligned} D' = & \theta^5 - 2z(2\theta + 1)(65\theta^4 + 130\theta^3 + 105\theta^2 + 40\theta + 6) \\ & + 16z^2(\theta + 1)(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5) \end{aligned} $ $A'_n = \binom{2n}{n} \sum_{j,k,l} \binom{n}{k} \binom{n}{j}^2 \binom{n}{l} \binom{j}{l} \binom{j}{k} \binom{k+l}{k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
190	$ \begin{aligned} D = & \theta^4 - z(4112\theta^4 + 1028\theta^3 + 262\theta^2 - 252\theta - 143) \\ & + z^2(6357088\theta^4 + 3178544\theta^3 + 949302\theta^2 + 146956\theta + 559863) \\ & - 2z^3(2198012032\theta^4 + 1648509024\theta^3 + 567945288\theta^2 \\ & + 254177786\theta - 23879293) \\ & + z^4(1168836133120\theta^4 + 1168836133120\theta^3 + 465472077920\theta^2 \\ & + 105421717520\theta + 12297022465) \\ & - 2^{13}z^5(2198012032\theta^4 + 2747515040\theta^3 + 2068510800\theta^2 \\ & + 1199997450\theta + 333111035) \\ & + 2^{24}z^6(6357088\theta^4 + 9535632\theta^3 + 8338518\theta^2 + 3563032\theta - 95477) \\ & - 2^{32}z^7(65792\theta^4 + 115136\theta^3 + 98608\theta^2 + 28784\theta + 4107) \\ & + 2^{44}z^8(2\theta + 1)^4 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(522\theta^5 + 1285\theta^4 + 1290\theta^3 + 650\theta^2 + 165\theta + 17) \\ & + 5 \cdot 2^2z^2(1032\theta^5 + 3080\theta^4 + 4622\theta^3 + 4618\theta^2 + 2881\theta + 805) \\ & - 5 \cdot 2^3z^3(4112\theta^5 + 14360\theta^4 + 25636\theta^3 + 27154\theta^2 + 14336\theta + 2101) \\ & + 5 \cdot 2^4z^4(8208\theta^5 + 32800\theta^4 + 63528\theta^3 + 65552\theta^2 + 31749\theta + 6554) \\ & - 2^5z^5(40992\theta^5 + 184400\theta^4 + 368720\theta^3 + 368680\theta^2 + 184330\theta + 36865) \\ & - 2^{20}z^6(\theta + 1)^5 \end{aligned} $
	$A'_n = \sum_k \binom{2n-2k}{n-k} \binom{2k}{k}^5$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
191	$ \begin{aligned} D = & \theta^4 - z(27664\theta^4 + 6916\theta^3 + 1422\theta^2 - 2036\theta - 1091) \\ & + z^2(287096928\theta^4 + 143548464\theta^3 + 35543478\theta^2 + 682140\theta + 24820311) \\ & - 2z^3(662746448000\theta^4 + 497059836000\theta^3 + 144112489416\theta^2 \\ & + 58159349850\theta - 9469320873) \\ & + z^4(2303683700982016\theta^4 + 2303683700982016\theta^3 + 768404767144032\theta^2 \\ & + 101811826189072\theta + 5044177631041) \\ & - 2^7 3 z^5(95435488512000\theta^4 + 119294360640000\theta^3 + 85218257140784\theta^2 \\ & + 49484972267112\theta + 14178690000383) \\ & + 2^{12} 3^2 z^6(5953241899008\theta^4 + 8929862848512\theta^3 + 7544293361280\theta^2 \\ & + 3145351380480\theta - 150788399663) \\ & - 2^{21} 3^5 z^7(1147281408\theta^4 + 2007742464\theta^3 + 1668935232\theta^2 \\ & + 456621984\theta + 58834537) \\ & + 2^{28} 3^8 z^8(24\theta + 7)(24\theta + 11)(24\theta + 13)(24\theta + 17) \end{aligned} $ the pullback of the 5th-order differential equation $D'y = 0$, where $ \begin{aligned} D' = & \theta^5 - 2z(3466\theta^5 + 8645\theta^4 + 8338\theta^3 + 3862\theta^2 + 857\theta + 73) \\ & + 2^2 z^2(34600\theta^5 + 103720\theta^4 + 152470\theta^3 + 149954\theta^2 + 93053\theta + 25841) \\ & - 2^3 z^3(138320\theta^5 + 483960\theta^4 + 851700\theta^3 + 889722\theta^2 + 460680\theta + 62757) \\ & + 2^4 z^4(276560\theta^5 + 1106080\theta^4 + 2117960\theta^3 + 2152016\theta^2 + 1013833) \\ & - 2^5 z^5(276512\theta^5 + 1244240\theta^4 + 2463440\theta^3 + 2420968\theta^2 \\ & + 1177258\theta + 226801) \\ & + 3 \cdot 2^{12} z^6(\theta + 1)(4\theta + 3)(4\theta + 5)(6\theta + 5)(6\theta + 7) \end{aligned} $ $A'_n = \sum_k \binom{2n-2k}{n-k} \frac{(3k)!(4k)!}{k!^7}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
192	$ \begin{aligned} D = & \theta^4 - z(442384\theta^4 + 110596\theta^3 + 18054\theta^2 - 37244\theta - 19343) \\ & + 3z^2(24463540256\theta^4 + 12231770128\theta^3 + 2506663954\theta^2 \\ & - 374614268\theta + 2063881213) \\ & - 2z^3(2705798010765440\theta^4 + 2029348508074080\theta^3 + 500517564438600\theta^2 \\ & + 178669097012730\theta - 39466352094909) \\ & + z^4(149673916892499149056\theta^4 + 149673916892499149056\theta^3 \\ & + 43170889056861388896\theta^2 + 2898231542674452496\theta \\ & - 195585894794578943) \\ & - 2^{11}3z^5(389634913550223360\theta^4 + 487043641937779200\theta^3 \\ & + 331475300236995440\theta^2 + 191938784333046600\theta \\ & + 55636145439400469) \\ & + 2^{20}3^2z^6(1521827912245248\theta^4 + 2282741868367872\theta^3 \\ & + 1865287735567488\theta^2 + 756921798415872\theta - 52441938125015) \\ & - 2^{33}3^5z^7(18346549248\theta^4 + 32106461184\theta^3 + 25927048512\theta^2 \\ & + 6633105696\theta + 751215433) \\ & + 2^{44}3^8z^8(24\theta + 5)(24\theta + 11)(24\theta + 13)(24\theta + 19) \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 - 2z(55306\theta^5 + 138245\theta^4 + 128650\theta^3 + 54730\theta^2 + 10469\theta + 721) \\ & + 2^2z^2(553000\theta^5 + 1658920\theta^4 + 2392390\theta^3 + 2315570\theta^2 \\ & + 1429445\theta + 394553) \\ & - 2^3z^3(2211920\theta^5 + 7741560\theta^4 + 13440180\theta^3 + 13847130\theta^2 \\ & + 7036800\theta + 889929) \\ & + 2^4z^4(4423760\theta^5 + 17694880\theta^4 + 33515720\theta^3 + 33546320\theta^2 \\ & + 15386905\theta + 2949122) \\ & - 2^5z^5(4423712\theta^5 + 19906640\theta^4 + 39045200\theta^3 + 37739560\theta^2 \\ & + 17863690\theta + 3317761) \\ & + 2^{18}3z^6(\theta + 1)(3\theta + 2)(3\theta + 4)(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A'_n = \sum_k \binom{2k}{k} \binom{2n-2k}{n-k} \frac{(4k)!}{k!^2(2k)!} \frac{(6k)!}{k!(2k)!(3k)!}$
193	$ \begin{aligned} D = & 7^2\theta^4 - 7z(1135\theta^4 + 2204\theta^3 + 1683\theta^2 + 581\theta + 77) \\ & + z^2(28723\theta^4 + 40708\theta^3 + 13260\theta^2 - 1337\theta - 896) \\ & - z^3(32126\theta^4 + 38514\theta^3 + 26511\theta^2 + 10731\theta + 1806) \\ & + 7 \cdot 11z^4(130\theta^4 + 254\theta^3 + 192\theta^2 + 65\theta + 8) + 11^2z^5(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{n+i+j}{n}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
194	$D = 17^2\theta^4 - 17z(1465\theta^4 + 2768\theta^3 + 2200\theta^2 + 816\theta + 119)$ $+ 2z^2(62015\theta^4 + 131582\theta^3 + 125017\theta^2 + 65926\theta + 15300)$ $- 2 \cdot 3^3z^3(4325\theta^4 + 10914\theta^3 + 12803\theta^2 + 7446\theta + 1700)$ $+ 3^6z^4(265\theta^4 + 836\theta^3 + 1118\theta^2 + 700\theta + 168) - 3^{10}z^5(\theta + 1)^4$
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j}^2$
195	$D = 29^2\theta^4 - 29z(3026\theta^4 + 5848\theta^3 + 4577\theta^2 + 1653\theta + 232)$ $+ z^2(258647\theta^4 + 424220\theta^3 + 239159\theta^2 + 57768\theta + 5568)$ $- z^3(272743\theta^4 + 532614\theta^3 + 581647\theta^2 + 336864\theta + 76560)$ $+ 2^217z^4(1922\theta^4 + 6193\theta^3 + 8121\theta^2 + 4894\theta + 1112)$ $- 2^23 \cdot 17^2z^5(\theta + 1)^2(3\theta + 2)(3\theta + 4)$
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{n+i}{n}$
196	$D = 47^2\theta^4 - 47z(2489\theta^4 + 4984\theta^3 + 4043\theta^2 + 1551\theta + 235)$ $- z^2(208867\theta^4 + 790072\theta^3 + 1135848\theta^2 + 701851\theta + 161022)$ $+ z^3(37085\theta^4 + 637644\theta^3 + 383912\theta^2 + 149319\theta + 38352)$ $+ z^4(291161\theta^4 - 511820\theta^3 - 4424049\theta^2 - 5161283\theta - 1770676)$ $- z^5(406192\theta^4 + 749482\theta^3 + 750755\theta^2 + 260936\theta - 2151)$ $+ 3^3z^6(5305\theta^4 + 90750\theta^3 + 152551\theta^2 + 91194\theta + 17914)$ $+ 2 \cdot 3^6z^7(106\theta^4 + 230\theta^3 + 197\theta^2 + 82\theta + 15) - 2^23^{10}z^8(\theta + 1)^4$
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{n+i-j}{n}$
197	$D = 13^2\theta^4 - 13^2z(41\theta^4 + 82\theta^3 + 67\theta^2 + 26\theta + 4)$ $- 13 \cdot 2^3z^2(471\theta^4 + 1788\theta^3 + 2555\theta^2 + 1534\theta + 338)$ $+ 13 \cdot 2^6z^3(251\theta^4 + 1014\theta^3 + 1798\theta^2 + 1413\theta + 405)$ $+ 2^9z^4(749\theta^4 + 436\theta^3 - 4908\theta^2 - 6266\theta - 2145)$ $- 2^{12}z^5(379\theta^4 + 1270\theta^3 + 967\theta^2 - 42\theta - 178)$ $+ 2^{15}z^6(-9\theta^4 + 156\theta^3 + 273\theta^2 - 156\theta - 28)$ $+ 2^{18}z^7(13\theta^4 + 26\theta^3 + 20\theta^2 + 7\theta + 1) - 2^{21}z^8(\theta + 1)^4$
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{j}{i} \binom{i+j}{j}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
198	$D = 11^2\theta^4 - 7 \cdot 11z(130\theta^4 + 266\theta^3 + 210\theta^2 + 77\theta + 11)$ $- z^2(32126\theta^4 + 89990\theta^3 + 103725\theta^2 + 55253\theta + 11198)$ $+ z^3(28723\theta^4 + 74184\theta^3 + 63474\theta^2 + 20625\theta + 1716)$ $- 7z^4(1135\theta^4 + 2336\theta^3 + 1881\theta^2 + 713\theta + 110) + 7^2z^5(\theta + 1)^4$
	the reflection of #193 at infinity
	$A_n = (-1)^n \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{2n-i}{n}$
199	$D = \theta^4 - z(265\theta^4 + 224\theta^3 + 200\theta^2 + 88\theta + 15)$ $+ 2 \cdot 3z^2(4325\theta^4 + 6386\theta^3 + 6011\theta^2 + 2718\theta + 468)$ $- 2 \cdot 3^2z^3(62015\theta^4 + 116478\theta^3 + 102361\theta^2 + 37422\theta + 4824)$ $+ 17 \cdot 3^6z^4(1465\theta^4 + 3092\theta^3 + 2686\theta^2 + 1140\theta + 200)$ $- 17^23^{10}z^5(\theta + 1)^4$
	the reflection of #194 at infinity
	a formula for A_n is not known
200	$D = 2^2\theta^4 - 2z(106\theta^4 + 194\theta^3 + 143\theta^2 + 46\theta + 6)$ $+ 3z^2(-5305\theta^4 + 6953\theta^3 + 87869\theta^2 + 37122\theta + 6174)$ $+ 3^2z^3(406192\theta^4 + 875286\theta^3 + 939461\theta^2 + 616896\theta + 144378)$ $+ 3^6z^4(-291161\theta^4 - 1676464\theta^3 + 1141623\theta^2 + 986711\theta + 230461)$ $- 3^{10}z^5(370857\theta^4 + 845784\theta^3 + 696122\theta^2 + 189001\theta + 6158)$ $+ 3^{14}z^6(208867\theta^4 + 45396\theta^3 + 18834\theta^2 + 35097\theta + 13814)$ $+ 47 \cdot 3^{18}z^7(2489\theta^4 + 4972\theta^3 + 4025\theta^2 + 1539\theta + 232)$ $- 47^23^{22}z^8(\theta + 1)^4$
	the reflection of #196 at infinity
	a formula for A_n is not known
201	$D = \theta^4 - 2^4z(13\theta^4 + 26\theta^3 + 20\theta^2 + 7\theta + 1)$ $+ 2^8z^2(9\theta^4 + 192\theta^3 + 249\theta^2 + 114\theta + 20)$ $+ 2^{12}z^3(379\theta^4 + 246\theta^3 - 569\theta^2 - 318\theta - 60)$ $+ 2^{16}z^4(-749\theta^4 - 2560\theta^3 + 1722\theta^2 + 1862\theta + 474)$ $- 13 \cdot 2^{20}z^5(251\theta^4 - 10\theta^3 + 262\theta^2 + 145\theta + 27)$ $+ 13 \cdot 2^{24}z^6(471\theta^4 + 96\theta^3 + 17\theta^2 + 96\theta + 42)$ $+ 13^22^{28}z^7(41\theta^4 + 82\theta^3 + 67\theta^2 + 26\theta + 4) - 13^22^{35}z^8(\theta + 1)^4$
	the reflection of #197 at infinity
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
202	$D = 19^2\theta^4 - 19z(1370\theta^4 + 2620\theta^3 + 2089\theta^2 + 778\theta + 114)$ $+ z^2(39521\theta^4 - 3916\theta^3 - 106779\theta^2 - 95266\theta - 25384)$ $+ 2^3z^3(1649\theta^4 + 19779\theta^3 + 29667\theta^2 + 17613\theta + 3876)$ $- 5 \cdot 2^4z^4(\theta + 1)(499\theta^3 + 1411\theta^2 + 1378\theta + 456) + 5^22^9z^5(\theta + 1)^4$
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{n+i-j}{n-j}$
203	$D = 5^2\theta^4 - 5z\theta(499\theta^3 + 86\theta^2 + 53\theta + 10)$ $+ 2^4z^2(1649\theta^4 - 13183\theta^3 - 19776\theta^2 - 11020\theta - 2200)$ $+ 2^6z^3(39521\theta^4 + 162000\theta^3 + 142095\theta^2 + 51540\theta + 6540)$ $- 19 \cdot 2^{11}z^4(1370\theta^4 + 2860\theta^3 + 2449\theta^2 + 1019\theta + 174)$ $+ 19^22^{16}z^5(\theta + 1)^4$
	the reflection of #202 at infinity
	a formula for A_n is not known
204	$D = \theta^4 - 16z(128\theta^4 + 256\theta^3 + 304\theta^2 + 176\theta + 39) + 2^{20}z^2(\theta + 1)^4$
	self-dual at infinity
	a formula for A_n is not known
205	$D = \theta^4 - z(59\theta^4 + 118\theta^3 + 105\theta^2 + 46\theta + 8)$ $+ 96z^2(\theta + 1)^2(3\theta + 2)(3 + 4)$
	a formula for A_n is not known
206	$D = \theta^4 - 4z(2\theta + 1)^2(17\theta^2 + 17\theta + 6)$ $- 224z^2(2\theta + 1)(2\theta + 3)(11\theta^2 + 22\theta + 12)$ $- 1008z^3(2\theta + 1)(2\theta + 3)^2(2\theta + 5)$ $- 24064z^4(2\theta + 1)(2\theta + 3)(2\theta + 5)(2\theta + 7)$
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
207	$D = \theta^4 + 2^4 z(-1072\theta^4 + 17824\theta^3 + 10888\theta^2 + 1976\theta + 145)$ $+ 2^{17} z^2(-51088\theta^4 - 116368\theta^3 + 45264\theta^2 + 14228\theta + 1397)$ $+ 13 \cdot 2^{28} z^3(73104\theta^4 + 1536\theta^3 - 488\theta^2 + 384\theta + 97)$ $- 13^2 2^{44} z^4(2\theta + 1)^4$
	the reflection of #99 at infinity
	a formula for A_n is not known
208	$D = 7^2 \theta^4 - 14z(1056\theta^4 + 1884\theta^3 + 1397\theta^2 + 455\theta + 56)$ $+ 2^2 z^2(68280\theta^4 + 41016\theta^3 - 67611\theta^2 - 54348\theta - 10752)$ $+ 2^4 z^3(-53312\theta^4 + 162120\theta^3 + 195172\theta^2 + 78561\theta + 11130)$ $- 19 \cdot 2^6 z^4(2\theta + 1)^2(1189\theta^2 + 2533\theta + 1646)$ $+ 19^2 2^{11} z^5(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k}^2 \binom{n+2k}{n}$
209	$D = 17^2 \theta^4 - 34z(1902\theta^4 + 3708\theta^3 + 2789\theta^2 + 935\theta + 119)$ $+ 2^2 z^2(62408\theta^4 + 68576\theta^3 - 10029\theta^2 - 24106\theta - 5661)$ $- 2^2 z^3(66180\theta^4 + 33048\theta^3 + 20785\theta^2 + 17799\theta + 4794)$ $+ 2^7 z^4(2\theta + 1)(196\theta^3 + 498\theta^2 + 487\theta + 169)$ $- 2^{12} z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{n+k}{n} \binom{n+2k}{n}$
210	$D = 5^2 \theta^4 - 20z(688\theta^4 + 1352\theta^3 + 981\theta^2 + 305\theta + 35)$ $+ 2^4 z^2(5856\theta^4 + 7008\theta^3 + 96\theta^2 - 1260\theta - 265)$ $+ 2^{10} z^3(176\theta^4 + 120\theta^3 + 69\theta^2 + 30\theta + 5)$ $+ 2^{12} z^4(2\theta + 1)^4$
	$A_n = \binom{2n}{n} \sum_k (-1)^k \binom{2n}{k}^4$
211	$D = \theta^4 + 2^4 z(704\theta^4 + 928\theta^3 + 612\theta^2 + 148\theta + 13)$ $+ 2^{12} z^2(5856\theta^4 + 4704\theta^3 - 1632\theta^2 - 972\theta - 121)$ $+ 2^{20} \cdot 5 z^3(2752\theta^4 + 96\theta^3 - 60\theta^2 + 24\theta + 7)$ $+ 2^{28} \cdot 5^2 z^4(2\theta + 1)^4$
	the reflection of #210 at infinity
	$A_n = (-1)^n \binom{2n}{n}^4 \left(\sum_{k=0}^n (-1)^k \binom{n}{k}^2 \binom{2k}{k} \binom{4n-2k}{2n-k} \binom{n+k}{n}^{-2} \binom{2n}{k}^{-1} \right)$ $+ \sum_{k=1}^n \binom{n}{k}^2 \binom{2n+k}{2n} \binom{4n+2k}{2n+k} \binom{n+k}{n}^{-2} \binom{2k}{k}^{-1}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
212	$ \begin{aligned} D = & 7^2\theta^4 - 14z(134\theta^4 + 286\theta^3 + 234\theta^2 + 91\theta + 14) \\ & - 2^2z^2(3183\theta^4 + 10266\theta^3 + 13501\theta^2 + 8225\theta + 1918) \\ & - 2^3z^3(2588\theta^4 + 8400\theta^3 + 10256\theta^2 + 5649\theta + 1190) \\ & - 2^4 \cdot 3z^4(256\theta^4 + 848\theta^3 + 1141\theta^2 + 717\theta + 174) \\ & - 2^8 \cdot 3^2z^5(\theta + 1)^4 \end{aligned} $
	the reflection of #117 at infinity
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{2n-i-j}{n}$
213	$ \begin{aligned} D = & 17^2\theta^4 - 34z(647\theta^4 + 1240\theta^3 + 977\theta^2 + 357\theta + 51) \\ & - 2^2z^2(14437\theta^4 + 89752\theta^3 + 147734\theta^2 + 92123\theta + 20400) \\ & + 2^2z^3(64614\theta^4 + 77040\theta^3 - 125937\theta^2 - 168453\theta - 52326) \\ & + 2^3z^4(51920\theta^4 + 166384\theta^3 - 83149\theta^2 - 217017\theta - 79362) \\ & + 2^4 \cdot 3z^5(-9360\theta^4 + 26784\theta^3 + 43813\theta^2 + 21965\theta + 3496) \\ & + 2^5 \cdot 3z^6(-10160\theta^4 + 96\theta^3 + 10535\theta^2 + 5385\theta + 438) \\ & - 2^8 \cdot 3^2z^7(288\theta^4 + 864\theta^3 + 1082\theta^2 + 641\theta + 147) \\ & - 2^{11} \cdot 3^2z^8(\theta + 1)^2(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} \binom{2i}{n}$
214	$ \begin{aligned} D = & \theta^4 - 2z(90\theta^4 + 188\theta^3 + 141\theta^2 + 47\theta + 6) \\ & - 2^2z^2(564\theta^4 + 1520\theta^3 + 1705\theta^2 + 934\theta + 192) \\ & - 2^4z^3(2\theta + 1)(286\theta^3 + 813\theta^2 + 851\theta + 294) \\ & - 2^6 \cdot 3z^4(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{n+k}{n} \binom{3k}{n}$
215	$ \begin{aligned} D = & 3^2\theta^4 - 12z(268\theta^4 + 632\theta^3 + 463\theta^2 + 147\theta + 18) \\ & + 2^7z^2(-448\theta^4 + 1616\theta^3 + 4280\theta^2 + 2418\theta + 441) \\ & + 2^{12}z^3(416\theta^4 + 2016\theta^3 + 756\theta^2 - 288\theta - 135) \\ & + 2^{19}z^4(2\theta + 1)^2(8\theta^2 - 28\theta - 33) - 2^{24}z^5(2\theta + 1)^2(2\theta + 3)^2 \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k}^2 \binom{4k}{2n}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
216	$ \begin{aligned} D = & \theta^4 - z\theta(27\theta^3 + 18\theta^2 + 11\theta + 2) \\ & - 2 \cdot 3^3 z^2(72\theta^4 + 414\theta^3 + 603\theta^2 + 330\theta + 64) \\ & + 2^2 \cdot 3^5 z^3(93\theta^4 - 720\theta^3 - 708\theta^2 - 184) \\ & + 2^3 \cdot 3^7 z^4(2\theta + 1)(54\theta^3 + 405\theta^2 + 544\theta + 200) \\ & - 2^4 \cdot 3^{10} z^5(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{3k}{n} \binom{3n-3k}{n}$
217	$ \begin{aligned} D = & 7^2\theta^4 + 7z\theta(13\theta^3 - 118\theta^2 - 73\theta - 14) \\ & - 2^3 \cdot 3z^2(3378\theta^4 + 13446\theta^3 + 18869\theta^2 + 11158\theta + 2352) \\ & - 2^4 \cdot 3^3 z^3(3628\theta^4 + 17920\theta^3 + 31668\theta^2 + 22596\theta + 5383) \\ & - 2^8 \cdot 3^3 z^4(2\theta + 1)(572\theta^3 + 2370\theta^2 + 2896\theta + 1095) \\ & - 2^{10} \cdot 3^4 z^5(2\theta + 1)(2\theta + 3)(6\theta + 5)(6\theta + 7) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{3k}{n} \binom{2n-2k}{n}$
218	$ \begin{aligned} D = & 7^2\theta^4 - 2 \cdot 3 \cdot 7z(192\theta^4 + 396\theta^3 + 303\theta^2 + 105\theta + 14) \\ & + 2^2 \cdot 3z^2(1188\theta^4 + 11736\theta^3 + 20431\theta^2 + 12152\theta + 2436) \\ & + 2^2 \cdot 3^3 z^3(532\theta^4 + 504\theta^3 - 3455\theta^2 - 3829\theta - 1036) \\ & - 2^4 \cdot 3^4 z^4(2\theta + 1)(36\theta^3 + 306\theta^2 + 421\theta + 156) \\ & - 2^6 \cdot 3^4 z^5(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4) \end{aligned} $
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{2k}{n} \binom{3k}{n}$
219	$ \begin{aligned} D = & 5^2\theta^4 - 2 \cdot 5z(464\theta^4 + 1036\theta^3 + 763\theta^2 + 245\theta + 30) \\ & - 2^2 \cdot 3^2 z^2(7064\theta^4 + 22472\theta^3 + 26699\theta^2 + 13200\theta + 2340) \\ & - 2^4 \cdot 3^4 z^3(3440\theta^4 + 13320\theta^3 + 18784\theta^2 + 10665\theta + 2070) \\ & - 2^6 \cdot 3^8 z^4(2\theta + 1)^2(19\theta^2 + 59\theta + 45) - 2^8 \cdot 3^9 z^5(2\theta + 1)^2(2\theta + 3)^2 \end{aligned} $
	$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k}^2 \binom{3k}{n}$
220	$ \begin{aligned} D = & \theta^4 - 2^4 z(20\theta^4 + 56\theta^3 + 38\theta^2 + 10\theta + 1) \\ & - 2^{10} z^2(84\theta^4 + 240\theta^3 + 261\theta^2 + 134\theta + 25) \\ & - 2^{16} z^3(2\theta + 1)^2(23\theta^2 + 55\theta + 39) \\ & - 2^{23} z^4(2\theta + 1)^2(2\theta + 3)^2 \end{aligned} $
	$ \begin{aligned} A_n = & \binom{2n}{n}^3 \left((-1)^n \sum_{k=0}^{[n/2]} \binom{n}{k}^2 \binom{n}{2k} \binom{2n}{4k}^{-1} \right. \\ & \left. + \sum_{k=[n/2]+1}^n \binom{n}{k}^2 \binom{4k}{2n} \binom{2k}{n}^{-1} \right) \end{aligned} $

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
221	$ \begin{aligned} D = & 5^2\theta^4 - 5 \cdot 2^2z(404\theta^4 + 1096\theta^3 + 773\theta^2 + 225\theta + 25) \\ & - 2^4z^2(66896\theta^4 + 137408\theta^3 + 1010960\theta^2 + 52800\theta + 11625) \\ & - 2^8 \cdot 3 \cdot 5z^3(2\theta + 1)(5672\theta^3 + 9500\theta^2 + 8422\theta + 2689) \\ & - 2^{15} \cdot 3^2z^4(2\theta + 1)(1208\theta^3 + 2892\theta^2 + 2842\theta + 969) \\ & - 2^{20} \cdot 3^3z^5(2\theta + 1)(2\theta + 3)(6\theta + 5)(6\theta + 7) \end{aligned} $ $ \begin{aligned} A_n = & \binom{2n}{n}^2 \left((-1)^n \sum_{k=0}^{[n/2]} \binom{n}{k}^2 \binom{n+k}{n} \binom{n}{2k} \binom{2n}{4k}^{-1} \right. \\ & \left. + \sum_{k=[n/2]+1}^n \binom{n}{k}^2 \binom{n+k}{n} \binom{4k}{2n} \binom{2k}{n}^{-1} \right) \end{aligned} $
222	$ \begin{aligned} D = & 5^2\theta^4 - 5z(407\theta^4 + 1198\theta^3 + 909\theta^2 + 310\theta + 40) \\ & - 2^7z^2(2103\theta^4 + 6999\theta^3 + 8358\theta^2 + 4050\theta + 680) \\ & - 2^{12}z^3(1387\theta^4 + 3840\theta^3 + 3081\theta^2 + 960\theta + 100) - 2^{21}z^4(2\theta + 1)^4 \end{aligned} $ $ A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k} \binom{2k}{n} \binom{2n}{n-k} $
223	$ \begin{aligned} D = & \theta^4 + 6z\theta(48\theta^3 - 12\theta^2 - 7\theta - 1) \\ & + 2^23^2z^2(392\theta^4 + 488\theta^3 + 775\theta^2 + 376\theta + 64) \\ & + 2^43^5z^3(1184\theta^4 + 3288\theta^3 + 3512\theta^2 + 1635\theta + 278) \\ & + 2^63^8z^4(2\theta + 1)^2(169\theta^2 + 361\theta + 238) + 2^{11}3^{11}z^5(2\theta + 1)^2(2\theta + 3)^2 \end{aligned} $ $ A_n = \binom{2n}{n} \sum_k (-1)^k 3^{2n-3k} \binom{2n}{3k} \binom{2k}{n} \frac{(3k)!}{k!^3} $
224	$ \begin{aligned} D = & 5^2\theta^4 - 5z(1057\theta^4 + 1058\theta^3 + 819\theta^2 + 290\theta + 40) \\ & + 2^5z^2(10123\theta^4 + 11419\theta^3 + 5838\theta^2 + 1510\theta + 180) \\ & - 2^8z^3(3098\theta^4 + 46560\theta^3 + 48211\theta^2 + 25500\theta + 5100) \\ & + 2^{14} \cdot 11z^4(2\theta + 1)(234\theta^3 + 591\theta^2 + 581\theta + 202) \\ & - 2^{20} \cdot 11^2z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3) \end{aligned} $ $ A_n = 2^{-n} \binom{2n}{n}^2 \sum_k (-1)^{n+k} \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2k}{n} \binom{2n}{2k}^{-1} $
225	$ \begin{aligned} D = & \theta^4 + 2^4z(22192\theta^4 - 17056\theta^3 - 9576\theta^2 - 1048\theta - 49) \\ & + 2^{20}z^2(33648\theta^4 - 44688\theta^3 + 16224\theta^2 + 1764\theta + 17) \\ & + 5 \cdot 2^{34}z^3(6512\theta^4 - 6144\theta^3 - 4440\theta^2 - 1536\theta - 193) \\ & - 5^2 \cdot 2^{55}z^4(2\theta + 1)^4 \end{aligned} $ the reflection of #222 at infinity $ \begin{aligned} A_n = & \binom{2n}{n}^2 \sum_k \binom{n}{k} \binom{n+2k}{k} \binom{3n-2k}{n-k} \binom{2n+4k}{n+2k} \binom{6n-4k}{3n-2k} \\ & \times (1 + k(-2H_k + 2H_{n-k} - H_{n+k} + H_{2n-k} - 2H_{n+2k} \\ & + 2H_{3n-2k} + 4H_{2n+4k} - 4H_{6n-4k})) \end{aligned} $

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
226	$D = 5^2\theta^4 - 2 \cdot 5z(328\theta^4 + 692\theta^3 + 551\theta^2 + 205\theta + 30)$ $+ 2^2 \cdot 3z^2(5352\theta^4 + 25416\theta^3 + 38387\theta^2 + 23020\theta + 4860)$ $- 2^4 \cdot 3^3z^3(352\theta^4 + 4520\theta^3 + 12108\theta^2 + 10205\theta + 2630)$ $- 2^6 \cdot 3^3z^4(2\theta + 1)(586\theta^4 + 3039\theta^2 + 3947\theta + 1527)$ $- 2^8 \cdot 3^4z^5(2\theta + 1)(2\theta + 3)(6\theta + 5)(6\theta + 7)$
	$A_n = \binom{2n}{n} \sum_k (-1)^{n+k} \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3k}{n}$
227	$D = \theta^4 - 2^2 \cdot 3^2z(132\theta^4 + 264\theta^3 + 201\theta^2 + 69\theta + 10)$ $+ 2^9 \cdot 3^6z^2(20\theta^4 + 80\theta^3 + 107\theta^2 + 54\theta + 10)$ $+ 2^{12} \cdot 3^{10}z^3(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = 432^n \binom{2n}{n} \sum_k (-1)^k \binom{n}{k} \binom{3k}{n} \binom{-1/6}{k} \binom{-5/6}{k}$
228	$D = \theta^4 - 2^2z(176\theta^4 + 352\theta^3 + 289\theta^2 + 113\theta + 18)$ $+ 2^{11}z^2(80\theta^4 + 320\theta^3 + 449\theta^2 + 258\theta + 54)$ $- 3 \cdot 2^{16}z^3(2\theta + 1)(2\theta + 5)(4\theta + 3)(4\theta + 9)$
	$A_n = 64^n \binom{2n}{n} \sum_k (-1)^{n+k} \binom{n}{k} \binom{3k}{n} \binom{-1/4}{k} \binom{-3/4}{k}$
229	$D = \theta^4 - 2^2z(256\theta^4 + 728\theta^3 + 506\theta^2 + 142\theta + 15)$ $+ 2^4 \cdot 3^2z^2(-2336\theta^4 - 2336\theta^3 + 1768\theta^2 + 1176\theta + 189)$ $+ 2^9 \cdot 3^4z^3(-512\theta^4 + 432\theta^3 + 404\theta^2 + 108\theta + 9)$ $+ 2^{12} \cdot 3^8z^4(2\theta + 1)^4$
	$A_n = \binom{2n}{n}^2 \sum_{i+j+k=n} \left(\frac{(2n)!}{i!j!k!} \right)^2$
230	$D = \theta^4 + 3z(945\theta^4 - 162\theta^3 - 49\theta^2 + 32\theta + 8)$ $+ 2 \cdot 3^2z^2(17928\theta^4 + 2970\theta^3 + 10187\theta^2 + 3376\theta + 408)$ $+ 2^2 \cdot 3^7z^3(156285\theta^4 + 200016\theta^3 + 19630\theta^2 + 84378\theta + 13964)$ $+ 2^4 \cdot 3^{10} \cdot 19z^4(2\theta + 1)^2(4743\theta^2 + 8199\theta + 4922)$ $+ 2^9 \cdot 3^{15} \cdot 19^2z^5(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = 27^n \binom{2n}{n}^2 \sum_k (-1)^k \binom{n}{k} \binom{2k}{n} \binom{n+k}{n}^{-1} \binom{-1/3}{k} \binom{-2/3}{k}$
231	$D = 9\theta^4 - 2^2z(84\theta^4 + 3048\theta^3 + 2217\theta^2 + 693\theta + 90)$ $+ 2^9z^2(-1168\theta^4 + 968\theta^3 + 9518\theta^2 + 5325\theta + 1005)$ $+ 2^{16}z^3(988\theta^4 + 8208\theta^3 - 743\theta^2 - 4230\theta - 1245)$ $+ 2^{24} \cdot 5z^4(2\theta + 1)^2(9\theta^2 - 279\theta - 277)$ $- 2^{33} \cdot 5^2z^5(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n}^3 \left(\sum_{k=0}^{[n/2]} (-1)^k \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{n}{2k} \binom{2n}{4k}^{-1} \binom{n+k}{n}^{-1} \right)$ $+ \sum_{k=[n/2]+1}^n (-1)^{n+k} \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{4k}{2n} \binom{2k}{n}^{-1} \binom{n+k}{n}^{-1}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
232	$D = 5^2\theta^4 - 5z(2617\theta^4 + 4658\theta^3 + 3379\theta^2 + 1050\theta + 120)$ $- 2^6 \cdot 3z^2(-673\theta^4 + 4871\theta^3 + 10282\theta^2 + 5410\theta + 860)$ $+ 2^{10} \cdot 3^2z^3(955\theta^4 + 4320\theta^3 + 3477\theta^2 + 1020\theta + 100)$ $- 2^{17} \cdot 3^3z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k}^2 \binom{3n}{n+k}$
233	$D = \theta^4 - 2^4z(83\theta^4 + 94\theta^3 + 71\theta^2 + 24\theta + 3)$ $+ 3 \cdot 2^{11}z^2(101\theta^4 + 191\theta^3 + 174\theta^2 + 71\theta + 10)$ $- 3^22^{16}z^3(203\theta^4 + 432\theta^3 + 333\theta^2 + 102\theta + 11)$ $+ 3^32^{23}z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \binom{2n}{n}^3 \sum_k \binom{n}{k}^2 \binom{3n}{n+k} \binom{2n}{2k}^{-1}$
234	$D = 7^2\theta^4 - 7 \cdot 2z\theta(192\theta^3 + 60\theta^2 + 37\theta + 7)$ $- 2^2z^2(17608\theta^4 + 115144\theta^3 + 166715\theta^2 + 94556\theta + 18816)$ $+ 2^4 \cdot 3^2z^3(20288\theta^4 + 57288\theta^3 + 27524\theta^2 - 7455\theta - 5026)$ $- 2^6 \cdot 3^5z^4(2\theta + 1)(458\theta^3 - 657\theta^2 - 1799\theta - 846)$ $- 2^{12} \cdot 3^8z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k}^2 \binom{2k}{n-k} \binom{2n-2k}{k}$
235	$D = 7^2\theta^4 - 2 \cdot 7z\theta(46\theta^3 + 52\theta^2 + 33\theta + 7)$ $- 2^2z^2(7332\theta^4 + 28848\theta^3 + 42633\theta^2 + 26670\theta + 6272)$ $- 2^4z^3(2860\theta^4 + 44760\theta^3 + 120483\theta^2 + 111279\theta + 35098)$ $+ 2^9z^4(2230\theta^4 + 5920\theta^3 - 741\theta^2 - 6509\theta - 3049)$ $+ 2^{14}z^5(174\theta^4 + 1320\theta^3 + 1971\theta^2 + 1095\theta + 190)$ $+ 2^{19}z^6(-22\theta^4 - 24\theta^3 + 9\theta^2 + 21\theta + 7)$ $- 2^{25}z^7(\theta + 1)^4$
	$A_n = \sum_k \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2k}{n-k} \binom{2n-2k}{k}$
236	$D = \theta^4 + 2^4z(22\theta^4 + 64\theta^3 + 51\theta^2 + 19\theta + 3)$ $+ 2^9z^2(-174\theta^4 + 624\theta^3 + 945\theta^2 + 417\theta + 80)$ $+ 2^{14}z^3(-2230\theta^4 - 3000\theta^3 + 5121\theta^2 + 3813\theta + 971)$ $+ 2^{19}z^4(2860\theta^4 - 33320\theta^3 + 3363\theta^2 + 6847\theta + 2402)$ $+ 2^{27}z^5(7332\theta^4 + 480\theta^3 + 81\theta^2 + 1380\theta + 719)$ $+ 2^{36} \cdot 7z^6(\theta + 1)(46\theta^3 + 86\theta^2 + 67\theta + 20)$ $+ 2^{45} \cdot 7^2z^7(\theta + 1)^4$
	the reflection of #235 at infinity
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
237	$D = \theta^4 - 2^4 z(46\theta^4 + 128\theta^3 + 91\theta^2 + 27\theta + 3)$ $+ 2^9 \cdot 3z^2(-74\theta^4 + 16\theta^3 + 231\theta^2 + 127\theta + 20)$ $+ 2^{14} \cdot 3^2 z^3(14\theta^4 + 216\theta^3 + 175\theta^2 + 51\theta + 5)$ $+ 2^{19} \cdot 3^3 z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k}^3 \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k} \binom{2n}{k}^{-1} \binom{2n}{n-k}^{-1}$
238	$D = \theta^4 + 2^2 z(500\theta^4 + 976\theta^3 + 677\theta^2 + 189\theta + 19)$ $+ 2^4 z^2(3968\theta^4 + 3968\theta^3 - 1336\theta^2 - 1164\theta - 177)$ $+ 2^{10} z^3(500\theta^4 + 24\theta^3 - 37\theta^2 + 6\theta + 3) + 2^{12} z^4(2\theta + 1)^4$
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n+2k}{n+k} \binom{2n+k}{2n-k}$
239	$D = \theta^4 - 2^4 \cdot 3z(-9\theta^4 + 198\theta^3 + 131\theta^2 + 32\theta + 39)$ $- 2^{11} \cdot 3^2 z^2(486\theta^4 + 1215\theta^3 + 81\theta^2 - 27\theta - 5)$ $- 2^{16} \cdot 3^5 z^3(891\theta^4 + 972\theta^3 + 675\theta^2 + 216\theta + 25)$ $- 2^{23} \cdot 3^8 z^4(3\theta + 1)^2(3\theta + 2)^2$
	$A_n = \frac{(3n)!}{n!^3} \sum_k \binom{n}{k} \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k}$
240	$D = 13^2 \theta^4 - 13z(1449\theta^4 + 4050\theta^3 + 3143\theta^2 + 1118\theta + 156)$ $+ 2^4 z^2(-22760\theta^4 + 27112\theta^3 + 121046\theta^2 + 82316\theta + 17589)$ $+ 2^8 z^3(3824\theta^4 + 39936\theta^3 - 34292\theta^2 - 63492\theta - 19539)$ $- 2^{16} \cdot 3z^4(2\theta + 1)(40\theta^3 + 684\theta^2 + 1013\theta + 399)$ $- 2^{20} \cdot 3^2 z^5(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$
	$A_n = \sum_k \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{n+2k}{n} \binom{3n-2k}{n}$
241	$D = \theta^4 - 2^4 z(152\theta^4 + 160\theta^3 + 110\theta^2 + 30\theta + 3)$ $+ 2^{10} \cdot 3z^2(428\theta^4 + 176\theta^3 - 299\theta^2 - 170\theta - 25)$ $+ 2^{17} \cdot 3^2 z^3(-136\theta^4 + 216\theta^3 + 180\theta^2 + 51\theta + 5)$ $- 2^{24} \cdot 3^3 z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \binom{2n}{n} \sum_k \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k} \binom{n+k}{n-k} \binom{2n-k}{k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
242	$ \begin{aligned} D = & \theta^4 + 2 \cdot 3z(72\theta^4 + 108\theta^3 + 91\theta^2 + 37\theta + 6) \\ & + 2^2 \cdot 3^3 z^2(648\theta^2 + 1800\theta^3 + 2211\theta^2 + 1248\theta + 260) \\ & + 2^4 \cdot 3^5 z^3(1344\theta^4 + 4968\theta^3 + 7320\theta^2 + 4749\theta + 1072) \\ & + 2^6 \cdot 3^7 z^4(2\theta + 1)(630\theta^3 + 2241\theta^2 + 2617\theta + 971) \\ & + 2^8 \cdot 3^{10} z^5(2\theta + 1)(2\theta + 3)(6\theta + 5)(6\theta + 7) \end{aligned} $
	$ \begin{aligned} A_n = & \binom{2n}{n} \sum_{k=[n/3]}^{[2n/3]} \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3k}{n} \binom{3n-3k}{n} \\ & \times (1 + k(-4H_k + 4H_{n-k} + 2H_{2k} - 2H_{2n-2k} \\ & + 3H_{3k} - 3H_{3n-3k} + 3H_{2n-3k} - 3H_{3k-n})) \\ & + 3 \binom{2n}{n} \sum_{k=0}^{[(n-1)/3]} (-1)^{n+k} \frac{n-2k}{n-3k} \binom{n}{k}^2 \\ & \times \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3n-3k}{n} \binom{n}{3k}^{-1} \end{aligned} $
243	$ \begin{aligned} D = & \theta^4 + z(295\theta^4 + 572\theta^3 + 424\theta^2 + 138\theta + 17) \\ & + 2z^2(843\theta^4 + 744\theta^2 - 473\theta^2 - 481\theta - 101) \\ & + 2z^3(1129\theta^4 - 516\theta^3 - 725\theta^2 - 159\theta + 4) \\ & - 3z^4(173\theta^4 + 352\theta^3 + 290\theta^2 + 114\theta + 18) - 3^2 z^5(\theta + 1)^4 \end{aligned} $
	the reflection of #27 at infinity
	a formula for A_n is not known
244	$ \begin{aligned} D = & \theta^4 + z(416\theta^4 + 104\theta^3 + 36\theta^2 - 16\theta - 11) \\ & + z^2(63168\theta^4 + 31584\theta^3 + 12158\theta^2 + 4828\theta + 6973) \\ & + 2^2 z^3(990080\theta^4 + 742560\theta^3 + 311828\theta^2 + 192658\theta - 33281) \\ & + z^4(62035456\theta^4 + 62035456\theta^3 + 26395808\theta^2 + 5661536\theta + 7849233) \\ & - 2^4 \cdot 3z^5(35642880\theta^4 + 44553600\theta^3 + 24468128\theta^2 \\ & + 1916112\theta + 4807463) \\ & + 2^5 \cdot 3^2 z^6(40932864\theta^4 + 61399296\theta^3 + 38293776\theta^2 \\ & + 9496512\theta + 4807463) \\ & - 2^8 \cdot 3^5 z^7(539136\theta^4 + 943488\theta^3 + 647568\theta^2 + 229320\theta + 23377) \\ & + 2^8 \cdot 3^8 z^8(12\theta + 5)^2(12\theta + 7)^2 \end{aligned} $
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$ \begin{aligned} D' = & \theta^5 + 2z(2\theta + 1)(26\theta^4 + 52\theta^3 + 44\theta^2 + 18\theta + 3) \\ & - 12z^2(\theta + 1)^3(6\theta + 5)(6\theta + 7) \end{aligned} $
	$ \begin{aligned} A'_n = & \sum_k \binom{n}{k}^6 \binom{2k}{k} \binom{2n-2k}{n-k} \\ & \times (1 + k(-8H_k + 8H_{n-k} + 2H_{2k} - 2H_{2n-2k})) \end{aligned} $

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
245	$D = \theta^4 - 3z(144\theta^4 + 72\theta^3 + 32\theta^2 - 4\theta - 5) \\ + 3^2 z^2(7776\theta^4 + 7776\theta^3 + 4086\theta^2 + 828\theta + 589) \\ - 2^3 \cdot 3^7 z^3(288\theta^4 + 432\theta^3 + 262\theta^2 + 81\theta + 6) \\ + 3^{12} z^4(4\theta + 1)^2(4\theta + 3)^2$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 6z(2\theta + 1)(18\theta^4 + 36\theta^3 + 34\theta^2 + 16\theta + 3) \\ + 2916z^2(\theta + 1)^3(2\theta + 1)(2\theta + 3)$
	$A'_n = 3 \binom{2n}{n}^2 \sum_{k=0}^{[n/3]} (-1)^k \frac{n-2k}{2n-3k} \binom{n}{k}^4 \binom{3n-3k}{2n} \binom{2n}{3k}^{-1}$
246	$D = 5^2 \theta^4 - 2^2 \cdot 5z(12\theta^4 + 48\theta^3 + 49\theta^2 + 25\theta + 5) \\ - 2^4 z^2(544\theta^4 + 1792\theta^3 + 2444\theta^2 + 1580\theta + 405) \\ + 2^9 z^3(112\theta^4 + 960\theta^3 + 2306\theta^2 + 2130\theta + 685) \\ + 2^{12} z^4(144\theta^4 + 768\theta^3 + 1308\theta^2 + 924\theta + 235) \\ + 2^{20} z^5(\theta + 1)^4$
	$A_n = \sum_k \binom{n}{k}^3 \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \\ \times (1 + k(-7H_k + 7H_{n-k} + 4H_{2k} - 4H_{2n-2k}))$
247	$D = \theta^4 - 2^4 z(-144\theta^4 + 192\theta^3 + 132\theta^2 + 36\theta + 5) \\ + 2^{13} z^2(112\theta^4 - 512\theta^3 + 98\theta^2 + 50\theta + 13) \\ - 2^{20} z^3(544\theta^4 + 384\theta^3 + 332\theta^2 + 108\theta + 21) \\ + 2^{30} \cdot 5z^4(-12\theta^4 + 23\theta^3 + 23\theta + 7) \\ + 2^{40} \cdot 5^2 z^5(\theta + 1)^4$
	the reflection of #246 at infinity
	$A_n = 4^{-n} \sum_k \binom{n}{k}^{-3} \binom{2k}{k}^5 \binom{2n-2k}{n-k}^5 \\ \times (1 + k(-7H_k + 7H_{n-k} + 10H_{2k} - 10H_{2n-2k}))$
248	$D = 3^2 \theta^4 - 3z(106\theta^4 + 146\theta^3 + 115\theta^2 + 42\theta + 6) \\ - z^2(4511\theta^4 + 24314\theta^3 + 37829\theta^2 + 23598\theta + 5286) \\ + 2^2 z^3(10457\theta^4 + 32184\theta^3 + 24449\theta^2 + 3627\theta - 1317) \\ - 2^2 \cdot 11z^4(1596\theta^4 + 2040\theta^3 - 101\theta^2 - 1085\theta - 386) \\ - 2^4 \cdot 11^2 z^5(\theta + 1)^2(4\theta + 3)(4\theta + 5)$
	$A_n = (-1)^n \sum_k \binom{n}{k}^3 \binom{n+k}{n} \binom{2n-k}{n} \binom{2k}{k} \binom{2n-2k}{n-k} \\ \times (1 + k(-6H_k + 6H_{n-k} + H_{n+k} - H_{2n-k} + 2H_{2k} - 2H_{2n-2k}))$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
249	$ \begin{aligned} D = & 5^2\theta^4 + 2^2 \cdot 5z(148\theta^4 + 392\theta^3 + 341\theta^2 + 145\theta + 25) \\ & + 2^4z^2(4096\theta^4 + 32128\theta^3 + 57016\theta^2 + 37920\theta + 9175) \\ & + 2^8z^3(-6656\theta^4 - 7680\theta^3 + 36960\theta^2 + 49920\theta + 16985) \\ & - 2^{15}z^4(512\theta^4 + 4864\theta^3 + 9136\theta^2 + 6464\theta + 1587) \\ & + 2^{20}z^5(4\theta + 3)^2(4\theta + 5)^2 \end{aligned} $
	$ \begin{aligned} A_n = & (-1)^n \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \\ & \times (1 + k(-6H_k + 6H_{n-k} + H_{n+k} - H_{2n-k} + 4H_{2k} - 4H_{2n-k})) \end{aligned} $
250	$ \begin{aligned} D = & 23^2\theta^4 - 23z(3271\theta^4 + 5078\theta^3 + 3896\theta^2 + 1357\theta + 184) \\ & + z^2(1357863\theta^4 + 999924\theta^3 - 787393\theta^2 - 850862\theta - 205712) \\ & + 2^3z^3(-775799\theta^4 + 272481\theta^3 + 218821\theta^2 - 176709\theta - 100234) \\ & + 2^4 \cdot 61z^4(-1005\theta^4 + 15654\theta^3 + 36317\theta^2 + 27938\theta + 7304) \\ & - 2^9 \cdot 61^2z^5(\theta + 1)^2(4\theta + 3)(4\theta + 5) \end{aligned} $
	$ \begin{aligned} A_n = & (-1)^n \sum_k \binom{n}{k}^3 \binom{2k}{k} \binom{2n-2k}{n-k} \binom{n+2k}{n} \binom{3n-2k}{n} \\ & \times (1 + k(-5H_k + 5H_{n-k} + 2H_{n+2k} - 2H_{3n-2k})) \end{aligned} $
251	$ \begin{aligned} D = & \theta^4 - 3z\theta(27\theta^3 + 18\theta^2 + 11\theta + 2) \\ & - 2 \cdot 3^2z^2(39\theta^4 + 480\theta^3 + 474\theta^2 + 276\theta + 64) \\ & + 2^3 \cdot 3^4z^3(348\theta^4 + 1152\theta^3 + 1759\theta^2 + 1110\theta + 260) \\ & - 2^3 \cdot 3^5z^4(3420\theta^4 + 15912\theta^3 + 28437\theta^2 + 20544\theta + 5296) \\ & + 2^4 \cdot 3^7z^5(1125\theta^4 + 12546\theta^3 + 31089\theta^2 + 26448\theta + 7480) \\ & + 2^5 \cdot 3^9z^6(1395\theta^4 + 3240\theta^3 - 3378\theta^2 - 7146\theta - 2696) \\ & - 2^7 \cdot 3^{11}z^7(351\theta^4 + 2646\theta^3 + 4767\theta^2 + 3309\theta + 800) \\ & - 2^7 \cdot 3^{13}z^8(3\theta + 2)(3\theta + 4)(6\theta + 5)(6\theta + 7) \end{aligned} $
	$ A_n = \sum_k \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3k}{n} \binom{3n-3k}{n} $
252	$ \begin{aligned} D = & 5^2\theta^4 - 5z(-36\theta^4 + 636\theta^3 + 488\theta^2 + 170\theta + 25) \\ & + 2^4z^2(-21301\theta^4 - 27148\theta^3 + 86889\theta^2 + 63110\theta + 14975) \\ & + 2^8z^3(-19535\theta^4 + 294315\theta^3 + 126425\theta^2 - 54390\theta - 35755) \\ & - 2^{10} \cdot 59z^4(-10981\theta^4 + 29878\theta^3 + 89811\theta^2 + 70372\theta + 17759) \\ & - 2^{15} \cdot 3 \cdot 59^2z^5(3\theta + 2)(3\theta + 4)(4\theta + 3)(4\theta + 5) \end{aligned} $
	$ \begin{aligned} A_n = & \sum_k \binom{n}{k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \binom{n+2k}{n} \binom{3n-2k}{n} \\ & \times (1 + k(-5H_k + 5H_{n-k} + 2H_{2k} - 2H_{2n-2k} + 2H_{n+2k} - 2H_{3n-2k})) \end{aligned} $

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
253	$D = \theta^4 - 2^2 z(64\theta^4 + 32\theta^3 + 15\theta^2 - \theta - 2)$ $+ 2^8 z^2(96\theta^4 + 96\theta^3 + 53\theta^2 + 13\theta + 8)$ $- 2^{12} z^3(256\theta^4 + 384\theta^3 + 244\theta^2 + 84\theta + 7)$ $+ 2^{18} z^4(2\theta + 1)^2(4\theta + 1)(4\theta + 3)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 - 4z(2\theta + 1)(16\theta^4 + 32\theta^3 + 31\theta^2 + 15\theta + 3)$ $+ 16z^2(\theta + 1)(4\theta + 3)^2(4\theta + 5)^2$
	$A'_n = \sum_k \binom{n}{k}^2 \binom{n+k}{n} \binom{2n-k}{n} \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2$ $\times (1 + k(-7H_k + 7H_{n-k} + H_{n+k} - H_{2n-k} + 4H_{2k} - 4H_{2n-2k}))$
254	$D = \theta^4 + 2^4 z(-2608\theta^4 + 544\theta^3 + 200\theta^2 - 72\theta - 15)$ $+ 2^{15} \cdot 3z^2(6128\theta^4 - 208\theta^3 + 2328\theta^2 + 452\theta + 25)$ $- 2^{24} \cdot 3^2 \cdot 5z^3(4592\theta^4 + 3456\theta^3 + 2632\theta^2 + 816\theta + 95)$ $+ 2^{38} \cdot 3^3 \cdot 5^2 z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \binom{2n+2k}{n+k}^2 \binom{4n-2k}{2n-k}^2$ $\times (1 + k(-2H_k + 2H_{n-k} - 3H_{n+k} + 3H_{2n-k} + 4H_{2n+2k} - 4H_{4n-2k}))$
255	$D = \theta^4 + 2^2 z(256\theta^4 + 128\theta^3 + 77\theta^2 + 13\theta - 2)$ $+ 2^7 z^2(3072\theta^4 + 3072\theta^3 + 1960\theta^2 + 536\theta + 141)$ $+ 2^{12} z^3(16384\theta^4 + 24576\theta^3 + 16576\theta^2 + 5184\theta + 491)$ $+ 2^{22} z^4(4\theta + 1)(4\theta + 3)(8\theta + 3)(8\theta + 5)$
	the pullback of the 5th-order differential equation $D'y = 0$, where
	$D' = \theta^5 + 4z(2\theta + 1)(64\theta^4 + 128\theta^3 + 141\theta^2 + 77\theta + 17)$ $+ 16z^2(\theta + 1)(8\theta + 5)(8\theta + 7)(8\theta + 9)(8\theta + 11)$
	$A'_n = \sum_k \binom{n}{k}^2 \binom{n+k}{n} \binom{2n-k}{n} \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!}$ $\times (1 + k(-5H_k + 5H_{n-k} + H_{n+k} - H_{2n-k} - 2H_{2k} + 2H_{2n-2k}))$
256	$D = \theta^4 + 2^5 z(24\theta^4 + 42\theta^3 + 30\theta^2 + 9\theta + 1)$ $+ 2^8 z^2(164\theta^4 + 104\theta^3 - 144\theta^2 - 100\theta - 17)$ $+ 2^{14} z^3(28\theta^4 - 48\theta^3 - 44\theta^2 - 12\theta - 1) - 2^{18} z^4(2\theta + 1)^4$
	$A_n = \sum_k \binom{n}{k}^3 \binom{n+k}{n} \binom{2n-k}{n} \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k}$ $\times (1 + k(-4H_k + 4H_{n-k} - H_{n+k} + H_{2n-k} + 2H_{2n+2k} - 2H_{4n-2k}))$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
257	$D = \theta^4 - 2^4 z(112\theta^4 + 416\theta^3 + 280\theta^2 + 72\theta + 7)$ $+ 2^{12} z^2(-656\theta^4 - 896\theta^3 + 216\theta^2 + 160\theta + 23)$ $- 2^{23} z^3(96\theta^4 + 24\theta^3 + 12\theta^2 + 6\theta + 1) - 2^{30} z^4(2\theta + 1)^4$
	the reflection of #256 at infinity
	$A_n = (-1)^n \binom{2n}{n}^2 \sum_k \binom{n}{k} \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k} \binom{n+k}{n-k} \binom{2n-k}{k}$ $\times (1 + k(-2H_k + 2H_{n-k} - H_{n+k} + H_{2n-k} - 2H_{2k}$ $+ 2H_{2n-2k} + 2H_{2n+2k} - 2H_{4n-2k}))$
258	$D = \theta^4 - 2^4 z(16\theta^4 + 224\theta^3 + 156\theta^2 + 44\theta + 5)$ $+ 2^{14} z^2(-48\theta^4 - 48\theta^3 + 120\theta^2 + 66\theta + 11)$ $+ 2^{22} z^3(-16\theta^4 + 192\theta^3 + 156\theta^2 + 48\theta + 5) + 2^{32} z^4(2\theta + 1)^4$
	the case is self-dual at infinity
	$A_n = \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k}$ $\times (1 + k(-4H_k + 4H_{n-k} - H_{n+k} + H_{2n-k} + 2H_{2k}$ $- 2H_{2n-2k} + 2H_{2n+2k} - 2H_{4n-2k}))$
259	$D = \theta^4 + 2 \cdot 3z(72\theta^4 + 108\theta^3 + 91\theta^2 + 37\theta + 6)$ $+ 2^2 \cdot 3^3 z^2(648\theta^4 + 1800\theta^3 + 2211\theta^2 + 1248\theta + 260)$ $+ 2^4 \cdot 3^5 z^3(1344\theta^4 + 4968\theta^3 + 7320\theta^2 + 4749\theta + 1072)$ $+ 2^6 \cdot 3^7 z^4(2\theta + 1)(630\theta^3 + 2241\theta^2 + 2617\theta + 971)$ $+ 2^8 \cdot 3^{10} z^5(2\theta + 1)(2\theta + 3)(6\theta + 5)(6\theta + 7)$
	$A_n = \binom{2n}{n} \sum_{k=[n/3]}^{[2n/3]} \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3k}{n} \binom{3n-3k}{n}$ $\times (1 + k(-4H_k + 4H_{n-k} + 2H_{2k} - 2H_{2n-2k}$ $+ 3H_{3k} - 3H_{3n-3k} + 3H_{2n-3k} - 3H_{3k-n}))$ $+ 3 \binom{2n}{n} \sum_{k=0}^{[(n-1)/3]} (-1)^{n+k} \frac{n-2k}{n-3k} \binom{n}{k}^2$ $\times \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3n-3k}{n} \binom{n}{3k}^{-1}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
260	$D = 5^2\theta^4 + 2^2 \cdot 5z(596\theta^4 + 544\theta^3 + 397\theta^2 + 125\theta + 15)$ $+ 2^4 \cdot 3z^2(30048\theta^4 + 14784\theta^3 - 13312\theta^2 - 10940\theta - 2115)$ $+ 2^8 \cdot 3^3z^3(6368\theta^4 - 6720\theta^3 - 9052\theta^2 - 4080\theta - 655)$ $- 2^{12} \cdot 3^6z^4(2\theta + 1)^2(76\theta^2 + 196\theta + 139)$ $- 2^{16} \cdot 3^9z^5(2\theta + 1)^2(2\theta + 3)^2$
	$A_n = \binom{2n}{n}^2 \sum_{k=[n/3]}^{[2n/3]} \binom{n}{k}^3 \binom{2k}{n-k} \binom{2n-2k}{k}$ $\times (1 + k(-4H_k + 4H_{n-k} + 2H_{2k} - 2H_{2n-2k} + 3H_{2n-3k} - 3H_{3k-n}))$ $+ 3 \binom{2n}{n}^2 \sum_{k=0}^{[(n-1)/3]} (-1)^{n+k} \frac{n-2k}{n-3k} \binom{n}{k}^3 \binom{2n-2k}{k} \binom{n-k}{2k}^{-1}$
261	$D = 5^2\theta^4 + 2^2 \cdot 5z(292\theta^4 + 368\theta^3 + 289\theta^2 + 105\theta + 15)$ $+ 2^4z^2(24736\theta^4 + 43648\theta^3 + 38936\theta^2 + 18980\theta + 3735)$ $+ 2^9 \cdot 3^2z^3(2512\theta^4 + 5760\theta^3 + 6328\theta^2 + 3330\theta + 655)$ $+ 2^{12} \cdot 3^4z^4(2\theta + 1)(232\theta^3 + 588\theta^2 + 590\theta + 207)$ $+ 2^{18} \cdot 3^6z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3)$
	$A_n = \binom{2n}{n} \sum_{k=[n/3]}^{[2n/3]} \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2k}{n-k} \binom{2n-2k}{k}$ $\times (1 + k(-5H_k + 5H_{n-k} + 4H_{2k} - 4H_{2n-2k} + 3H_{2n-3k} - 3H_{3k-n}))$ $+ 3 \binom{2n}{n} \sum_{k=0}^{[(n-1)/3]} (-1)^{n+k} \frac{n-2k}{n-3k} \binom{n}{k}^2$ $\times \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2n-2k}{k} \binom{n-k}{2k}^{-1}$
262	$D = 5^2\theta^4 + 2^2 \cdot 5z(136\theta^4 + 224\theta^3 + 197\theta^2 + 85\theta + 15)$ $+ 2^4z^2(5584\theta^4 + 16192\theta^3 + 21924\theta^2 + 14800\theta + 3955)$ $+ 2^{11}z^3(608\theta^4 + 2280\theta^3 + 3642\theta^2 + 2745\theta + 780)$ $+ 2^{14}z^4(464\theta^4 + 1888\theta^3 + 2956\theta^2 + 2012\theta + 501) + 2^{24}z^5(\theta + 1)^4$
	$A_n = \sum_{k=[8n/3]}^{[2n/3]} \binom{n}{k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \binom{2k}{n-k} \binom{2n-2k}{k}$ $\times (1 + k(-6H_k + 6H_{n-k} + 6H_{2k} - 6H_{2n-2k} + 3H_{2n-3k} - 3H_{3k-n}))$ $+ 3 \sum_{k=0}^{[(n-1)/3]} (-1)^{n+k} \frac{n-2k}{n-3k} \binom{n}{k} \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \binom{2n-2k}{k} \binom{n-k}{2k}^{-1}$
263	$D = \theta^4 + 2^4z(464\theta^4 - 32\theta^3 + 76\theta^2 + 92\theta + 21)$ $+ 2^{15}z^2(608\theta^4 + 152\theta^3 + 450\theta^2 + 131\theta + 5)$ $+ 2^{22}z^3(5584\theta^4 + 6144\theta^3 + 6852\theta^2 + 2808\theta + 471)$ $+ 2^{34} \cdot 5z^4(136\theta^4 + 320\theta^3 + 341\theta^2 + 181\theta + 39) + 2^{46}z^5(\theta + 1)^4$
	the reflection of #262 at infinity
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
264	$D = \theta^4 - 2^4 z(-3392\theta^4 + 9344\theta^3 + 5764\theta^2 + 1092\theta + 93)$ $+ 2^{17} \cdot 3z^2(-1952\theta^4 - 15200\theta^3 + 7758\theta^2 + 2593\theta + 323)$ $+ 2^{26} \cdot 3^2 \cdot 7z^3(-11584\theta^4 + 6912\theta^3 + 5364\theta^2 + 1632\theta + 167)$ $+ 2^{42} \cdot 3^3 \cdot 7^2 z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = 16^{-n} \binom{2n}{n}^2 \left(\sum_{k=0}^n \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2n+2k}{n+k}^2 \binom{4n-2k}{2n-k}^2 \right.$ $\times \binom{2n}{k}^{-1} \binom{2n}{n-k}^{-1} (1 + k(-2H_k + 2H_{n-k} - 3H_{n+k} + 3H_{2n-k}$ $+ 2H_{2k} - 2H_{2n-2k} + 4H_{2n+2k} - 4H_{4n-2k}))$ $+ \sum_{k=1}^n \frac{n+2k}{k} \binom{2n+k}{2n} \binom{2n+2k}{n+k} \binom{2n-2k}{n-k}^2 \binom{4n+2k}{2n+k}^2$ $\times \binom{2k}{k}^{-1} \binom{n+k}{n}^{-1} \binom{2n}{n+k}^{-1} \Big)$
265	$D = \theta^4 - 2^4 \cdot 3z(-96\theta^4 + 96\theta^3 + 6\theta^2 + 12\theta + 1)$ $+ 2^{13} \cdot 3z^2(288\theta^4 - 144\theta^3 + 526\theta^2 + 206\theta + 27)$ $+ 2^{20} \cdot 3^3 z^3(288\theta^4 + 864\theta^3 + 652\theta^2 + 204\theta + 23)$ $+ 2^{30} \cdot 3^5 z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \sum_k \binom{n}{k}^3 \binom{n+k}{n} \binom{2n-k}{n} \binom{2n+2k}{n+k}^2 \binom{4n-2k}{2n-k}^2$ $\times \binom{2n}{k}^{-1} \binom{2n}{n-k}^{-1} \cdot (1 + k(-3H_k + 3H_{n-k} - 2H_{n+k}$ $+ 2H_{2n-k} + 4H_{2n+2k} - 4H_{4n-2k}))$
266	$D = 5^2\theta^4 - 3 \cdot 5z(27\theta^4 + 108\theta^3 + 124\theta^2 + 70\theta + 15)$ $- 3^2 z^2(2754\theta^4 + 9072\theta^2 + 13014\theta^2 + 8910\theta + 2440)$ $+ 3^5 z^3(1134\theta^4 + 9720\theta^3 + 23166\theta^2 + 21330\theta + 6890)$ $+ 3^8 z^4(729\theta^4 + 3888\theta^3 + 6606\theta^2 + 4662\theta + 1184) + 3^{15} z^5(\theta + 1)^4$
	$A_n = \sum_k \binom{n}{k}^3 \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3}$ $\times (1 + k(-6H_k + 6H_{n-k} + 3H_{3k} - 3H_{3n-3k}))$
267	$D = \theta^4 + 3^2 z(729\theta^4 - 972\theta^3 - 684\theta^2 - 198\theta - 31)$ $+ 2 \cdot 3^8 z^2(567\theta^4 - 2592\theta^3 + 405\theta^2 + 189\theta + 70)$ $- 2 \cdot 3^{14} z^3(1377\theta^4 + 972\theta^3 + 1161\theta^2 + 459\theta + 113)$ $+ 3^{22} \cdot 5z^4(-27\theta^4 + 38\theta^2 + 38\theta + 12) + 3^{30} \cdot 5^2 z^5(\theta + 1)^4$
	the reflection of #266 at infinity
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
268	$D = 5^2\theta^4 - 2^2 \cdot 3 \cdot 5z(108\theta^4 + 432\theta^3 + 661\theta^2 + 445\theta + 105)$ $- 2^4 \cdot 3^2 z^2(44064\theta^4 + 14515\theta^3 + 239004\theta^2 + 186300\theta + 58045)$ $+ 2^9 \cdot 3^5 z^3(9072\theta^4 + 7760\theta^3 + 180954\theta^2 + 164970\theta + 53965)$ $+ 2^{12} \cdot 3^8 z^4(11664\theta^4 + 62208\theta^3 + 104940\theta^2 + 73836\theta + 18659)$ $+ 2^{20} \cdot 3^{15} z^5(\theta + 1)^4$
	$A_n = \sum_k \binom{n}{k}^3 \frac{(6k)!}{k!(2k)!(3k)!} \frac{(6n-6k)!}{(n-k)!(2n-2k)!(3n-3k)!}$ $\times (1 + k(-4H_k + 4H_{n-k} + 6H_{6k} - 6H_{6n-6k} - 3H_{3k} + 3H_{3n-3k} - 2H_{2k} + 2H_{2n-2k}))$
269	$D = \theta^4 + 2^4 \cdot 3^2 z(11664\theta^4 - 15552\theta^3 - 11700\theta^2 - 3924\theta - 781)$ $+ 2^{13} \cdot 3^8 z^2(9072\theta^4 - 41472\theta^3 + 2106^2 - 54\theta + 1261)$ $- 2^{20} \cdot 3^{14} z^3(44064\theta^4 + 31104\theta^3 + 67932\theta^2 + 32508\theta + 9661)$ $- 2^{30} \cdot 3^{22} \cdot 5z^4(108\theta^4 + 13\theta^2 + 13\theta - 3)$ $+ 2^{40} \cdot 3^{30} \cdot 5^2 z^5(\theta + 1)^4$
	the reflection of #268 at infinity
	a formula for A_n is not known
270	$D = 5^2\theta^4 - 2^2 \cdot 5z(48\theta^4 + 192\theta^3 + 251\theta^2 + 155\theta + 35)$ $- 2^4 z^2(8704\theta^4 + 28672\theta^3 + 43664\theta^2 + 31760\theta + 9265)$ $+ 2^{11} z^3(1792\theta^4 + 15360\theta^3 + 36248\theta^2 + 33240\theta + 10795)$ $+ 2^{16} z^4(2304\theta^4 + 12288\theta^3 + 20816\theta^2 + 14672\theta + 3719) + 2^{30} z^5(\theta + 1)^4$
	$A_n = \sum_k \binom{n}{k}^3 \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!}$ $\times (1 + k(-5H_k + 5H_{n-k} + 4H_{4k} - 4H_{4n-4k} + 2H_{2n-2k} - 2H_{2k}))$
271	$D = \theta^4 - 2^4 z(-2304\theta^4 + 3072\theta^3 + 2224\theta^2 + 668\theta + 121)$ $+ 2^{17} z^2(1792\theta^4 - 8192\theta^3 + 920\theta^2 + 344\theta + 235)$ $- 2^{28} z^3(8704\theta^4 + 6144\theta^3 + 9872\theta^2 + 4368\theta + 1201)$ $+ 2^{44} \cdot 5z^4(-48\theta^4 + 37\theta^2 + 37\theta + 13) + 2^{60} \cdot 5^2 z^5(\theta + 1)^4$
	the reflection of #270 at infinity
	a formula for A_n is not known
272	$D = 5^2\theta^4 - 2^2 \cdot 3 \cdot 5z(1332\theta^4 + 3528\theta^3 + 3289\theta^2 + 1525\theta + 285)$ $+ 2^4 \cdot 3^2 z^2(331776\theta^4 + 1602368\theta^3 + 453333\theta^2 + 2996640\theta + 724415)$ $+ 2^8 \cdot 3^5 z^3(539136\theta^4 + 622080\theta^3 - 3024864\theta^2 - 4008960\theta - 1315985)$ $- 2^{15} \cdot 3^8 z^4(41472\theta^4 + 393984\theta^3 + 735984\theta^2 + 510912\theta + 120811)$ $- 2^{20} \cdot 3^{11} z^5(12\theta + 7)(12\theta + 11)(12\theta + 13)(12\theta + 17)$
	$A_n = \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \frac{(6k)!}{k!(2k)!(3k)!}$ $\times \frac{(6n-6k)!}{(n-k)!(2n-2k)!(3n-3k)!} \cdot (1 + k(-3H_k + 3H_{n-k} + H_{n+k} - H_{2n-k} + 6H_{6k} - 6H_{6n-6k} - 3H_{3k} + 3H_{3n-3k} - 2H_{2k} + 2H_{2n-2k}))$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
273	$D = 5^2\theta^4 - 3 \cdot 5z(333\theta^4 + 882\theta^3 + 781\theta^2 + 340\theta + 60)$ $+ 2^2 \cdot 3^2z^2(5184\theta^4 + 40662\theta^3 + 71829\theta^2 + 47700\theta + 11540)$ $+ 2^2 \cdot 3^5z^3(8424\theta^4 + 9720\theta^3 - 46899\theta^2 - 63045\theta - 21260)$ $- 2^4 \cdot 3^8z^4(1296\theta^4 + 12312\theta^3 + 23094\theta^2 + 16263\theta + 3956)$ $- 2^6 \cdot 3^{11}z^5(3\theta + 2)(3\theta + 4)(6\theta + 5)(6\theta + 7)$ $A_n = \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3}$ $\times (1 + k(-5H_k + 5H_{n-k} + H_{n+k} - H_{2n-k} + 3H_{3k} - 3H_{3n-3k}))$
274	$D = 5^2\theta^4 - 5z(757\theta^4 + 1298\theta^3 + 1049\theta^2 + 400\theta + 60)$ $+ 2^2 \cdot 3^2z^2(5456\theta^4 + 17498\theta^3 + 22121\theta^2 + 11940\theta + 2340)$ $- 2^2 \cdot 3^4z^3(15128\theta^4 + 68040\theta^3 + 112171\theta^2 + 73845\theta + 16380)$ $+ 2^4 \cdot 3^8z^4(2\theta + 1)(216\theta^3 + 864\theta^2 + 1015\theta + 356)$ $- 2^6 \cdot 3^{10}z^5(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4)$ $A_n = 9^{-n} \binom{2n}{n} \sum_{k=[n/3]}^{[2n/3]} \binom{3k}{n} \binom{3n-3k}{n} \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3}$ $\times (1 + k(-3H_k + 3H_{n-k} + 6H_{3k} - 6H_{3n-3k} + 3H_{2n-3k} - 3H_{3k-n}))$ $+ 3 \cdot 9^{-n} \binom{2n}{n} \sum_{k=0}^{[(n-1)/3]} (-1)^k \frac{n-2k}{n-3k} \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3} \binom{3n-3k}{n} \binom{n}{3k}^{-1}$
275	$D = 5^2\theta^4 - 2^2 \cdot 5z(592\theta^4 + 1568\theta^3 + 1419\theta^2 + 635\theta + 115)$ $+ 2^4z^2(6553\theta^4 + 514048\theta^3 + 902816\theta^2 + 598400\theta + 144735)$ $+ 2^{10}z^3(106496\theta^4 + 122880\theta^3 - 594816\theta^2 - 794880\theta - 265065)$ $- 2^{19}z^4(8192\theta^4 + 77824\theta^3 + 145728\theta^2 + 102016\theta + 24527)$ $- 2^{26}z^5(8\theta + 5)(8\theta + 7)(8\theta + 9)(8\theta + 11)$ $A_n = \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \frac{(4k)!}{k!^2(2k)!} \frac{(4n-4k)!}{(n-k)!^2(2n-2k)!}$ $\times (1 + k(-4H_k + 4H_{n-k} + H_{n+k} - H_{2n-k} + 4H_{4k} - 4H_{4n-4k} + 2H_{2n-2k} - 2H_{2k}))$
276	$D = \theta^4 + 2^4 \cdot 3z(-18432\theta^4 + 4608\theta^3 + 1024\theta^2 - 1280\theta - 221)$ $+ 2^{17} \cdot 3^4z^2(25344\theta^4 - 2304\theta^3 + 11680\theta^2 + 1472\theta - 33)$ $- 2^{28} \cdot 3^8z^3(18432\theta^4 + 13824\theta^3 + 11392\theta^2 + 3264\theta + 359)$ $+ 2^{46} \cdot 3^{12}z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$ $A_n = \binom{2n}{n} \sum_k \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k} \frac{(6k)!}{k!(2k)!(3k)!}$ $\times \frac{(6n-6k)!}{(n-k)!(2n-2k)!(3n-3k)!} \cdot (1 + k(-H_k + H_{n-k} - 2H_{2k} + 2H_{2n-2k} - 3H_{3k} + 3H_{3n-3k} + 6H_{6k} - 6H_{6n-6k} + 2H_{2n+2k} - 2H_{4n-2k}))$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
277	$D = \theta^4 - 2^4 z(-576\theta^4 + 1152\theta^3 + 724\theta^2 + 148\theta + 13)$ $+ 2^{17} z^2(-32\theta^4 - 992\theta^2 + 166\theta^2 + 57\theta + 6)$ $- 2^{26} \cdot 3z^3(832\theta^4 + 768\theta^3 + 556\theta^2 + 192\theta + 25) - 2^{40} \cdot 3^2 z^4(2\theta + 1)^4$
	the reflection of #55 at infinity
	$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k} \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k}$ $\times (1 + k(-3H_k + 3H_{n-k} + 2H_{2k} - 2H_{2n-2k} + 2H_{2n-k} + 2H_{n+k} + 2H_{2n+2k} - 2H_{4n-2k}))$
278	$D = \theta^4 - 3z(279\theta^4 + 882\theta^3 + 641\theta^2 + 200\theta + 24)$ $+ 2 \cdot 3^5 z^2(-72\theta^4 + 1710\theta^3 + 3665\theta^2 + 1864\theta + 296)$ $+ 2^2 \cdot 3^9 z^3(909\theta^4 + 3888\theta^3 + 3082\theta^2 + 918\theta + 92)$ $+ 2^4 \cdot 3^{15} z^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2)$
	$A_n = \binom{2n}{n}^2 \binom{3n}{n} \sum_k \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3} \binom{n+k}{n}^{-1} \binom{2n-k}{n}^{-1}$
279	$D = 17^2 \theta^4 + 17z(286\theta^4 + 734\theta^3 + 656\theta^2 + 289\theta + 51)$ $+ 3^2 z^2(4110\theta^4 + 22074\theta^3 + 37209\theta^2 + 26265\theta + 6800)$ $- 3^5 z^3(1521\theta^4 + 7344\theta^3 + 12936\theta^2 + 9945\theta + 2822)$ $+ 3^8 z^4(123\theta^4 + 552\theta^3 + 879\theta^2 + 603\theta + 152) - 3^{12} z^5(\theta + 1)^4$
	$A_n = 3 \sum_{k=[2(n+1)/3]}^n (-1)^k \binom{n}{k} \frac{n-2k}{n-3k} \frac{(3k)!}{k!^3} \frac{(3n-3k)!}{(n-k)!^3} \binom{n}{3n-3k} \binom{3k}{n}^{-1}$
280	$D = \theta^4 + 3^2 z(123\theta^4 - 60\theta^3 - 39\theta^2 - 9\theta - 1)$ $+ 3^5 z^2(1521\theta^2 - 1260\theta^3 + 30\theta^2 - 21\theta - 10)$ $+ 3^8 z^3(4110\theta^4 - 5634\theta^3 - 4353\theta^2 - 1629\theta - 220)$ $- 3^{12} \cdot 17z^4(286\theta^4 + 410\theta^3 + 170\theta^2 - 35\theta - 30) + 3^{18} \cdot 17^2 z^5(\theta + 1)^4$
	the reflection of #279 at infinity
	a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
281	$ \begin{aligned} D = & \theta^4 + z(328\theta^4 + 82\theta^3 + 33\theta^2 - 8\theta - 7) \\ & + z^2(52844\theta^4 + 26422\theta^3 + 11771\theta^2 - 3472\theta - 3676) \\ & + z^3(5280472\theta^4 + 3960354\theta^3 + 1932893\theta^2 - 37864\theta + 197467) \\ & + z^4(355955926\theta^4 + 355955926\theta^3 + 188833541\theta^2 \\ & + 39149936\theta + 53236566) \\ & + 5z^5(3300295000\theta^4 + 4125368750\theta^3 + 2363161375\theta^2 \\ & + 789584250\theta + 300065632) \\ & + 5^6 z^6(33027500\theta^4 + 49541250\theta^3 + 30476125\theta^2 + 10997750\theta + 305303) \\ & + 5^{11} z^7(205000\theta^4 + 358750\theta^3 + 235875\theta^2 + 71750\theta + 6104) \\ & + 5^{16} z^8(5\theta + 1)(5\theta + 2)(5\theta + 3)(5\theta + 4) \end{aligned} $ the pullback of the 5th-order differential equation $D'y = 0$, where $ \begin{aligned} D' = & \theta^5 + z(2\theta + 1)(41\theta^4 + 82\theta^3 + 74\theta^2 + 33\theta + 6) \\ & + 5z^2(\theta + 1)(5\theta + 3)(5\theta + 4)(5\theta + 6)(5\theta + 7) \end{aligned} $ $ \begin{aligned} A'_n = & 3 \sum_{k=[2(n+1)/3]}^n (-1)^k \frac{n-2k}{n-3k} \binom{n+k}{n} \binom{2n-k}{n} \\ & \times \frac{(3k)! (3n-3k)!}{k!^3 (n-k)!^3} \binom{n}{3n-3k} \binom{3k}{n}^{-1} \end{aligned} $
282	$ \begin{aligned} D = & 5^2\theta^4 - 2^2 \cdot 5z(1348\theta^4 + 752\theta^3 + 521\theta^2 + 145\theta + 15) \\ & + 2^4 \cdot 3^4 z^2(5696\theta^4 - 1792\theta^3 - 7304\theta^2 - 3740\theta - 585) \\ & + 2^{10} \cdot 3^8 z^3(-20\theta^4 + 360\theta^3 + 289\theta^2 + 90\theta + 10) \\ & - 2^{12} \cdot 3^{13} z^4(2\theta + 1)^4 \end{aligned} $ $ \begin{aligned} A_n = & (-1)^n \binom{2n}{n} \sum_{k=[n/3]}^{[2n/3]} \binom{2n}{3k} \binom{2n}{3n-3k} \frac{(3k)! (3n-3k)!}{k!^3 (n-k)!^3} \\ & \times (1 + k(-3H_k + 3H_{n-k} + 3H_{2n-3k} - 3H_{3k-n})) \\ & + 3 \binom{2n}{n} \sum_{k=0}^{[(n-1)/3]} (-1)^{n+k} \frac{n-2k}{2n-3k} \frac{(3k)! (3n-3k)!}{k!^3 (n-k)!^3} \binom{2n}{3n-3k} \binom{3k}{2n}^{-1} \end{aligned} $
283	$ \begin{aligned} D = & \theta^4 + 2^2 z(20\theta^4 + 400\theta^3 + 281\theta^2 + 81\theta + 9) \\ & - 2^4 \cdot 3z^2(5696\theta^4 + 13184\theta^3 + 3928\theta^2 + 628\theta + 39) \\ & + 2^{10} \cdot 3^2 \cdot 5z^3(1348\theta^4 + 1944\theta^3 + 1415\theta^2 + 486\theta + 63) \\ & - 2^{12} \cdot 3^7 \cdot 5^2 z^4(2\theta + 1)^4 \end{aligned} $ the reflection of #282 at infinity a formula for A_n is not known

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
284	$ \begin{aligned} D = & 38^2\theta^4 - 38z(3014\theta^4 + 5878\theta^3 + 4725\theta^2 + 1786\theta + 266) \\ & + z^2(689717\theta^4 + 2305502\theta^3 + 3057079\theta^2 + 1810054\theta + 402002) \\ & - z^3(1438808\theta^4 + 5812350\theta^3 + 9142457\theta^2 + 6295992\theta + 1576582) \\ & + z^4(1395491\theta^4 + 5075392\theta^3 + 6297445\theta^2 + 3375833\theta + 663471) \\ & - z^5(834163\theta^4 + 2657224\theta^3 + 2407768\theta^2 + 604005\theta - 52928) \\ & + z^6(277543\theta^4 + 692484\theta^3 + 572576\theta^2 + 148359\theta - 4832) \\ & + 11z^7(4625\theta^4 + 9100\theta^3 + 6395\theta^2 + 1845\theta + 178) - 11^2z^8(\theta + 1)^4 \end{aligned} $ $A_n = \sum_{k,l} \binom{n}{k} \binom{n}{l} \binom{n+k}{n} \binom{n+l}{n} \binom{n}{k+l} \binom{n+k-l}{n-l}$
285	$ \begin{aligned} D = & 11^2\theta^4 + 11z(4625\theta^4 + 9400\theta^3 + 6845\theta^2 + 2145\theta + 253) \\ & - z^2(834163\theta^4 + 417688\theta^3 + 160382\theta^2 + 29513\theta + 4444) \\ & + z^3(834163\theta^4 + 679428\theta^3 - 558926\theta^2 - 423489\theta - 72226) \\ & - z^4(1395491\theta^4 + 506572\theta^3 - 555785\theta^2 - 425155\theta - 94818) \\ & + z^5(1438808\theta^4 + 57118\theta^3 + 338255\theta^2 + 307104\theta + 49505) \\ & - z^6(689717\theta^4 + 453366\theta^3 + 278875\theta^2 + 146466\theta + 33242) \\ & + 38z^7(3014\theta^4 + 6178\theta^3 + 5175\theta^2 + 2086\theta + 341) - 38^2z^8(\theta + 1)^4 \end{aligned} $ the reflection of #284 at infinity a formula for A_n is not known
286	$ \begin{aligned} D = & 3^2\theta^4 - 3^2z(38\theta^4 + 82\theta^3 + 67\theta^2 + 26\theta + 4) \\ & + 3z^2(2045\theta^4 + 5702\theta^3 + 7535\theta^2 + 4170\theta + 852) \\ & + 2^3 \cdot 3z^3(2208\theta^4 + 5925\theta^3 + 7925\theta^2 + 5607\theta + 1512) \\ & + 2^3z^4(60287\theta^4 + 56374\theta^3 - 215983\theta^2 - 268986\theta - 85452) \\ & - 2^4z^5(205651\theta^4 + 605608\theta^3 + 603579\theta^2 + 204622\theta + 8104) \\ & + 2^7z^6(-51414\theta^4 + 273267\theta^3 + 502700\theta^2 + 305649\theta + 63398) \\ & + 2^8 \cdot 37z^7(7909\theta^4 + 18122\theta^3 + 17595\theta^2 + 8462\theta + 1672) \\ & - 2^{13} \cdot 37^2z^8(\theta + 1)^2(4\theta + 3)(4\theta + 5) \end{aligned} $ $A_n = \sum_{k,l} \binom{n}{k} \binom{n}{l} \binom{2n-k}{n} \binom{n}{k-l} \binom{2k}{n} \binom{2l}{n}$
287	$ \begin{aligned} D = & 21^2\theta^4 - 21z(3289\theta^4 + 6098\theta^3 + 4645\theta^2 + 1596\theta + 210) \\ & + 2^2z^2(38560\theta^4 - 230840\theta^3 - 534425\theta^2 - 337050\theta - 68145) \\ & + 2^4z^3(106636\theta^4 + 493416\theta^3 + 420211\theta^2 + 116361\theta + 6090) \\ & - 5 \cdot 2^8z^4(2\theta + 1)(1916\theta^3 + 2622\theta^2 + 1077\theta + 91) \\ & - 5^2 \cdot 2^{12}z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3) \end{aligned} $ $A_n = \binom{2n}{n} \sum_{i,j} \binom{n}{i} \binom{n}{j} \binom{n+j}{n} \binom{i+j}{n} \binom{n}{i-j}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
288	$D = \theta^4 - 2^4 z(496\theta^4 + 1568\theta^3 + 1060\theta^2 + 276\theta + 27)$ $+ 3 \cdot 2^{15} z^2(-32\theta^4 + 760\theta^3 + 1570\theta^2 + 651\theta + 81)$ $+ 3^2 2^{22} z^3(1616\theta^4 + 6912\theta^3 + 5092\theta^2 + 1416\theta + 135)$ $+ 3^3 2^{34} z^4(3\theta + 1)(3\theta + 2)(4\theta + 1)(4\theta + 3)$
	$A_n = \binom{2n}{n} \binom{3n}{n} \binom{4n}{n} \sum_k \frac{(4k)!}{k!^2 (2k)!} \frac{(4n-4k)!}{(n-k)!^2 (2n-2k)!}$ $\times \binom{n+k}{n}^{-1} \binom{2n-k}{n}^{-1}$
289	$D = \theta^4 - 2^4 z(400\theta^4 + 2720\theta^3 + 1752\theta^2 + 392\theta + 33)$ $+ 2^{15} z^2(-4272\theta^4 - 6288\theta^3 + 3184\theta^2 + 1484\theta + 177)$ $+ 2^{24} \cdot 5 z^3(-4688\theta^4 + 1536\theta^3 + 1384\theta^2 + 336\theta + 27)$ $+ 2^{36} \cdot 5^2 z^4(2\theta + 1)^2(4\theta + 1)(4\theta + 3)$
	$A_n = (-1)^n \binom{2n}{n} \binom{4n}{2n} \sum_k \binom{n}{k} \binom{n+k}{n} \binom{2n-k}{n} \binom{2n}{k} \binom{2n}{n-k}$ $\times \binom{2n+2k}{n+k} \binom{4n-2k}{2n-k} \binom{2k}{k}^{-1} \binom{2n-2k}{n-k}^{-1} (1 + k(-H_k + H_{n-k}$ $- 2H_{n+k} + 2H_{2n-k} - 2H_{2k} + 2H_{2n-2k} + 2H_{2n+2k} - 2H_{4n-2k}))$
290	$D = \theta^4 + 3z(-279\theta^4 + 252\theta^3 + 160\theta^2 + 34\theta + 3)$ $+ 2 \cdot 3^5 z^2(423\theta^4 - 468\theta^3 + 457\theta^2 + 215\theta + 37)$ $- 2 \cdot 3^9 z^3(531\theta^4 + 1296\theta^3 + 1243\theta^2 + 567\theta + 104)$ $+ 3^{15} \cdot 5 z^4(51\theta^4 + 120\theta^3 + 126\theta^2 + 66\theta + 14)$ $- 3^{20} \cdot 5^2 z^5(\theta + 1)^4$
	the reflection of #17 at infinity
	a formula for A_n is not known
291	$D = \theta^4 - z(566\theta^4 + 34\theta^3 + 62\theta^2 + 45\theta + 9)$ $+ 3z^2(39370\theta^4 + 17302\theta^3 + 22493\theta^2 + 8369\theta + 1140)$ $- 3^2 z^3(1215215\theta^4 + 1432728\theta^3 + 1274122\theta^2 + 538245\theta + 93222)$ $+ 3^7 \cdot 61 z^4(3029\theta^4 + 6544\theta^3 + 6135\theta^2 + 2863\theta + 548)$ $- 3^{12} \cdot 61^2 z^5(\theta + 1)^4$
	the reflection of #124 at infinity
	a formula for A_n is not known
292	$D = 3^2 \theta^4 - 2^2 \cdot 3z(4636\theta^2 + 7928\theta^3 + 5347\theta^2 + 1383\theta + 126)$ $+ 2^9 z^2(59048\theta^4 + 50888\theta^3 - 26248\theta^2 - 16827\theta - 2205)$ $+ 2^{16} \cdot 7 z^3(-9004\theta^4 + 2304\theta^3 + 2511\theta^2 + 504\theta + 27)$ $- 2^{24} \cdot 7^2 z^4(2\theta + 1)^2(4\theta + 1)(4\theta + 3)$
	$A_n = \binom{2n}{n} \sum_k (-1)^{n+k} 4^{n-k} \binom{n}{k} \binom{2n+k}{n}^2 \binom{4n+2k}{2n+k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
293	$ \begin{aligned} D = & \theta^4 - 2^2 z(54\theta^4 + 66\theta^3 + 49\theta^2 + 16\theta + 2) \\ & + 2^4 z^2(417\theta^4 - 306\theta^3 - 1219\theta^2 - 776\theta - 154) \\ & + 2^8 z^3(166\theta^4 + 1920\theta^3 + 1589\theta^2 + 432\theta + 23) \\ & - 2^{12} \cdot 7 z^4(2\theta + 1)(38\theta^3 + 45\theta^2 + 12\theta - 2) \\ & - 2^{14} \cdot 7^2 z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3) \end{aligned} $
	$A_n = \sum_{k,l} \binom{n}{k} \binom{n}{l} \binom{k+l}{n} \binom{2k}{n} \binom{2l}{n} \binom{2n}{k+l}$
294	$ \begin{aligned} D = & \theta^4 + 2^4 z(-18800\theta^4 + 14624\theta^3 + 8184\theta^2 + 872\theta + 33) \\ & + 2^{18} z^2(101744\theta^4 - 107920\theta^3 + 74968\theta^2 + 15100\theta + 1191) \\ & - 2^{30} \cdot 17 z^3(40048\theta^4 + 49152\theta^3 + 35848\theta^2 + 10752\theta + 1143) \\ & + 2^{50} \cdot 17^2 z^4(2\theta + 1)^2(4\theta + 1)(4\theta + 3) \end{aligned} $
	$ \begin{aligned} A_n = & \frac{(4n)!}{n!^2(2n)!} \sum_k \binom{2n}{2k} \binom{n+2k}{k} \binom{3n-2k}{n-k} \binom{2n+4k}{n+2k} \binom{6n-4k}{3n-3k} \\ & \times (1 + k(-H_k + H_{n-k} - H_{n+k} + H_{2n-k} - 2H_{2k} + 2H_{2n-2k} \\ & + 2H_{3n-2k} - 2H_{n+2k} + 4H_{2n+4k} - 4H_{6n-4k})) \end{aligned} $
295	$ \begin{aligned} D = & \theta^4 + 2^4 z(816\theta^4 - 1440\theta^3 - 904\theta^2 - 184\theta - 17) \\ & + 2^{18} z^2(80\theta^4 - 592\theta^3 + 432\theta^2 + 164\theta + 23) \\ & + 2^{30} z^3(-80\theta^4 + 384\theta^3 + 296\theta^2 + 96\theta + 11) \\ & + 2^{45} z^4(2\theta + 1)^4 \end{aligned} $
	$ \begin{aligned} A_n = & \binom{3n}{n} \sum_k \binom{2k}{k} \binom{2n-2k}{n-k} \binom{2n}{2k} \binom{n+2k}{k} \\ & \times \binom{3n-2k}{n-k} \binom{2n+4k}{n+2k} \binom{6n-4k}{3n-2k} \binom{3n}{n+k}^{-1} \\ & \times (1 + k(-3H_k + 3H_{n-k} - 2H_{n+2k} + 2H_{3n-2k} + 4H_{2n+4k} - 4H_{6n-4k})) \end{aligned} $
296	$ \begin{aligned} D = & \theta^4 - 2^4 z(5\theta^4 + 34\theta^3 + 25\theta^2 + 8\theta + 1) \\ & + 2^{11} z^2(5\theta^4 + 47\theta^3 + 90\theta^2 + 47\theta + 8) \\ & 2^{16} z^3(51\theta^4 + 192\theta^3 + 155\theta^2 + 48\theta + 5) \\ & + 2^{23} z^4(2\theta + 1)^4 \end{aligned} $
	the reflection of #295 at infinity
	a formula for A_n is not known
297	$ \begin{aligned} D = & 7^2 \theta^4 - 2 \cdot 7 z \theta(520\theta^3 + 68\theta^2 + 41\theta + 7) \\ & - 2^2 \cdot 3 z^2(9480\theta^4 + 153912\theta^3 + 212893\theta^2 + 108080\theta + 18816) \\ & + 2^4 \cdot 3^3 z^3(93968\theta^4 + 341544\theta^3 + 319592\theta^2 + 125853\theta + 18242) \\ & - 2^6 \cdot 3^7 z^4(2\theta + 1)^2(2257\theta^2 + 3601\theta + 1942) \\ & + 2^{11} \cdot 3^{11} z^4(2\theta + 1)^2(2\theta + 3)^2 \end{aligned} $
	$A_n = \binom{2n}{n} \sum_k \binom{n}{k} \binom{2n}{2k} \binom{2k}{n-k} \binom{2n-2k}{k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
298	$ \begin{aligned} D = & 3^4\theta^4 - 3^2z(1993\theta^4 + 3218\theta^3 + 2437\theta^2 + 828\theta + 108) \\ & + 2^5z^2(17486\theta^4 + 25184\theta^3 + 12239\theta^2 + 2790\theta + 297) \\ & - 2^8z^3(23620\theta^4 + 34776\theta^3 + 28905\theta^2 + 12447\theta + 2106) \\ & + 2^{15}z^4(2\theta + 1)(340\theta^3 + 618\theta^2 + 455\theta + 129) \\ & - 2^{22}z^5(\theta + 1)^2(2\theta + 1)(2\theta + 3) \end{aligned} $ $A_n = \sum_{k,l} \binom{n}{k} \binom{n}{l} \binom{k+l}{k} \binom{k+l}{n}^2 \binom{2n}{k+l}$
299	$ \begin{aligned} D = & \theta^4 - 6z(144\theta^4 + 36\theta^3 + 47\theta^2 + 29\theta + 6) \\ & + 6^2z^2(8376\theta^4 + 6648\theta^3 + 8157\theta^2 + 3900\theta + 724) \\ & - 6^4z^3(42672\theta^4 + 68616\theta^3 + 81056\theta^2 + 44841\theta + 9964) \\ & + 2^6 \cdot 3^5z^4(374028\theta^4 + 962040\theta^3 + 1262091\theta^2 + 794463\theta + 195335) \\ & - 2^8 \cdot 3^7z^5(633840\theta^4 + 2243328\theta^3 + 3405968\theta^2 + 2385208\theta + 6529129) \\ & + 2^{12} \cdot 3^8z^6(438960\theta^4 + 1884384\theta^3 + 3176664\theta^2 + 2380392\theta + 652943) \\ & - 2^{19} \cdot 3^{10}z^7(5760\theta^4 + 25128\theta^3 + 39548\theta^2 + 26606\theta + 6517) \\ & + 12^{11}z^8(6\theta + 5)^2(6\theta + 7)^2 \end{aligned} $ $ \begin{aligned} A_n = & (-1)^n \binom{2n}{n} \sum_{k=\lfloor n/3 \rfloor}^{\lfloor 2n/3 \rfloor} \binom{n}{k}^3 \binom{2k}{k} \binom{2n-2k}{n-k} \binom{3k}{n} \binom{3n-3k}{n} \binom{2n}{2k}^{-1} \\ & \times (1 + k(-5H_k + 5H_{n-k} + 4H_{2k} - 4H_{2n-2k} \\ & + 3H_{3k} - 3H_{3k-n} + 3H_{2n-3k} - 3H_{3n-3k})) \\ & + 3 \binom{2n}{n} \sum_{k=0}^{\lfloor (n-1)/3 \rfloor} (-1)^k \frac{n-2k}{n-3k} \binom{n}{k}^3 \binom{2k}{k} \binom{2n-2k}{n-k} \\ & \times \binom{3n-3k}{n} \binom{2n}{2k}^{-1} \binom{n}{3k}^{-1} \end{aligned} $
300	$ \begin{aligned} D = & \theta^4 + 2^4z(371\theta^4 + 862\theta^3 + 591\theta^2 + 160\theta + 15) \\ & + 5 \cdot 2^{11}z^2(224\theta^4 + 2069\theta^3 + 3277\theta^2 + 1363\theta + 159) \\ & - 5^2 \cdot 2^{16}z^3(2089\theta^4 + 7500\theta^3 + 5533\theta^2 + 1500\theta + 135) \\ & + 5^3 \cdot 2^{23}z^4(5\theta + 1)(5\theta + 2)(5\theta + 3)(5\theta + 4) \end{aligned} $ $A_n = \binom{2n}{n} \sum_k (-1)^k 4^{n-k} \binom{n}{k} \binom{2n+k}{n} \binom{2k}{n} \binom{4n+2k}{2n+k}$
301	$ \begin{aligned} D = & 11^2\theta^4 - 11z(1517\theta^4 + 3136\theta^3 + 2393\theta^2 + 825\theta + 110) \\ & - z^2(90362\theta^4 + 207620\theta^3 + 202166\theta^2 + 106953\theta + 24266) \\ & - z^3(245714\theta^4 + 507996\theta^3 + 415082\theta^2 + 217437\theta + 53130) \\ & - z^4(407863\theta^4 + 785972\theta^3 + 564786\theta^2 + 183269\theta + 15226) \\ & - z^5(434831\theta^4 + 790148\theta^3 + 728323\theta^2 + 279826\theta + 25160) \\ & - 2^3z^6(36361\theta^4 + 70281\theta^3 + 73343\theta^2 + 37947\theta + 7644) \\ & - 2^4 \cdot 5z^7(1307\theta^4 + 3430\theta^3 + 3877\theta^2 + 2162\theta + 488) \\ & - 2^9 \cdot 5^2z^8(\theta + 1)^4 \end{aligned} $ $A_n = \sum_{k,l} \binom{n}{k} \binom{n}{l} \binom{n+k}{n} \binom{k+l}{k} \binom{2l}{n} \binom{n}{l-k}$

#	differential operator D and coefficients A_n , $n = 0, 1, 2, \dots$
302	$ \begin{aligned} D = & 5^2\theta^4 + 5z(1307\theta^4 + 1798\theta^3 + 1429\theta^2 + 530\theta + 80) \\ & + 2^4z^2(36361\theta^4 + 75163\theta^3 + 80666\theta^2 + 43340\theta + 9120) \\ & + 2^6z^3(434831\theta^4 + 949176\theta^3 + 966865\theta^2 + 545700\theta + 118340) \\ & + 2^{11}z^4(407863\theta^4 + 845480\theta^3 + 654048\theta^2 + 219839\theta + 18634) \\ & + 2^{16}z^5(245714\theta^4 + 474860\theta^3 + 365378\theta^2 + 71595\theta - 11507) \\ & + 2^{21}z^6(90362\theta^4 + 153828\theta^3 + 121478\theta^2 + 35967\theta + 2221) \\ & + 2^{26} \cdot 11z^7(1517\theta^4 + 2932\theta^3 + 2087\theta^2 + 621\theta + 59) \\ & - 2^{31} \cdot 11^2z^8(\theta + 1)^4 \end{aligned} $
	the reflection of #301 at infinity
	a formula for A_n is not known
303	$ \begin{aligned} D = & 13^2\theta^4 - 13z(1505\theta^4 + 2746\theta^3 + 2127\theta^2 + 754\theta + 104) \\ & + 2^2z^2(22961\theta^4 - 2086\theta^3 - 55741\theta^2 - 41574\theta - 9256) \\ & + 2^5z^3(7524\theta^4 + 28098\theta^3 + 16131\theta^2 + 2691\theta - 52) \\ & - 2^7z^4(7241\theta^4 + 6214\theta^3 + 17522\theta^2 + 15423\theta + 4146) \\ & + 2^8z^5(-6087\theta^4 - 1806\theta^3 + 3905\theta^2 + 3796\theta + 1036) \\ & + 2^{10}z^6(553\theta^4 + 4062\theta^3 + 4405\theta^2 + 1752\theta + 220) \\ & + 2^{14}z^7(82\theta^4 + 230\theta^3 + 275\theta^2 + 160\theta + 37) \\ & + 2^{18}z^8(\theta + 1)^4 \end{aligned} $
	$A_n = \sum_{i,j} \binom{n}{k} \binom{n}{l} \binom{k+l}{n} \binom{2k}{n} \binom{2l}{n} \binom{n}{l-k}$
304	$ \begin{aligned} D = & \theta^4 + z(82\theta^4 + 98\theta^3 + 77\theta^2 + 28\theta + 4) \\ & - z^2(-553\theta^4 + 1850\theta^3 + 4463\theta^2 + 2916\theta + 636) \\ & - 2^2z^3(6087\theta^4 + 22542\theta^3 + 27199\theta^2 + 14916\theta + 3136) \\ & - 2^5z^4(7241\theta^4 + 22750\theta^3 + 42326\theta^2 + 29943\theta + 7272) \\ & + 2^7z^5(7524\theta^4 + 1998\theta^3 - 23019\theta^2 - 24627\theta - 7186) \\ & + 2^8z^6(22961\theta^4 + 93930\theta^3 + 88283\theta^2 + 28194\theta + 1624) \\ & - 2^{10} \cdot 13z^7(1505\theta^4 + 3274\theta^3 + 2919\theta^2 + 1282\theta + 236) \\ & + 2^{14} \cdot 13^2z^8(\theta + 1)^4 \end{aligned} $
	the reflection of #303 at infinity
	a formula for A_n is not known

Remark 1. Double sums expressing coefficients A_n , $n = 0, 1, 2, \dots$, occur in cases #28, 102, 193–198, 202, 212, 213 and several others. In all given examples we have

$$A_n = \sum_{i,j} \binom{n}{i}^2 \binom{n}{j}^2 \binom{i+j}{j} B_n(i, j), \quad n = 0, 1, 2, \dots,$$

with a suitable binomial factor $B_n(i, j)$.

Remark 2. In the table there are 25 pairs of differential equations dual at 0 and ∞ , respectively. For the pairs (#22, #118), (#21, #71), (#23, #56) we know a simple formula for the coefficients. Then we have the sporadic pairs (#193, #198), (#210, #211), (#222, #225), (#246, #247), (#55, #277), where a

formula for A_n at ∞ was found by chance. It would be desirable to find formulas for A_n at ∞ also for the remaining 17 cases. An even more interesting question is: What is the geometric meaning of the instanton numbers at ∞ , e.g., for the cases #17, #21, #22, #23, #27, where we know the manifold?

Remark 3. Most of the differential equations #242–290 were found by using Maple’s Zeilberger on

$$“A_n” = \sum_k (n - 2k)C(n, k)$$

(cf. Subsection 2.2), where a binomial expression $C(n, k)$ satisfies $C(n, n - k) = C(n, k)$. Using this symmetry one sees that “ A_n ” = 0, but Maple finds several bona fide differential equations and *usually* the correct coefficients are found by differentiation

$$A_n = \frac{d“A_n”}{dk},$$

which is the case in #244, #246, #247, etc., leading to harmonic sums containing $H_n = \sum_{j=1}^n j^{-1}$. However in some cases (like #242, #245, #259, #260, #261, #262, #264, #274, #279, #281, #282, #299) this is not enough. This will be explained in a forthcoming paper [2].

B Table of powers

We found that in most of the cases $q(z)/z$ was a high power of a power series in $\mathbb{Z}[[z]]$ (and $z(q)/q$ the same power of a power series in $\mathbb{Z}[[q]]$). Similarly, $y_0(z)$ was a (not so high) power of an integral power series. In the following table we present (conjectured) powers r, s in

$$q = z(1 + C_1z + \cdots)^r, \quad y_0 = (1 + D_1z + \cdots)^s;$$

the first column is reserved for numeration of cases.

#	r	s	#	r	s	#	r	s	#	r	s
1	10	4	7*	32	8	39	8	2	73	18	6
2	960	24	7**	32	8	40	32	8	74	6	4
3	64	8	8*	72	2	41	2	1	75	2	1
4	180	6	8**	216	2	42	4	4	76	1	1
5	108	4	9*	288	8	43	32	8	77	2	1
6	32	8	9**	864	8	44	4	12	78	1	8
7	64	8	10*	32	8	45	4	4	79	1	1
8	64	12	10**	32	8	46	18	4	80	2	1
9	576	24	13*	432	8	47	144	8	81	1	1
10	960	24	13**	144	8	48	12	4	82	1	6
11	60	12	14*	48	24	49	72	2	83	8	8
12	96	24	$\widehat{1}$	10	—	50	6	3	84	4	4
13	2880	120	$\widehat{2}$	320	8	51	4	2	85	2	1
14	1728	4	$\widehat{3}$	64	8	52	12	4	86	1	1
15	6	2	$\widehat{4}$	18	—	53	6	3	87	2	1
16	4	4	$\widehat{5}$	24	6	54	1	1	88	2	1
17	6	1	$\widehat{6}$	64	8	55	12	4	89	1	24
18	12	2	$\widehat{7}$	64	8	56	8	8	90	2	1
19	1	2	$\widehat{8}$	72	6	57	2	2	91	1	1
20	3	2	$\widehat{9}$	576	8	58	8	2	92	2	2
21	2	2	$\widehat{10}$	64	8	59	6	4	93	108	1
22	5	1	$\widehat{11}$	24	2	60	2	4	94	1	2
23	4	4	$\widehat{12}$	192	8	61	288	24	95	1	1
24	3	3	$\widehat{13}$	576	24	62	12	4	96	32	1
25	4	2	$\widehat{14}$	192	24	63	12	6	97	192	1
26	2	4	30	64	8	64	24	6	98	24	1
27	14	1	31	64	8	65	24	8	99	2	2
28	8	1	32	78	1	66	24	6	100	12	2
29	2	1	33	4	4	67	288	8	101	30	1
2*	160	16	34	8	1	68	4	4	102	1	1
3*	16	8	35	12	4	69	24	6	103	24	3
4*	54	1	36	24	8	70	3	3	104	1	1
4**	18	4	37	24	2	71	32	8	105	1	1
6*	64	8	38	8	8	72	64	1	106	2	2

#	r	s	#	r	s	#	r	s	#	r	s
107	16	8	150	12	4	193	3	1	236	8	8
108	1	1	151	6	1	194	4	1	237	24	8
109	6	8	152	4	4	195	1	4	238	4	2
110	12	12	153	12	4	196	1	1	239	480	24
111	32	8	154	180	6	197	2	2	240	2	2
112	96	24	155	192	8	198	3	1	241	48	8
113	1	3	156	2	1	199	4	1	242	6	6
114	4	4	157	1	1	200	1	1	243	14	1
115	64	8	158	2	1	201	8	8	244	28	1
116	48	8	159	2	1	202	1	1	245	24	1
117	4	4	160	3	1	203	2	4	246	4	2
118	10	4	161	6	3	204	320	8	247	32	8
119	4	4	162	3	3	205	2	4	248	2	1
120	8	6	163	6	2	206	4	4	249	12	2
121	4	2	164	12	6	207	64	8	250	1	2
122	8	8	165	6	3	208	6	8	251	6	12
123	2	2	166	2880	24	209	2	1	252	2	2
124	1	1	167	3	1	210	12	2	253	4	4
125	1	1	168	3	3	211	96	8	254	192	8
126	2	1	169	12	3	212	2	2	255	12	4
127	72	1	170	6	2	213	2	1	256	16	8
128	2	1	171	24	6	214	2	2	257	64	8
129	1	6	172	9	3	215	4	4	258	96	8
130	12	1	173	4	2	216	6	12	259	6	6
131	1	1	174	1	1	217	2	12	260	4	2
132	10	1	175	3	3	218	6	2	261	4	2
133	12	2	176	4	4	219	2	2	262	4	2
134	9	3	177	4	4	220	48	8	263	32	8
135	12	6	178	3	3	221	4	2	264	96	8
136	36	6	179	3	3	222	6	4	265	96	8
137	4	4	180	24	8	223	6	24	266	6	3
138	6	6	181	6	3	224	2	4	267	18	3
139	12	12	182	2	1	225	192	8	268	60	6
140	12	12	183	4	2	226	2	2	269	1440	24
141	216	2	184	2	1	227	36	12	270	12	2
142	9	3	185	6	1	228	4	12	271	192	8
143	72	6	186	2	1	229	8	2	272	12	6
144	12	6	187	6	9	230	360	4	273	6	6
145	72	3	188	2	2	231	4	4	274	4	2
146	2	2	189	2	1	232	6	4	275	4	2
147	8	4	190	320	1	233	96	8	276	576	8
148	5	1	191	24	1	234	2	16	277	96	8
149	12	12	192	192	1	235	2	16	278	12	12

#	r	s	#	r	s	#	r	s	#	r	s
279	1	1	286	1	2	293	16	4	300	160	8
280	9	3	287	2	1	294	64	8	301	1	1
281	10	1	288	96	72	295	64	8	302	2	4
282	4	2	289	64	8	296	32	8	303	2	2
283	4	6	290	9	3	297	2	48	304	2	2
284	1	1	291	1	3	298	4	2			
285	1	1	292	4	4	299	6	6			

C Superseeker of Calabi–Yau differential equations

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
1	1	2	8		184
2	1	2	104		41
3	3	3	28		28
4	1	4	44		253
5	3	4	44		23
6	1	4	84		84
7	5	4	108		246
8	1	4	284		206
9	1	4	3252	(p)*(p)	3*
10	1	5	100	(a)*(a)	100
11	1	6	104	(k)*(k)	4*
12	1	6	170		245
13	1	6	325	(a)*(f)	160
14	1	9	748	(f)*(f)	165
15	17	10	170		279
16	7	10	295		22
17	7	10	508		235
18	1	10	664	(c)*(c)	103
19	1	10	870		60
20	6	12	140		130
21	2	12	208		46
22	10	12	236		17
23	3	12	644		16
24	1	12	3204	(A)*(a)	45
25	1	13	2650	(b)*(b)	101
26	2	16	208	(a)*(d)	105
27	10	16	304		21
28	1	16	1232	(A)*(d)	36
29	1	16	1744	(d)*(d)	107
30	2	16	2000		42
31	2	16	2106	(a)*(b)	102
32	1	16	3280		56
33	6	18	490		20
34	9	18	3820		199
35	7	18	5676		234
36	20	20	100		205
37	4	20	1680		281
38	5	20	1820		18
39	2	20	2036		244
40	1	20	5924	(a)*(e)	114, 150

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
41	1	20	8220		25
42	1	21	15894		15
43	3	24	1552		188
44	7	26	55644		297
45	3	27	14201		216
46	1	27	18089	$(B)*(\iota)$	70
47	1	28	1036	$(b)*(e)$	121
48	5	28	1268		262
49	6	28	1820		27
50	3	28	3892		119
51	1	32	1440	$(l)*(l), (A)*(\beta)$	$10^{**}, 40$
52	1	32	7584		201
53	1	32	26016	$(C)*(d), (A)*(\theta)$	3, 30, 31, 72
54	3	32	38880	$(A)*(d)$	111
55	3	33	3600	$(b)*(c)$	113
56	7	35	2184		28
57	9	36	556		183
58	10	36	1284		266
59	3	36	3020	$(d)*(f)$	163
60	3	36	3284	$(A)*(f)$	133
61	6	36	3648		185
62	1	36	8076	$(B)*(e)$	110
63	1	36	41421	$(B)*(b)$	24
64	1	36	128217204	$(B)*(j)$	
65	1	37	15270	$(g)*(g)$	144
66	13	39	1621		197
67	8	40	5128		304
68	2	40	26376		293
69	6	42	2542	$(a)*(c)$	104
70	44	44	308		182
71	5	44	2480		249
72	1	44	22500	$(C)*(g)$	139
73	4	48	1424	$(b)*(d)$	106
74	1	48	2864	$(B)*(\beta), (A)*(h)$	$8^{**}, 49, 141$
75	1	48	9104		296
76	3	48	11056	$(A)*(c)$	58
77	2	48	11664	$(B)*(d)$	48
78	1	48	32368	$(d)*(e), (A)*(\varepsilon)$	122
79	1	48	73328	$(C)*(d)$	38
80	19	49	1761		186
81	21	51	5095		217
82	20	52	1356		203

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
83	3	52	52284		117
84	1	52	220220		68
85	1	54	40552		50
86	3	54	64744		223
87	1	55	11655		118
88	24	56	3552		248
89	5	59	22503		224
90	3	60	1684	(A)*(g)	137
91	3	60	28820		255
92	1	60	134292		5, 90, 91, 93, 157
93	1	60	307860	(B)*(i)	
94	1	63	96866	(f)*(h)	172
95	1	64	23360		116
96	1	64	246848	(C)*(c)	69
97	2	66	59386	(B)*(δ)	151
98	4	68	95246	(c)*(g)	175
99	24	72	3496		286
100	12	72	3900	(a)*(f)	160
101	4	72	20708	(B)*(f)	134
102	5	76	10500		270
103	5	76	24836		261
104	3	76	144196		55
105	1	76	415420	(a)*(i), (C)*(δ)	152
106	28	80	2912		212
107	23	80	4655		19
108	1	80	104976		233
109	1	80	174096		83
110	1	80	249872		236
111	11	84	9052		198
112	6	84	20848		29
113	2	84	83412		243
114	3	84	113304		291
115	1	90	151648		73
116	1	92	585396	(C)*(b)	51
117	1	96	12064	(C)*(β)	7*, 43
118	3	96	26208	(c)*(e), (A)*(α)	39, 120
119	1	100	126580	(b)*(i), (C)*(η)	
120	6	108	910		242
121	12	108	968	(c)*(f)	162
122	6	108	3136		259
123	12	108	4916	(b)*(f)	161
124	4	108	10472	(B)*(g)	138

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
125	12	108	12580		251
126	6	108	19598	$(f)*(g)$	178
127	3	108	62596	$(e)*(f), (A)*(\zeta)$	164
128	2	108	81104	$(c)*(h)$	169
129	3	108	206716	$(C)*(f)$	135
130	1	108	49457556	$(D)*(g)$	140
131	5	109	16777		302
132	1	112	378800		71
133	19	113	8515		202
134	5	116	186172		275
135	1	117	713814		4
136	1	117	844872		280
137	17	126	11700		194
138	1	128	382592		220
139	1	128	800384		256
140	1	128	5282176	$(C)*(\alpha)$	37
141	7	129	41441		193
142	12	132	9736	$(a)*(g)$	173
143	4	132	52204		32
144	4	132	118772	$(a)*(h)$	167
145	7	138	42984		218
146	14	140	24136		26
147	12	144	7312	$(c)*(d)$	123
148	6	144	30896	$(d)*(g)$	176
149	15	147	6032		274
150	13	151	26293		303
151	1	160	9310		19
152	2	160	539680	$(e)*(e), (c)*(i), (C)*(\theta)$	$\hat{3}, 6^*, 115$
153	1	160	1956896		6, 75, 96, 146
154	1	160	5870688		76
155	3	162	197216		299
156	61	163	4795		124
157	4	180	28320	$(b)*(h)$	168
158	4	180	110940	$(B)*(h)$	142
159	1	180	21847076	$(D)*(f)$	136
160	15	186	20300		226
161	5	188	450516		260
162	47	189	9277		196
163	12	192	156	$(b)*(g)$	174
164	11	193	48570		301
165	1	196	2993772		33
166	3	204	18628		228

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
167	1	208	1218192		237
168	1	208	1863312	$(d)*(i), (C)*(\varepsilon)$	
169	9	209	97622		298
170	3	220	267636		215
171	3	228	278988	$(e)*(g), (A)*(\gamma)$	44, 177
172	13	231	38037		240
173	5	232	122168		252
174	9	234	103520		214
175	34	236	22848		213
176	56	240	6944		59
177	8	240	117056		74
178	1	240	19105840	$(D)*(d)$	65
179	38	241	17458		284
180	1	243	513936		278
181	20	252	56064		273
182	4	252	387464	$(B)*(\zeta)$	
183	1	252	1162036		154
184	3	252	1522388		35
185	2	270	835370	$(g)*(h)$	53, 179
186	7	274	281388		208
187	20	276	116324		222
188	29	285	40626		195
189	1	288	2339616	$(e)*(h)$	$\widehat{5}, 98, 171$
190	1	288	96055968	$(D)*(e), (A)*(\kappa)$	112
191	1	291	7935104		230
192	23	308	70799		250
193	1	320	19748928		241
194	2	324	1502052		290
195	1	324	10792428		11, 95
196	4	336	595280	$(d)*(h), (B)*(\varepsilon)$	170
197	21	361	120472		287
198	5	364	6324580		282
199	1	372	71562236	$(D)*(a)$	62
200	5	379	1364199		232
201	1	384	164736	$(m)*(m)$	13*, 47
202	3	384	1546624	$(c)*(i), (C)*(\alpha)$	37
203	14	420	159040		189
204	4	428	485244		187
205	1	432	78259376	$(D)*(c)$	64
206	5	444	1501908		210
207	1	444	19050964		238
208	3	460	894404		231

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
209	3	468	3687996	$(f)*(i), (C)*(\zeta)$	
210	5	468	59427420		272
211	17	478	285760		209
212	1	480	4215904		258
213	1	480	16034720	$(D)*(\beta)$	$9^*, 9^{**}, 67$
214	5	492	872164		221
215	1	492	136094428	$(a)*(j), (D)*(\delta)$	153
216	15	498	360988		219
217	1	522	9879192	$(h)*(h)$	$\widehat{4}, 145$
218	3	564	2422620		283
219	1	575	63441275		1, 79, 87, 128
220	1	608	22293216		247
221	1	612	51318900	$(b)*(j), (D)*(\eta)$	
222	1	624	43406256		180
223	13	647	942613		99
224	72	684	398428		200
225	1	684	195638820	$(D)*(b)$	63
226	1	736	26911072	$(e)*(i)$	$\widehat{6}, 77, 78, 97$
227	5	828	4270932		268
228	1	864	147560800	$(c)*(j), (D)*(\alpha)$	66
229	1	900	8364884		227
230	1	928	170809536		10, 54
231	1	992	63721056		257
232	1	1008	607849200	$(D)*(h), (B)*(\kappa)$	143
233	3	1020	15174100	$(g)*(i), (C)*(\gamma)$	52
234	3	1056	15001120		229
235	1	1056	138459552		265
236	1	1116	349462868	$(f)*(j), (D)*(\zeta)$	
237	1	1248	683015008		14, 85, 86, 156
238	1	1312	58156708		263
239	1	1344	109320512	$(h)*(i)$	$\widehat{11}, 94$
240	7	1434	18676572		109
241	1	1488	517984144	$(d)*(j), (D)*(\varepsilon)$	
242	1	1584	171534960		239
243	1	1616	283183120		300
244	1	1818	467810538		267
245	4	2300	253765100		148
246	1	2400	2956977632		211
247	1	2450	623291900		$\widehat{1}, 131$
248	1	2484	1327731388	$(g)*(j), (D)*(\gamma)$	149
249	1	2592	81451104	$(C)*(j)$	
250	1	2628	3966805740		8, 92, 125, 158

	N_0	$ N_1 $	$ N_3 $	Hadamard product	# in Table A
251	44	3180	6378752		285
252	1	3488	1142687008	(i)*(i)	$\widehat{10}$, 155
253	1	3616	264403872		288
254	1	3936	10892932064	(D)*(i)	
255	1	4192	2124587232		277
256	3	4300	1701817028		292
257	2	4900	1246583800		80, 81
258	1	5408	4296119968		295
259	1	5408	22147077792		254
260	1	5472	6444589536	(e)*(j)	$\widehat{14}$, 88, 89
261	1	7776	66942277344		12
262	1	8224	15542388128		289
263	1	10080	24400330080	(h)*(j)	$\widehat{8}$, 82, 126, 127, 129
264	1	10912	71557619232		271
265	1	14752	711860273440		7, 147
266	1	26400	230398034080	(i)*(j)	$\widehat{12}$
267	1	37216	464865119712		264
268	1	41184	5124430612320	(D)*(j), (D)*(κ)	61
269	1	57760	3869123234080		$\widehat{7}$
270	1	67104	28583248229280		13, 57, 108
271	1	70944	3707752060576		207
272	1	80416	15561562691488		294
273	1	93984	25265152551072		225
274	1	188832	101990911789344		276
275	1	201888	40177844666400	(j)*(j)	$\widehat{13}$, 166
276	1	231200	1700894366474400		2, 159
277	1	549216	5134247872650720		269
278	1	678816	69080128815414048		9
279	1	791200	4288711075194400		$\widehat{2}$
280	1	2710944	302270555492914464		$\widehat{9}$

Comments. We list $|N_1|$ and $|N_3|$. The reason for not using N_2 is that it is not invariant under the transformation $q \mapsto -q$. There are many differential equations in the table that are just transformations (by [1], Proposition 8)

$$y_0(z) \mapsto \frac{1}{1-pz} y_0\left(\left(\frac{z}{1-pz}\right)^r\right)$$

which transforms the Yukawa coupling $K(q) \mapsto K(q^r)$. These differential equations should not be in the table but we did not know this transformation when we found them. In the Superseeker table we identify $K(q)$ and $K(q^r)$ as the instanton numbers are just thinned out.

References

- [1] G. Almkvist and W. Zudilin, *Differential equations, mirror maps and zeta values*, Calabi–Yau Varieties and Mirror Symmetry, The BIRS workshop volume (06–11 December 2003) (J. Lewis, S.-T. Yau and N. Yui, eds.), Amer. Math. Soc. & International Press, Providence, RI, 2005; `math.NT/0402386` (2004).
- [2] G. Almkvist and C. Krattenthaler, *Some harmonic sums related to Calabi–Yau differential equations*, in preparation.
- [3] V. V. Batyrev and D. van Straten, *Generalized hypergeometric functions and rational curves on Calabi–Yau complete intersections in toric varieties*, Commun. Math. Phys. **168** (1995), 493–533; `alg-geom/9310001` (1993).
- [4] V. V. Batyrev, I. Ciocan-Fontanine, B. Kim, and D. van Straten, *Conifold transitions and mirror symmetry for Calabi–Yau complete intersections in Grassmannians*, Nuclear Phys. B **514** (1998), no. 3, 640–666; `alg-geom/9710022` (1997).
- [5] C. van Enkevort and D. van Straten, *Monodromy calculations of fourth order equations of Calabi–Yau type*, Calabi–Yau Varieties and Mirror Symmetry, The BIRS workshop volume (06–11 December 2003) (J. Lewis, S.-T. Yau and N. Yui, eds.), Amer. Math. Soc. & International Press, Providence, RI, 2005; `math.AG/0412539` (2004).
- [6] V. Golyshev, *Modularity of equations $D3$ and the Iskovskikh classification*, Modularity of the $D3$ equations and the Iskovskikh classification. Dokl. Akad. Nauk **396** (2004), no. 6, 733–739; `math.AG/0405039` (2004).
- [7] A. Klemm and S. Theisen, *Considerations of one-modulus Calabi–Yau compactifications: Picard–Fuchs equations, Kähler potentials and mirror maps*, Nuclear Phys. B **389** (1993), no. 1, 153–180; `hep-th/9205041` (1992).
- [8] A. Klemm and S. Theisen, *Mirror maps and instanton sums for complete intersections in weighted projective space*, Modern Phys. Lett. A **9** (1994), no. 20, 1807–1817; `hep-th/9304034` (1993).
- [9] C. Krattenthaler, A personal communication (2005).
- [10] A. Libgober and J. Teitelbaum, *Lines in Calabi–Yau complete intersections, mirror symmetry, and Picard–Fuchs equations*, Intern. Math. Research Notices **1** (1993), no. 1, 29–39; `alg-geom/9301001` (1993).
- [11] D. R. Morrison, *Picard–Fuchs equations and mirror maps for hypersurfaces*, Essays on mirror manifolds, Hong Kong, International Press, 1992, pp. 241–264; `hep-th/9111025` (1991).
- [12] E. A. Rødland, *The Pfaffian Calabi–Yau, its mirror, and their link to the Grassmannian $G(2, 7)$* , Compositio Math. **122** (2000), no. 2, 135–149; `math.AG/9801092` (1998).
- [13] E. N. Tjøtta, *Quantum cohomology of a Pfaffian Calabi–Yau variety: verifying mirror symmetry predictions*, `math.AG/9906119` (1999).

- [14] H. A. Verrill, *Sums of squares of binomial coefficients, with applications to Picard–Fuchs equations*, **math.CO/0407327** (2004).
- [15] H. S. Wilf and D. Zeilberger, *An algorithmic proof theory for hypergeometric (ordinary and “q”) multisum/integral identities*, *Invent. Math.* **108** (1992), no. 3, 575–633.
- [16] D. Zagier, *Integral solutions of Apéry-like recurrence equations*, Manuscript (2003).

MATEMATIKCENTRUM
 LUNDS UNIVERSITET
 MATEMATIK MNF, BOX 118
 SE-221 00 LUND, SWEDEN
E-mail address: `gert@maths.lth.se`

FACHBEREICH MATHEMATIK 17
 AG ALGEBRAISCHE GEOMETRIE
 JOHANNES GUTENBERG-UNIVERSITÄT
 D-55099 MAINZ, GERMANY
E-mail address: `enckevor@mathematik.uni-mainz.de`
E-mail address: `straten@mathematik.uni-mainz.de`

DEPARTMENT OF MECHANICS AND MATHEMATICS
 MOSCOW LOMONOSOV STATE UNIVERSITY
 VOROBIOVY GORY, GSP-2
 119992 MOSCOW, RUSSIA
E-mail address: `wadim@ips.ras.ru`